


NMDA

Cat. No.:	HY-17551		
CAS No.:	6384-92-5		
Molecular Formula:	C ₅ H ₉ NO ₄		
Molecular Weight:	147.13		
Target:	iGluR; Endogenous Metabolite		
Pathway:	Membrane Transporter/Ion Channel; Neuronal Signaling; Metabolic Enzyme/Protease		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro

H₂O : 33.33 mg/mL (226.53 mM; Need ultrasonic)
 DMSO : 10 mg/mL (67.97 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent		Mass		
	Concentration		1 mg	5 mg	10 mg
	1 mM		6.7967 mL	33.9836 mL	67.9671 mL
	5 mM		1.3593 mL	6.7967 mL	13.5934 mL
	10 mM		0.6797 mL	3.3984 mL	6.7967 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: PBS
Solubility: 36.67 mg/mL (249.24 mM); Clear solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 1 mg/mL (6.80 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 1 mg/mL (6.80 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 1 mg/mL (6.80 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

NMDA is a specific agonist for NMDA receptor mimicking the action of glutamate, the neurotransmitter which normally acts at that receptor.

IC₅₀ & Target

Human Endogenous	NMDA Receptor
------------------	---------------

	Metabolite
In Vitro	NMDA exerts a significant augmentation of the adrenal binding independently of the incubation temperature in a concentration-dependent manner ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	NMDA (0.2 nmol) shows significant effects on MF, IF, IL, and EL, respectively, decreasing the mount and intromission frequencies, and shortening the intromission and ejaculation latencies. NMDA and AP-5 significantly, respectively, facilitates and inhibits the ejaculatory behavior during the copulation testing 30 min. Bilateral microinjection of NMDA into PVN significantly increases the baseline LSNA, the peaking increment of LSNA occurred within 5 min from the time of NMDA microinjected into PVN ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Kinase Assay ^[2]	Adrenal membranous homogenate suspensions are incubated with 10 nM [³ H]Glu in 500/μl 50 mM Tris-acetate buffer (pH 7.4) at 2°C or 30°C in the presence and absence of various compounds. Incubation is terminated by the addition of 3 mL ice-cold buffer and subsequent filtration through a Whatman GF/B glass filter under a constant vacuum of 15 mm Hg. After washing the filter 4 times with 3 mL icecold buffer, the radioactivity trapped on the filter is measured by a liquid scintillation spectrometer using 5 mL modified Triton-toluene scintillant at a counting efficiency of 40-42%. The radioactivity found in the presence of 1 mM non-radioactive Glu is subtracted from each experimental value to obtain the specific binding of [³ H]Glu in accordance with the γ-aminobutyric acid (GABA) receptor binding assay system. The kinetic parameters of [³ H]Glu binding, K _d and B _{max} , are calculated by Scatchard analysis of the specific binding using a personal computer with a programme for non-linear regression analysis developed in our own laboratory. MCE has not independently confirmed the accuracy of these methods. They are for reference only.
Animal Administration ^[1]	Thirty male rats are paired with different receptive females for a total of three times (once every 3 days) a week prior to the experiment, only the males that ejaculated at least three times during this period are included. After selecting the male rats with normal ejaculatory ability. Saline (100 nL), NMDA (0.20 nmol in 100 nL saline), and AP-5 (10.0 nmol in 100 nL saline) are administration into the bilateral PVN of each male rat in random order. After 5 min, the behavioral testing is performed and recorded as described above. Copulatory behaviors occur once a week and the entire experiment lasted 4 weeks. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Biomark Res. 2023 Aug 8;11(1):74.
- Mol Neurobiol. 2023 Dec 8.
- Mol Neurobiol. 2022 Dec 17.
- Mol Neurobiol. 2022 Oct 13.
- Heliyon. 2023 May 29, e16631.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Xia JD, et al. Centrally mediated ejaculatory response via sympathetic outflow in rats: role of N-methyl-D-aspartic acid receptors in paraventricular nucleus. *Andrology*. 2016 Nov 16.

[2]. Yoneda Y, et al. Enhancement of [3H]glutamate binding by N-methyl-D-aspartic acid in rat adrenal. Brain Res. 1987 Mar 17;406(1-2):24-31.

[3]. Jiang L, et al. Decrease of growth and differentiation factor 10 contributes to neuropathic pain through N-methyl-D-aspartate receptor activation. Neuroreport. 2017 May 24;28(8):444-450.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA