


CFDA-SE

Cat. No.:	HY-D0938
CAS No.:	150347-59-4
Molecular Formula:	C ₅₈ H ₃₈ N ₂ O ₂₂
Molecular Weight:	557.46
Storage:	-20°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (89.69 mM; Need ultrasonic)						
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg	
				1 mM	1.7939 mL	8.9693 mL	17.9385 mL
				5 mM	0.3588 mL	1.7939 mL	3.5877 mL
				10 mM	0.1794 mL	0.8969 mL	1.7939 mL
Please refer to the solubility information to select the appropriate solvent.							
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (3.73 mM); Clear solution 2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (3.73 mM); Clear solution						

BIOLOGICAL ACTIVITY

Description	<p>CFDA-SE is a fluorescent dye that can penetrate the cell membrane. It can react with the free amine group in the cytoskeleton protein inside the cell, and finally form a protein complex with fluorescence. After entering the cell, CFDA-SE locates in the cell membrane, cytoplasm and nucleus, and the fluorescence staining is strongest in the nucleus^[1]. CFDA-SE dye can be uniformly inherited by the cells with cell division and proliferation, and its attenuation is proportional to the number of cell divisions. This phenomenon can be detected and analyzed by flow cytometry under the excitation light of 488 nm, and can be used to detect the proliferation of cells^[1].</p>
In Vitro	<p>Preparation of CFDA-SE working solution</p> <p>1.1 Preparation of the stock solution</p> <p>Dissolve 1 mg of CFDA-SE in 0.1794 mL of DMSO to obtain 10 mM of CFSE.</p> <p>Note: It is recommended to store the stock solution at -20 °C or -80 °C away from light and avoid repetitive freeze-thaw cycles.</p>

1.2 Preparation of CFDA-SE working solution

Dilute the stock solution in serum-free cell culture medium or PBS to obtain 5-10 μM of CFDA-SE working solution.

Note: Please adjust the concentration of CFDA-SE working solution according to the actual situation.

Cell staining

2.1 For suspension cells: Centrifuge at 1000 g at 4 \times for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time.

For adherent cells: Discard the cell culture medium, and add trypsin to dissociate cells to make a single-cell suspension. Centrifuge at 1000 g at 4 \times for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time.

2.2 Add 1 mL of CFDA-SE working solution, and then incubate at room temperature for 30 minutes.

2.3 Centrifuge at 400 g at 4 \times for 3-4 minutes and then discard the supernatant.

2.4 Wash twice with PBS, 5 minutes each time.

2.5 Resuspend cells with serum-free cell culture medium or PBS, and then detect by fluorescence microscope or flow cytometer.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[2]

Single-cell suspensions of splenocytes (or PBMC) are stained with 1 μM CFSE in PBS for 9 min at 37 $^{\circ}\text{C}$, combined with 20 mL of 10% FBS RPMI-1640 medium at RT for 2 min, centrifuged, washed, and counted. 2×10^6 CFSE-labeling splenocytes are added to the top of the slices in 24-well membrane culture insert^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cell Res. 2023 Jun;33(6):464-478.
- Adv Mater. 2023 Nov;35(46):e2303614.
- Nature Cancer. 222-234 (2020).
- Adv Funct Mater. 2023 May 14.
- Autophagy. 2021 Nov;17(11):3592-3606.

See more customer validations on www.MedChemExpress.com

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA