

IL-17A Protein, Mouse (Biotinylated, HEK293, His-Avi)

Cat. No.:	HY-P78158
Synonyms:	CTLA-8; IL17; IL17A; IL-17CTLA-8; interleukin 17A
Species:	Mouse
Source:	HEK293
Accession:	Q62386 (A26-A158)
Gene ID:	16171
Molecular Weight:	20-28 kDa

PROPERTIES

Biological Activity	Immobilized Mouse IL-17R alpha at 2 µg/mL (100µL/Well) on the plate. Dose response curve for Biotinylated Mouse IL-17A His with the EC ₅₀ < 20 ng/mL determined by ELISA.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

Interleukin-17A (IL-17A), also known as CTLA-8, belongs to the IL-17 cytokine family. IL-17A is expressed in memory Th17 cells and is a product of memory CD4⁺ T cells. IL-17A is also produced by a wide variety of immune cells, including CD8⁺ T cells, γδT cells, natural killer T (NKT) cells, monocytes, and neutrophils^{[1][2][3]}.

The mouse IL-17A shares 63.23% amino acid sequence identity with human and 87.33% identity with rat.

IL-17A plays a critical role in host defense mechanisms against many bacterial and fungal pathogens as well as allergic and autoimmune responses. IL-17A induces the production of antimicrobial peptides (defensins and S100 proteins), cytokines (IL-6, G-CSF, and GM-CSF), chemokines (CXCL1, CXCL5, IL-8, CCL2, and CCL7), and matrix metalloproteinases (MMP1, MMP3, and MMP13). IL-17A is detrimental in viral infection through promoting neutrophilic inflammation. IL-17A is a homodimeric cytokine and shares similar biological activities with IL-17F. IL-17A binds to IL-17RA with high affinity, and IL-17RA is required for the biological activity of IL-17A. In tumorigenesis, IL-17A recruits myeloid derived suppressor cells (MDSCs) to dampen anti-tumor immunity. IL-17A also enhances tumor growth in vivo through the induction of IL-6^{[1][2]}.

IL-17A can be used for the research of autoimmune diseases, infection and cancer^{[1][4]}.

REFERENCES

- [1]. Chen K, et al. Interleukin-17A (IL17A). *Gene*. 2017 May 30;614:8-14.
 - [2]. Iwakura Y, et al. The roles of IL-17A in inflammatory immune responses and host defense against pathogens. *Immunol Rev*. 2008 Dec;226:57-79.
 - [3]. Cua DJ, et al. Innate IL-17-producing cells: the sentinels of the immune system. *Nat Rev Immunol*. 2010 Jul;10(7):479-89.
 - [4]. Wright JF, et al. The human IL-17F/IL-17A heterodimeric cytokine signals through the IL-17RA/IL-17RC receptor complex. *J Immunol*. 2008 Aug 15;181(4):2799-805.
 - [5]. Pelidou SH, et al. Enhancement of acute phase and inhibition of chronic phase of experimental autoimmune neuritis in Lewis rats by intranasal administration of recombinant mouse interleukin 17: potential immunoregulatory role. *Exp Neurol*. 2000 May;163(1):165-72.
 - [6]. Liu Y, et al. IL-17A and TNF- α exert synergistic effects on expression of CXCL5 by alveolar type II cells in vivo and in vitro. *J Immunol*. 2011 Mar 1;186(5):3197-205.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA