



www.MedChemExpress.com

Inhibitors, Screening Libraries, Proteins

ADC Linkers











Antibody-drug conjugates linkers











Antibody-drug conjugates (ADCs) consist of a desirable monoclonal antibody, an active cytotoxic drug and an appropriate linker. An appropriate linker between the antibody and the cytotoxic drug provides a specific bridge, and thus helps the antibody to selectively deliver the cytotoxic drug to tumor cells and accurately releases the cytotoxic drug at tumor sites. In addition to conjugation, the linkers maintain ADCs' stability during the preparation and storage stages of the ADCs and during the systemic circulation period.











The ADCs currently undergoing clinical evaluation contain linkers are mostly classified into two categories: cleavable and noncleavable. Cleavable linkers rely on processes inside the cell to liberate the toxin, such as reduction in the cytoplasm, exposure to acidic conditions in the lysosome, or cleavage by specific proteases within the cell. Noncleavable linkers require proteolytic degradation of the antibody portion of the ADC for release of the cytotoxic molecule, which will retain the linker and the amino acid by which it was attached to the antibody.









The selection of linker is target dependent, based on the knowledge of the internalization and degradation of the antibody-target antigen complex, and a preclinical in vitro and in vivo activity comparison of conjugates. Moreover, the choice of a linker is also influenced by which cytotoxin is used, as each molecule has different chemical constraints, and frequently the drug structure lends itself to a specific linker.


ADC Linkers Chemicals

<p>(2-pyridyldithio)-PEG1-hydrazine</p> <p>Cat. No.: HY-136135</p>	<p>(2-pyridyldithio)-PEG4 acid</p> <p>Cat. No.: HY-135964</p>
<p>(2-pyridyldithio)-PEG1-hydrazine is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 50 mg, 100 mg</p>	<p>(2-pyridyldithio)-PEG4 acid is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>(2-Pyridyldithio)-PEG6 acid</p> <p>Cat. No.: HY-132086</p>	<p>(Ac)Phe-Lys(Alloc)-PABC-PNP</p> <p>Cat. No.: HY-20560</p>
<p>2-Pyridyldithio-PEG6 acid is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>(Ac)Phe-Lys(Alloc)-PABC-PNP is a useful cleavable chemical linker in antibody drug conjugates.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>(R)-8-Azido-2-(Fmoc-amino)octanoic acid</p> <p>Cat. No.: HY-131082</p>	<p>(R)-Azetidine-2-carboxylic acid</p> <p>Cat. No.: HY-W017755</p>
<p>(R)-8-Azido-2-(Fmoc-amino)octanoic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>(R)-Azetidine-2-carboxylic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). (R)-Azetidine-2-carboxylic acid is also a alkyl chain-based PROTAC linker that can be.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>1,6-Bis(mesyloxy)hexane</p> <p>Cat. No.: HY-138327</p>	<p>1-Boc-azetidine-3-carboxylic acid</p> <p>Cat. No.: HY-40141</p>
<p>16-Bismesyloxyhexane is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>1-Boc-azetidine-3-carboxylic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-Boc-azetidine-3-carboxylic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs^{2/sup}.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg, 1 g</p>
<p>1-Boc-azetidine-3-yl-methanol</p> <p>Cat. No.: HY-40152</p>	<p>1-Cbz-3-Hydroxyazetidine</p> <p>Cat. No.: HY-77475</p>
<p>1-Boc-azetidine-3-yl-methanol is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-Boc-azetidine-3-yl-methanol is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs^{2/sup}.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg, 1 g</p>	<p>1-Cbz-3-Hydroxyazetidine is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-Cbz-3-Hydroxyazetidine is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g, 5 g</p>









<p>1-Cbz-azetidine-3-carboxylic acid</p> <p>Cat. No.: HY-W004868</p> <p>1-Cbz-azetidine-3-carboxylic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-Cbz-azetidine-3-carboxylic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs^{+/sup}.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg</p> 	<p>1-N-Boc-3-hydroxyazetidine</p> <p>Cat. No.: HY-40142</p> <p>1-N-Boc-3-hydroxyazetidine is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-N-Boc-3-hydroxyazetidine is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg, 1 g</p> 
<p>2-Aminoethyl-mono-amide-DOTA-tris(tBu ester)</p> <p>Cat. No.: HY-100138</p> <p>2-Aminoethyl-mono-amide-DOTA-tris(tBu ester) is a macrocycle DOTA derivative for tumor pretargeting.</p> <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 10 mg, 25 mg, 50 mg, 100 mg</p> 	<p>2-Hydroxyethyl disulfide mono-tosylate</p> <p>Cat. No.: HY-140125</p> <p>2-Hydroxyethyl disulfide mono-tosylate is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>20-(tert-Butoxy)-20-oxoicosanoic acid</p> <p>Cat. No.: HY-W034597</p> <p>20-(tert-Butoxy)-20-oxoicosanoic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 20-(tert-Butoxy)-20-oxoicosanoic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs <su.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 250 mg, 500 mg</p> 	<p>22-(tert-Butoxy)-22-oxodocosanoic acid</p> <p>Cat. No.: HY-W046348</p> <p>22-(tert-Butoxy)-22-oxodocosanoic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 22-(tert-Butoxy)-22-oxodocosanoic acid is also a alkyl chain-based PROTAC linker that can be used in t.</p> <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p> 
<p>4-Methyl-4-(methylidisulfanyl)pentanoic acid</p> <p>Cat. No.: HY-133408</p> <p>4-Methyl-4-(methylidisulfanyl)pentanoic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>4-Methyl-4-(pyridin-2-ylidisulfanyl)pentanoic acid</p> <p>Cat. No.: HY-133409</p> <p>4-Methyl-4-(pyridin-2-ylidisulfanyl)pentanoic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>4-N3Pfp-NHS ester</p> <p>Cat. No.: HY-126525</p> <p>4-N3Pfp-NHS ester is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>4-Succinimidyl-oxycarbonyl-α-(2-pyridyldithio)toluene</p> <p>Cat. No.: HY-133538</p> <p>4-Succinimidyl-oxycarbonyl-α-(2-pyridyldithio)toluene is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 

<p>5-Maleimidovaleric acid</p> <p>Cat. No.: HY-140987</p> <p>5-Maleimidovaleric acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>6-Azido-hexylamine</p> <p>Cat. No.: HY-138387</p> <p>6-Azido-hexylamine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>6-Maleimidohexanoic acid N-hydroxysuccinimide ester (EMCS)</p> <p>Cat. No.: HY-78961</p> <p>6-Maleimidohexanoic acid N-hydroxysuccinimide ester (EMCS) is a heterobifunctional cross-linking reagent. EMCS is used as a unique and useful reagent for preparation of hapten conjugate and enzyme immunoconjugates.</p>  <p>Purity: 99.62% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 500 mg</p>	<p>6-O-2-Propyn-1-yl-D-galactose</p> <p>Cat. No.: HY-128930</p> <p>6-O-2-Propyn-1-yl-D-galactose is a noncleavable glycolinker for the functionalization of cytotoxic drugs and applications in antibody-drug conjugation.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>6-Oxohexanoic acid</p> <p>Cat. No.: HY-141595</p> <p>6-Oxohexanoic acid is a non-cleavable modified MMAF-C5-COOH linker and can be used in the synthesis of modified MMAF-C5-COOH, a drug-linker conjugate for ADC.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>AcBut (4-(4-Acetyl-phenoxy)-butyric acid)</p> <p>Cat. No.: HY-132261</p> <p>AcBut is a cleavable Ozogamicin linker used in the synthesis of Ozogamicin, a drug-linker conjugate for ADC.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 50 mg, 100 mg</p>
<p>Acid-C3-SSPy</p> <p>Cat. No.: HY-141597</p> <p>Acid-C3-SSPy is a cleavable DBA-DM4 linker used in the synthesis of DBA-DM4 (HY-128960), a drug-linker conjugate for ADC.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Acid-PEG1-bis-PEG3-BCN</p> <p>Cat. No.: HY-136088</p> <p>Acid-PEG1-bis-PEG3-BCN is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Acid-PEG2-SS-PEG2-acid</p> <p>Cat. No.: HY-140112</p> <p>Acid-PEG2-SS-PEG2-acid is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 100 mg, 250 mg, 500 mg</p>	<p>Acid-PEG3-SS-PEG3-acid</p> <p>Cat. No.: HY-140113</p> <p>Acid-PEG3-SS-PEG3-acid is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>




<p>Acid-propionylamino-Val-Cit-OH</p> <p>Cat. No.: HY-130930</p> <p>Acid-propionylamino-Val-Cit-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>AEEA-AEEA</p> <p>Cat. No.: HY-W125504</p> <p>AEEA-AEEA is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). AEEA-AEEA is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Ala-Ala-Asn-PAB</p> <p>Cat. No.: HY-129360</p> <p>Ala-Ala-Asn-PAB is a peptide cleavable ADC linker for antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g</p>	<p>Ala-CO-amide-C4-Boc</p> <p>Cat. No.: HY-145367</p> <p>Ala-CO-amide-C4-Boc is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Ald-CH2-PEG3-azide</p> <p>Cat. No.: HY-130144</p> <p>Ald-CH2-PEG3-azide is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Ald-CH2-PEG3-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Ald-CH2-PEG5-azide</p> <p>Cat. No.: HY-140634</p> <p>Ald-CH2-PEG5-azide is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Ald-PEG23-SPDP</p> <p>Cat. No.: HY-136309</p> <p>Ald-PEG23-SPDP is a cleavable 23 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>ALD-PEG4-OPFP</p> <p>Cat. No.: HY-136127</p> <p>ALD-PEG4-OPFP is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Ald-Ph-amido-C2-nitrate</p> <p>Cat. No.: HY-130096</p> <p>Ald-Ph-amido-C2-nitrate (Example XXIVb) is a thiazolidine derivative, used as a noncleavable ADC linker.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Ald-Ph-amido-PEG1-C2-NHS ester</p> <p>Cat. No.: HY-130106</p> <p>Ald-Ph-amido-PEG1-C2-NHS ester is a noncleavable 1-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>

<p>Ald-Ph-amido-PEG1-C2-Pfp ester</p> <p>Cat. No.: HY-130105</p>	<p>Ald-Ph-amido-PEG11-C2-NH2</p> <p>Cat. No.: HY-133546</p>
<p>Ald-Ph-amido-PEG1-C2-Pfp ester is a noncleavable 1-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg</p>	<p>Ald-Ph-amido-PEG11-C2-NH2 is a non-cleavable 11 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Ald-Ph-amido-PEG11-NH-Boc</p> <p>Cat. No.: HY-133572</p>	<p>Ald-Ph-amido-PEG2</p> <p>Cat. No.: HY-130099</p>
<p>Ald-Ph-amido-PEG11-NH-Boc is a non-cleavable 11 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Ald-Ph-amido-PEG2 is a noncleavable ADC linker for antibody-drug conjugate.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 1 g</p>
<p>Ald-Ph-amido-PEG2-C2-NHS ester</p> <p>Cat. No.: HY-130104</p>	<p>Ald-Ph-amido-PEG2-C2-Pfp ester</p> <p>Cat. No.: HY-130103</p>
<p>Ald-Ph-amido-PEG2-C2-NHS ester is a noncleavable 2-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>Ald-Ph-amido-PEG2-C2-Pfp ester is a noncleavable 2-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>
<p>Ald-Ph-amido-PEG23-OPSS</p> <p>Cat. No.: HY-130962</p>	<p>Ald-Ph-amido-PEG3-C-COOH</p> <p>Cat. No.: HY-130098</p>
<p>Ald-Ph-amido-PEG23-OPSS is a cleavable 23 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Ald-Ph-amido-PEG3-C-COOH is a noncleavable linker used for the antibody-drug conjugate (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Ald-Ph-amido-PEG3-C1-Boc</p> <p>Cat. No.: HY-130100</p>	<p>Ald-Ph-amido-PEG3-C2-Pfp ester</p> <p>Cat. No.: HY-130102</p>
<p>Ald-Ph-amido-PEG3-C1-Boc is an ADC linker, which belongs to a polyethylene glycol (PEG) linker.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Ald-Ph-amido-PEG3-C2-Pfp ester is a noncleavable ADC linker, which belongs to a polyethylene glycol (PEG) linker.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>

<p>Ald-Ph-amido-PEG3-NHS ester</p> <p>Cat. No.: HY-133579</p> <p>Ald-Ph-amido-PEG3-NHS ester is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Ald-Ph-amido-PEG4-C2-acid</p> <p>Cat. No.: HY-130097</p> <p>Ald-Ph-amido-PEG4-C2-acid is a noncleavable linker used for the antibody-drug conjugate (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>
<p>Ald-Ph-amido-PEG4-C2-NHS ester</p> <p>Cat. No.: HY-130101</p> <p>Ald-Ph-amido-PEG4-C2-NHS ester is a noncleavable 4-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Ald-Ph-amido-PEG4-propargyl (Ald-benzyl-amide-PEG4-propargyl)</p> <p>Cat. No.: HY-133426</p> <p>Ald-Ph-amido-PEG4-propargyl is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Ald-Ph-NHS ester</p> <p>Cat. No.: HY-130107</p> <p>Ald-Ph-NHS ester is a noncleavable linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 250 mg</p>	<p>Ald-Ph-PEG4-bis-PEG3-methyltetrazine</p> <p>Cat. No.: HY-130974</p> <p>Ald-Ph-PEG4-bis-PEG3-methyltetrazine is a cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Ald-Ph-PEG4-bis-PEG3-N3</p> <p>Cat. No.: HY-130969</p> <p>Ald-Ph-PEG4-bis-PEG3-N3 is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Ald-Ph-PEG4-bis-PEG4-propargyl</p> <p>Cat. No.: HY-130967</p> <p>Ald-Ph-PEG4-bis-PEG4-propargyl is a cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Alkyne-PEG4-SS-PEG4-alkyne</p> <p>Cat. No.: HY-135970</p> <p>Alkyne-PEG4-SS-PEG4-alkyne is a cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Aloc-D-Ala-Phe-Lys(Aloc)-PAB-PNP</p> <p>Cat. No.: HY-129351</p> <p>Aloc-D-Ala-Phe-Lys(Aloc)-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>



<p>AMAS</p> <p style="text-align: right;">Cat. No.: HY-128925</p> <p>AMAS is a noncleavable heterobifunctional crosslinker with NHS ester and maleimide groups that allows covalent conjugation of amine- and sulfhydryl-containing molecules.</p>  <p>Purity: ≥97.0% Clinical Data: Size: 10 mg, 25 mg, 50 mg</p>	<p>Amino-bis-PEG3-BCN</p> <p style="text-align: right;">Cat. No.: HY-136085</p> <p>Amino-bis-PEG3-BCN is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Amino-bis-PEG3-DBCO</p> <p style="text-align: right;">Cat. No.: HY-130972</p> <p>Amino-bis-PEG3-DBCO is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Amino-bis-PEG3-TCO</p> <p style="text-align: right;">Cat. No.: HY-130955</p> <p>Amino-bis-PEG3-TCO is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Amino-ethyl-SS-PEG3-NHBoc</p> <p style="text-align: right;">Cat. No.: HY-140099</p> <p>Amino-ethyl-SS-PEG3-NHBoc is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Amino-PEG10-OH</p> <p style="text-align: right;">Cat. No.: HY-120761</p> <p>Amino-PEG10-OH is non-cleavable 10 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG10-OH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Amino-PEG11-OH</p> <p style="text-align: right;">Cat. No.: HY-130298</p> <p>Amino-PEG11-OH is non-cleavable 11 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG11-OH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Amino-PEG2-C2-acid</p> <p style="text-align: right;">Cat. No.: HY-W040168</p> <p>Amino-PEG2-C2-acid is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG2-C2-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>Amino-PEG3-C2-acid</p> <p style="text-align: right;">Cat. No.: HY-W040165</p> <p>Amino-PEG3-C2-acid is a cleavable PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG3-C2-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 250 mg, 500 mg</p>	<p>Amino-PEG3-SS-acid</p> <p style="text-align: right;">Cat. No.: HY-135974</p> <p>Amino-PEG3-SS-acid is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>











<p>Amino-PEG4-alcohol</p> <p style="text-align: right;">Cat. No.: HY-W008005</p>	<p>Amino-PEG4-bis-PEG3-methyltetrazine</p> <p style="text-align: right;">Cat. No.: HY-130970</p>
<p>Amino-PEG4-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Amino-PEG4-alcohol is also a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 100 mg</p>	<p>Amino-PEG4-bis-PEG3-methyltetrazine is a cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Amino-PEG4-bis-PEG3-N3</p> <p style="text-align: right;">Cat. No.: HY-136090</p>	<p>Amino-PEG4-bis-PEG3-propargyl</p> <p style="text-align: right;">Cat. No.: HY-130968</p>
<p>Amino-PEG4-bis-PEG3-N3 is a cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Amino-PEG4-bis-PEG3-propargyl is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Amino-PEG4-CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-130524</p>	<p>Amino-PEG4-Val-Cit-PAB-MMAE</p> <p style="text-align: right;">Cat. No.: HY-141154</p>
<p>Amino-PEG4-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Amino-PEG4-CH2COOH is also a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Amino-PEG4-Val-Cit-PAB-MMAE is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: 98.04% Clinical Data: No Development Reported Size: 1 mg, 5 mg, 10 mg, 25 mg</p>
<p>Amino-PEG5-C2-acid</p> <p style="text-align: right;">Cat. No.: HY-115384</p>	<p>Amino-PEG6-alcohol</p> <p style="text-align: right;">Cat. No.: HY-126942</p>
<p>Amino-PEG5-C2-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Amino-PEG5-C2-acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Amino-PEG6-alcohol is a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG6-alcohol is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Amino-PEG6-amido-bis-PEG5-N3</p> <p style="text-align: right;">Cat. No.: HY-130957</p>	<p>Amino-PEG8-Boc</p> <p style="text-align: right;">Cat. No.: HY-W019799</p>
<p>Amino-PEG6-amido-bis-PEG5-N3 is a cleavable 11 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Amino-PEG8-Boc is a cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG8-Boc is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>











<p>Amino-PEG9-acid</p> <p style="text-align: right;">Cat. No.: HY-130166</p>	<p>Amino-SS-PEG12-acid</p> <p style="text-align: right;">Cat. No.: HY-140097</p>
<p>Amino-PEG9-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Amino-PEG9-acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 10 mg, 25 mg, 50 mg, 100 mg</p>	<p>Amino-SS-PEG12-acid is a cleavable 12 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Amino-Tri-(carboxyethoxymethyl)-methane</p> <p style="text-align: right;">Cat. No.: HY-117519</p>	<p>Aminoethyl-SS-ethylalcohol</p> <p style="text-align: right;">Cat. No.: HY-117409</p>
<p>Amino-Tri-(carboxyethoxymethyl)-methane is a cleavable PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-Tri-(carboxyethoxymethyl)-methane is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Aminoethyl-SS-ethylalcohol is a glutathione cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Aminoethyl-SS-propionic acid</p> <p style="text-align: right;">Cat. No.: HY-140096</p>	<p>Aminoxy-amido-PEG4-propargyl</p> <p style="text-align: right;">Cat. No.: HY-133435</p>
<p>Aminoethyl-SS-propionic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 10 mg, 25 mg, 50 mg</p>	<p>Aminoxy-amido-PEG4-propargyl is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Aminoxy-PEG2-alcohol</p> <p style="text-align: right;">Cat. No.: HY-126951</p>	<p>Aminoxy-PEG2-azide</p> <p style="text-align: right;">Cat. No.: HY-113931</p>
<p>Aminoxy-PEG2-alcohol is a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Aminoxy-PEG2-alcohol is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Aminoxy-PEG2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Aminoxy-PEG2-azide is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Aminoxy-PEG2-BCN</p> <p style="text-align: right;">Cat. No.: HY-145593</p>	<p>Aminoxy-PEG2-bis-PEG3-BCN</p> <p style="text-align: right;">Cat. No.: HY-136089</p>
<p>Aminoxy-PEG2-BCN is a cleavable PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Aminoxy-PEG2-bis-PEG3-BCN is a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>











<p>Aminoxy-PEG3-azide</p> <p style="text-align: right;">Cat. No.: HY-126949</p> <p>Aminoxy-PEG3-azide is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Aminoxy-C2-PEG3-azide is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Aminoxy-PEG4-alcohol</p> <p style="text-align: right;">Cat. No.: HY-124123</p> <p>Aminoxy-PEG4-alcohol is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Aminoxy-PEG4-alcohol is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Aminoxyacetamide-PEG3-azide</p> <p style="text-align: right;">Cat. No.: HY-133434</p> <p>Aminoxyacetamide-PEG3-azide is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>APN-PEG36-tetrazine</p> <p style="text-align: right;">Cat. No.: HY-139859</p> <p>APN-PEG36-tetrazine is an analogue of APN-PEG4-tetrazine. APN-PEG4-tetrazine is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: 96.05% Clinical Data: No Development Reported Size: 5 mg, 10 mg</p>
<p>APN-PEG4-Amine hydrochloride</p> <p style="text-align: right;">Cat. No.: HY-130939</p> <p>APN-PEG4-Amine (hydrochloride) is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>APN-PEG4-BCN</p> <p style="text-align: right;">Cat. No.: HY-136044</p> <p>APN-PEG4-BCN is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>APN-PEG4-DBCO</p> <p style="text-align: right;">Cat. No.: HY-136049</p> <p>APN-PEG4-DBCO is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>APN-PEG4-tetrazine</p> <p style="text-align: right;">Cat. No.: HY-136045</p> <p>APN-PEG4-tetrazine is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Azetidin-3-ol hydrochloride</p> <p style="text-align: right;">Cat. No.: HY-40144</p> <p>Azetidin-3-ol hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azetidin-3-ol hydrochloride is also an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg, 1 g</p>	<p>Azetidine-3-carboxylic acid</p> <p style="text-align: right;">Cat. No.: HY-Y0530</p> <p>Azetidine-3-carboxylic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azetidine-3-carboxylic acid is also an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs¹².</p> <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>











<p>Azide-C2-Azide</p> <p style="text-align: right;">Cat. No.: HY-138535</p>	<p>Azide-C2-SS-C2-biotin</p> <p style="text-align: right;">Cat. No.: HY-140127</p>
<p>Azide-C2-Azide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Azide-C2-SS-C2-biotin is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 25 mg, 50 mg, 100 mg</p>
<p>Azide-PEG1-Val-Cit-PABC-OH</p> <p style="text-align: right;">Cat. No.: HY-136137</p>	<p>Azide-PEG3-Tos</p> <p style="text-align: right;">Cat. No.: HY-140004</p>
<p>Azide-PEG1-Val-Cit-PABC-OH is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Azide-PEG3-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Azide-PEG3-Tos is also a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Azide-PEG5-Tos</p> <p style="text-align: right;">Cat. No.: HY-140352</p>	<p>Azido-C2-SS-PEG2-C2-acid</p> <p style="text-align: right;">Cat. No.: HY-140101</p>
<p>Azide-PEG5-Tos is a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Azido-C2-SS-PEG2-C2-acid is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Azido-C6-OH</p> <p style="text-align: right;">Cat. No.: HY-138521</p>	<p>Azido-PEG1-Val-Cit-OH</p> <p style="text-align: right;">Cat. No.: HY-136034</p>
<p>Azido-C6-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Azido-PEG1-Val-Cit-OH is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Azido-PEG1-Val-Cit-PABC-PNP</p> <p style="text-align: right;">Cat. No.: HY-136105</p>	<p>Azido-PEG2-C2-amine (N3-PEG2-CH2CH2NH2)</p> <p style="text-align: right;">Cat. No.: HY-140213</p>
<p>Azido-PEG1-Val-Cit-PABC-PNP is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Azido-PEG2-C2-amine (N3-PEG2-CH2CH2NH2) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Azido-PEG2-C2-amine is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 250 mg, 500 mg</p>










<p>Azido-PEG3-maleimide</p> <p>Cat. No.: HY-140811</p>	<p>Azido-PEG3-SS-NHS</p> <p>Cat. No.: HY-135966</p>
<p>Azido-PEG3-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Azido-PEG3-maleimide is also a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 25 mg</p>	<p>Azido-PEG3-SS-NHS is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 25 mg, 50 mg</p>
<p>Azido-PEG3-SSPy</p> <p>Cat. No.: HY-136038</p>	<p>Azido-PEG3-Val-Cit-PAB-OH</p> <p>Cat. No.: HY-140148</p>
<p>Azido-PEG3-SSPy is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Azido-PEG3-Val-Cit-PAB-OH is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Azido-PEG3-Val-Cit-PAB-PNP</p> <p>Cat. No.: HY-140150</p>	<p>Azido-PEG4-C2-acid</p> <p>Cat. No.: HY-130653</p>
<p>Azido-PEG3-Val-Cit-PAB-PNP is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG3-Val-Cit-PAB-PNP is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>	<p>Azido-PEG4-C2-acid a PEG-based PROTAC linker can be used in the synthesis of vRucaparib-TP4. Azido-PEG4-C2-acid is also a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg, 250 mg, 500 mg</p>
<p>Azido-PEG4-CH2-Boc</p> <p>Cat. No.: HY-42618</p>	<p>Azido-PEG4-Val-Cit-PAB-OH</p> <p>Cat. No.: HY-140149</p>
<p>Azido-PEG4-CH2-Boc is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG4-CH2-Boc is also a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>Azido-PEG4-Val-Cit-PAB-OH is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG4-Val-Cit-PAB-OH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>
<p>Azido-PEG5-acid</p> <p>Cat. No.: HY-130572</p>	<p>Azido-PEG5-Ala-Ala-Asn-PAB</p> <p>Cat. No.: HY-141150</p>
<p>Azido-PEG5-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs, such as the conjugate CPT-APO (CPT: Camptothecin (HY-16560)). Azido-PEG5-acid is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Azido-PEG5-Ala-Ala-Asn-PAB is a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>











<p>Azido-PEG5-alcohol</p> <p style="text-align: right;">Cat. No.: HY-130211</p> <p>Azido-PEG5-alcohol is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG5-alcohol is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Azido-PEG5-CH2CO2H</p> <p style="text-align: right;">Cat. No.: HY-130194</p> <p>Azido-PEG5-CH2CO2H is a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG5-CH2CO2H is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: 99.60% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>Azido-PEG6-alcohol</p> <p style="text-align: right;">Cat. No.: HY-130537</p> <p>Azido-PEG6-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Azido-PEG6-alcohol is also a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Azido-PEG6-amine</p> <p style="text-align: right;">Cat. No.: HY-140215</p> <p>Azido-PEG6-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Azido-PEG6-amine is also a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 200 mg, 500 mg</p>
<p>Azido-PEG6-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130474</p> <p>Azido-PEG6-NHS ester is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG6-NHS ester is also a PEG- and Alkyl/ether based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: 98.85% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Azido-PEG7-amine</p> <p style="text-align: right;">Cat. No.: HY-130324</p> <p>Azido-PEG7-amine is a non-cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG7-amine is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>
<p>Azido-PEG8-acid</p> <p style="text-align: right;">Cat. No.: HY-140454</p> <p>Azido-PEG8-acid is a non-cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG8-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>Azido-PEG8-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130184</p> <p>Azido-PEG8-NHS ester is a cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG8-NHS ester is also a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Azido-PEG9-acid</p> <p style="text-align: right;">Cat. No.: HY-130475</p> <p>Azido-PEG9-acid is a non-cleavable 9 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG9-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Azido-PEG9-amine</p> <p style="text-align: right;">Cat. No.: HY-130169</p> <p>Azido-PEG9-amine is a non-cleavable 9 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG9-amine is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>


<p>Azidoethyl-SS-ethylamine</p> <p>Cat. No.: HY-140104</p>	<p>Azidoethyl-SS-ethylazide</p> <p>Cat. No.: HY-140105</p>
<p>Azidoethyl-SS-ethylamine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Azidoethyl-SS-ethylazide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Azidoethyl-SS-propionic acid</p> <p>Cat. No.: HY-140100</p>	<p>Azidoethyl-SS-propionic NHS ester</p> <p>Cat. No.: HY-140102</p>
<p>Azidoethyl-SS-propionic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Azidoethyl-SS-propionic NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>BCN-PEG1-Val-Cit-OH</p> <p>Cat. No.: HY-130922</p>	<p>BCN-PEG1-Val-Cit-PABC-OH</p> <p>Cat. No.: HY-130923</p>
<p>BCN-PEG1-Val-Cit-OH is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>BCN-PEG1-Val-Cit-PABC-OH is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>BCN-PEG3-Biotin</p> <p>Cat. No.: HY-130924</p>	<p>BCN-PEG3-oxyamine</p> <p>Cat. No.: HY-130926</p>
<p>BCN-PEG3-Biotin is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: Size: 5 mg, 10 mg, 25 mg, 50 mg, 100 mg</p>	<p>BCN-PEG3-oxyamine is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>BCN-PEG3-Val-Cit</p> <p>Cat. No.: HY-140151</p>	<p>BCN-PEG3-VC-PFP ester</p> <p>Cat. No.: HY-140152</p>
<p>BCN-PEG3-Val-Cit is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. BCN-PEG3-Val-Cit is also a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>BCN-PEG3-VC-PFP ester is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>










<p>BCN-PEG4-acid</p> <p style="text-align: right;">Cat. No.: HY-135971</p> <p>BCN-PEG4-acid is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg, 250 mg</p>	<p>BCN-PEG4-HyNic</p> <p style="text-align: right;">Cat. No.: HY-136061</p> <p>BCN-PEG4-HyNic is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>BCN-PEG4-OH</p> <p style="text-align: right;">Cat. No.: HY-130925</p> <p>BCN-PEG4-OH is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>BCN-PEG4-Ts</p> <p style="text-align: right;">Cat. No.: HY-130927</p> <p>BCN-PEG4-Ts is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>BCN-SS-amine</p> <p style="text-align: right;">Cat. No.: HY-135972</p> <p>BCN-SS-amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>BCN-SS-NHS</p> <p style="text-align: right;">Cat. No.: HY-135973</p> <p>BCN-SS-NHS is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 25 mg</p>
<p>BCOT-PEF3-OPFP</p> <p style="text-align: right;">Cat. No.: HY-136125</p> <p>BCOT-PEF3-OPFP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>BDP FL DBCO</p> <p style="text-align: right;">Cat. No.: HY-140296</p> <p>BDP FL DBCO is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Biotin-C4-amide-C5-NH2</p> <p style="text-align: right;">Cat. No.: HY-W096148</p> <p>Biotin-C4-amide-C5-NH2 is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Biotin-PEG1-azide</p> <p style="text-align: right;">Cat. No.: HY-W096133</p> <p>Biotin-PEG1-azide is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>Biotin-PEG1-NH2</p> <p style="text-align: right;">Cat. No.: HY-W096135</p>	<p>Biotin-PEG2-acid</p> <p style="text-align: right;">Cat. No.: HY-126958</p>
<p>Biotin-PEG1-NH2 is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Biotin-PEG2-acid is a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Biotin-PEG2-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: 96.14% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 250 mg</p>
<p>Biotin-PEG2-methyl ethanethioate</p> <p style="text-align: right;">Cat. No.: HY-138508</p>	<p>Biotin-PEG3-aldehyde</p> <p style="text-align: right;">Cat. No.: HY-136051</p>
<p>Biotin-PEG2-methyl ethanethioate is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Biotin-PEG3-aldehyde is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Biotin-PEG3-SS-azide</p> <p style="text-align: right;">Cat. No.: HY-140944</p>	<p>Biotin-PEG4-Dde-TAMRA-PEG3-Azide</p> <p style="text-align: right;">Cat. No.: HY-141091</p>
<p>Biotin-PEG3-SS-azide is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥98.0% Clinical Data: Size: 5 mg</p>	<p>Biotin-PEG4-Dde-TAMRA-PEG3-Azide is a cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Biotin-PEG4-PFP ester</p> <p style="text-align: right;">Cat. No.: HY-138488</p>	<p>Biotin-PEG4-SS-azide</p> <p style="text-align: right;">Cat. No.: HY-139107</p>
<p>Biotin-PEG4-PFP ester is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Biotin-PEG4-SS-azide is a cleavable, biotin-labeled, ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 25 mg, 50 mg</p>
<p>Biotin-sar-oh</p> <p style="text-align: right;">Cat. No.: HY-W096127</p>	<p>Bis-(PEG6-acid)-SS</p> <p style="text-align: right;">Cat. No.: HY-140115</p>
<p>Biotin-sar-oh is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bis-(PEG6-acid)-SS is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>

<p>Bis-PEG1-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130089</p> <p>Bis-PEG1-NHS ester is a noncleavable 1-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>	<p>Bis-PEG1-PFP ester</p> <p style="text-align: right;">Cat. No.: HY-112561</p> <p>Bis-PEG1-PFP ester is a non-cleavable (1 unit PEG) ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Bis-PEG10-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130824</p> <p>Bis-PEG10-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG10-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bis-PEG13-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130825</p> <p>Bis-PEG13-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG13-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Bis-PEG17-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130826</p> <p>Bis-PEG17-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG17-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>bis-PEG2-endo-BCN</p> <p style="text-align: right;">Cat. No.: HY-140078</p> <p>bis-PEG2-endo-BCN is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.10% Clinical Data: Size: 50 mg</p>
<p>Bis-PEG2-PFP ester</p> <p style="text-align: right;">Cat. No.: HY-112560</p> <p>Bis-PEG2-PFP ester is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Bis-PEG2-PFP ester is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bis-PEG21-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130827</p> <p>Bis-PEG21-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG21-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Bis-PEG25-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130828</p> <p>Bis-PEG25-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG25-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bis-PEG3-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130087</p> <p>Bis-PEG3-NHS ester is a noncleavable 3-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>Bis-PEG5-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126889</p> <p>Bis-PEG5-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG5-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bis-PEG6-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130410</p> <p>Bis-PEG6-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG6-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>
<p>Bis-PEG7-acid</p> <p style="text-align: right;">Cat. No.: HY-126892</p> <p>Bis-PEG7-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG6-propionic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bis-PEG7-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126890</p> <p>Bis-PEG7-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG7-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>Bis-PEG8-acid</p> <p style="text-align: right;">Cat. No.: HY-126893</p> <p>Bis-PEG8-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG8-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bis-PEG9-acid</p> <p style="text-align: right;">Cat. No.: HY-126894</p> <p>Bis-PEG9-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG9-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Bis-PEG9-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-117009</p> <p>Bis-PEG9-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG9-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>Bis-SS-C3-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133584</p> <p>Bis-SS-C3-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Bis-SS-C3-sulfo-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133585</p> <p>Bis-SS-C3-sulfo-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bis-sulfone-PEG3-Azide</p> <p style="text-align: right;">Cat. No.: HY-138745</p> <p>Bis-sulfone-PEG3-Azide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 5 mg</p>











<p>Bis-Tos-(2-hydroxyethyl disulfide)</p> <p style="text-align: right;">Cat. No.: HY-140126</p> <p>Bis-Tos-(2-hydroxyethyl disulfide) is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bis-PEG2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130088</p> <p>Bis-PEG2-NHS ester is a noncleavable 2-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 50 mg</p>
<p>Bis-PEG4-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130086</p> <p>Bis-PEG4-NHS ester is a noncleavable 4-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>BMPS</p> <p style="text-align: right;">Cat. No.: HY-42146</p> <p>BMPS is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.93% Clinical Data: No Development Reported Size: 100 mg, 500 mg, 1 g</p>
<p>BnO-PEG6-OH</p> <p style="text-align: right;">Cat. No.: HY-W042654</p> <p>BnO-PEG6-OH is a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). BnO-PEG6-OH is also a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: 99.88% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Boc-amino-PEG3-SS-acid</p> <p style="text-align: right;">Cat. No.: HY-136037</p> <p>Boc-amino-PEG3-SS-acid is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Boc-amino-PEG3-SSPy</p> <p style="text-align: right;">Cat. No.: HY-136041</p> <p>Boc-amino-PEG3-SSPy is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Boc-aminoxy-amide-PEG4-propargyl</p> <p style="text-align: right;">Cat. No.: HY-133436</p> <p>Boc-aminoxy-amide-PEG4-propargyl is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Boc-aminoxy-ethyl-SS-propanol</p> <p style="text-align: right;">Cat. No.: HY-140117</p> <p>Boc-aminoxy-ethyl-SS-propanol is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Boc-Aminoxy-PEG2-bromide</p> <p style="text-align: right;">Cat. No.: HY-135962</p> <p>Boc-Aminoxy-PEG2-bromide is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>

<p>Boc-C14-COOH</p> <p style="text-align: right;">Cat. No.: HY-W034599</p> <p>Boc-C14-COOH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-C14-COOH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Boc-C16-COOH</p> <p style="text-align: right;">Cat. No.: HY-W045598</p> <p>Boc-C16-COOH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-C16-COOH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<su.</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 250 mg</p>
<p>Boc-C2-Urea-bis(Boc)-C4-Urea-4-phenylacetic acid</p> <p style="text-align: right;">Cat. No.: HY-108379</p> <p>Boc-C2-Urea-bis(Boc)-C4-Urea-4-phenylacetic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Boc-Cystamine</p> <p style="text-align: right;">Cat. No.: HY-140098</p> <p>Boc-Cystamine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Boc-Dap-NE</p> <p style="text-align: right;">Cat. No.: HY-78931</p> <p>Boc-Dap-NE, a dipeptide, is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 50 mg, 100 mg</p>	<p>Boc-Gly-Gly-Phe-Gly-OH</p> <p style="text-align: right;">Cat. No.: HY-P1449</p> <p>Boc-Gly-Gly-Phe-Gly-OH, a self-assembly of N- and C-protected tetrapeptide, is a protease cleavable linker used for the antibody-drug conjugate (ADC).</p>  <p>Purity: 99.10% Clinical Data: No Development Reported Size: 10 mg</p>
<p>Boc-Gly-Gly-Phe-Gly-OH TFA</p> <p style="text-align: right;">Cat. No.: HY-P1449A</p> <p>Boc-Gly-Gly-Phe-Gly-OH TFA, a self-assembly of N- and C-protected tetrapeptide, is a protease cleavable linker used for the antibody-drug conjugate (ADC).</p>  <p>Purity: 98.27% Clinical Data: No Development Reported Size: 10 mg</p>	<p>Boc-gly-PEG3-endo-BCN</p> <p style="text-align: right;">Cat. No.: HY-140081</p> <p>Boc-gly-PEG3-endo-BCN is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Boc-gly-PEG3-endo-BCN is also a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Boc-Hyp-OH</p> <p style="text-align: right;">Cat. No.: HY-10781</p> <p>Boc-Hyp-OH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-Hyp-OH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>5</sup>.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g, 5 g</p>	<p>Boc-Hyp-OMe</p> <p style="text-align: right;">Cat. No.: HY-65039</p> <p>Boc-Hyp-OMe is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-Hyp-OMe is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g, 5 g</p>







<p>Boc-NH-C6-Br</p> <p style="text-align: right;">Cat. No.: HY-W011561</p> <p>Boc-NH-C6-Br is a non-cleavable linker used for antibody-drug conjugates (ADC).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Boc-NH-ethyl-SS-propionic acid</p> <p style="text-align: right;">Cat. No.: HY-140116</p> <p>Boc-NH-ethyl-SS-propionic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 100 mg, 250 mg, 500 mg</p>
<p>Boc-NH-PEG1-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-120775</p> <p>Boc-NH-PEG1-CH2CH2COOH is a cleavable (1 unit PEG) ADC linker and also a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of antibody-drug conjugates (ADCs) or PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Boc-NH-PEG3-C2-triazole-DBCO-PEG4-VC-PAB-DMEA</p> <p style="text-align: right;">Cat. No.: HY-126677</p> <p>Boc-NH-PEG3-C2-triazole-DBCO-PEG4-VC-PAB-DMEA is a double cleavable 3-unit and 4-unit PEG linker for antibody-drug-conjugation (ADC). Boc-NH-PEG3-C2-triazole-DBCO-PEG4-VC-PAB-DMEA also is a PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Boc-NH-PEG4-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W040132</p> <p>Boc-NH-PEG4-CH2CH2COOH is a PEG-based PROTAC linker can be used in the synthesis of PROTAC. Boc-NH-PEG4-CH2CH2COOH is also a cleavable ADC linker used as a linker for antibody-drug conjugates (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Boc-NH-PEG4-CH2CH2NH2</p> <p style="text-align: right;">Cat. No.: HY-W008352</p> <p>Boc-NH-PEG4-CH2CH2NH2 a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-NH-PEG4-CH2CH2NH2 is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Boc-NH-PEG4-CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-42640</p> <p>Boc-NH-PEG4-CH2COOH is a cleavable ADC linker used as a linker for antibody-drug conjugates (ADC). Boc-NH-PEG4-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>Boc-NH-PEG6-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W040244</p> <p>Boc-NH-PEG6-CH2CH2COOH is a cleavable ADC linker used as a linker for antibody-drug conjugates (ADC). Boc-NH-PEG6-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Boc-NMe-Val-Val-Dil-Dap-OH</p> <p style="text-align: right;">Cat. No.: HY-130956</p> <p>Boc-NMe-Val-Val-Dil-Dap-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Boc-Phe-(Alloc)Lys-PAB-PNP</p> <p style="text-align: right;">Cat. No.: HY-129353</p> <p>Boc-Phe-(Alloc)Lys-PAB-PNP is used as a cleavable linker for antibody-drug conjugates (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g</p>

<p>Boc-trans-D-Hyp-OMe</p> <p>Cat. No.: HY-W017882</p> <p>Boc-trans-D-Hyp-OMe is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-trans-D-Hyp-OMe is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>	<p>Boc-Val-Ala-PAB-PNP</p> <p>Cat. No.: HY-130932</p> <p>Boc-Val-Ala-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>
<p>Boc-Val-Cit-OH</p> <p>Cat. No.: HY-W038702</p> <p>Boc-Val-Cit-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 250 mg, 500 mg, 1 g</p>	<p>Boc-Val-Cit-PAB</p> <p>Cat. No.: HY-141141</p> <p>Boc-Val-Cit-PAB is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.99% Clinical Data: Size: 25 mg, 50 mg, 100 mg</p>
<p>Boc-Val-Cit-PAB-PNP</p> <p>Cat. No.: HY-141142</p> <p>Boc-Val-Cit-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.74% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 50 mg, 100 mg</p>	<p>Boc-Val-Dil-Dap-OH</p> <p>Cat. No.: HY-130961</p> <p>Boc-Val-Dil-Dap-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.86% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Boc-Val-Dil-Dap-Phe-OMe</p> <p>Cat. No.: HY-130975</p> <p>Boc-Val-Dil-Dap-Phe-OMe is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Bocaminoxyacetamide-PEG2-Azido</p> <p>Cat. No.: HY-136099</p> <p>Bocaminoxyacetamide-PEG2-Azido is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Bocaminoxyacetamide-PEG3-alkyne</p> <p>Cat. No.: HY-136101</p> <p>Bocaminoxyacetamide-PEG3-alkyne is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Br-PEG4-C2-Boc</p> <p>Cat. No.: HY-130315</p> <p>Br-PEG4-C2-Boc is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 250 mg</p>











<p>Bromo-PEG2-C2-azide</p> <p>Cat. No.: HY-130485</p> <p>Bromo-PEG2-C2-azide is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Bromo-PEG2-C2-azide is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: 98.10% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>	<p>Bromoacetamido-PEG4-acid</p> <p>Cat. No.: HY-141382</p> <p>Bromoacetamido-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Bromoacetamido-PEG4-acid is also a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>BS2G Crosslinker disodium</p> <p>Cat. No.: HY-130547</p> <p>BS2G Crosslinker (disodium) is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>BS3 Crosslinker</p> <p>Cat. No.: HY-124329</p> <p>BS3 Crosslinker is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>BS3 Crosslinker disodium</p> <p>Cat. No.: HY-124329A</p> <p>BS3 Crosslinker disodium is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Cbz-Phe-(Alloc)Lys-PAB-PNP</p> <p>Cat. No.: HY-129352</p> <p>Cbz-Phe-(Alloc)Lys-PAB-PNP is a cleavable linker for antibody-drug conjugates (ADC) design.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>
<p>cis-4-Hydroxy-D-proline hydrochloride</p> <p>Cat. No.: HY-76104</p> <p>cis-4-Hydroxy-D-proline hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). cis-4-Hydroxy-D-proline hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g, 5 g</p>	<p>cis-4-Hydroxy-L-proline hydrochloride</p> <p>Cat. No.: HY-W019213</p> <p>cis-4-Hydroxy-L-proline hydrochloride is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). cis-4-Hydroxy-L-proline hydrochloride is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 250 mg, 500 mg</p>
<p>CL2 Linker</p> <p>Cat. No.: HY-128947</p> <p>CL2 Linker is a cleavable ADC linker. CL2-SN-38 and CL2A-SN-38 are equivalent in drug substitution (~6), cell binding (K_d ~1.2 nM), cytotoxicity (IC_{50} ~2.2 nM), and serum stability in vitro ($t_{1/2}$ ~20 hours).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>CL2A</p> <p>Cat. No.: HY-128945</p> <p>CL2A is a cleavable complicated PEG8- and triazole-containing PABC-peptide-mc linker. CL2A is cleavable through pH sensitivity, giving rise to bystander effect, and binds the antibody at a cysteine residue via a disulfide bond. Labetuzumab govitcan used this linker.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 5 mg, 10 mg</p>











<p>Cyclooctyne-O-amido-PEG2-PFP ester</p> <p>Cat. No.: HY-133573</p>	<p>Cyclooctyne-O-amido-PEG3-PFP ester</p> <p>Cat. No.: HY-133575</p>
<p>Cyclooctyne-O-amido-PEG2-PFP ester is a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Cyclooctyne-O-amido-PEG3-PFP ester is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Cyclooctyne-O-amido-PEG4-PFP ester</p> <p>Cat. No.: HY-133576</p>	<p>Cyclooctyne-O-NHS ester</p> <p>Cat. No.: HY-126517</p>
<p>Cyclooctyne-O-amido-PEG4-PFP ester is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Cyclooctyne-O-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 10 mg, 100 mg, 500 mg, 1 g</p>
<p>Cyclooctyne-O-PFP ester</p> <p>Cat. No.: HY-126518</p>	<p>D-Proline, 4-hydroxy-, methyl ester hydrochloride</p> <p>Cat. No.: HY-76105</p>
<p>Cyclooctyne-O-PFP ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 95.05% Clinical Data: No Development Reported Size: 100 mg, 1 g</p>	<p>D-Proline, 4-hydroxy-, methyl ester hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-(PEG2-Val-Cit-PAB)2</p> <p>Cat. No.: HY-126676</p>	<p>DBCO-acid</p> <p>Cat. No.: HY-42972</p>
<p>DBCO-(PEG2-Val-Cit-PAB)2 is a dual cleavable ADC linker for antibody-drug conjugates (ADCs). DBCO-(PEG2-Val-Cit-PAB)2 is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>DBCO-acid is a cleavable ADC linker used in the synthesis of ADC linker DBCO-NHS ester (HY-115524 and HY-115545), and drug-linker conjugates DBCO-PEG-MMAE (HY-111012 and HY-126690).</p>  <p>Purity: 99.63% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 10 mg, 50 mg, 100 mg</p>
<p>DBCO-amine</p> <p>Cat. No.: HY-W000423</p>	<p>DBCO-C3-Acid</p> <p>Cat. No.: HY-120903</p>
<p>DBCO-amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.86% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>	<p>DBCO-C3-Acid is a Click Chemistry intermediate used in the synthesis of antibody-drug conjugate (ADC) linker.</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>

<p>DBCO-CONH-S-S-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133413</p> <p>DBCO-CONH-S-S-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 95.04% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>DBCO-Maleimide</p> <p style="text-align: right;">Cat. No.: HY-116270</p> <p>DBCO-Maleimide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 96.41% Clinical Data: No Development Reported Size: 10 mg, 50 mg, 100 mg, 250 mg</p>
<p>DBCO-N-bis(PEG4-NHS ester)</p> <p style="text-align: right;">Cat. No.: HY-145090</p> <p>DBCO-N-bis(PEG4-NHS ester) is a PEG linker which contains two PEG4-NHS ester and a DBCO group. DBCO-N-bis(PEG4-NHS ester) is useful for protein modification or labeling.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>DBCO-NHCO-PEG4-acid</p> <p style="text-align: right;">Cat. No.: HY-125541</p> <p>DBCO-Amide-PEG5-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-Amide-PEG5-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-NHCO-PEG4-amine</p> <p style="text-align: right;">Cat. No.: HY-124386</p> <p>DBCO-NHCO-PEG4-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-NHCO-PEG4-amine is a cleavable ADC linker used to conjugate MMAE (HY-15162) and antibody (e.g).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>DBCO-NHCO-PEG4-NH-Boc</p> <p style="text-align: right;">Cat. No.: HY-126884</p> <p>DBCO-NHCO-PEG4-NH-Boc is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-NHCO-PEG4-NH-Boc is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-NHCO-PEG4-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-111456</p> <p>DBCO-NHCO-PEG4-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-NHCO-PEG4-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>DBCO-NHCO-S-S-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133412</p> <p>DBCO-NHCO-S-S-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-42973</p> <p>DBCO-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.53% Clinical Data: No Development Reported Size: 10 mg, 50 mg, 100 mg</p>	<p>DBCO-NHS ester 2</p> <p style="text-align: right;">Cat. No.: HY-115524</p> <p>DBCO-NHS ester 2 is a cleavable linker that is used for making antibody-drug conjugate (ADC). DBCO-NHS ester 2 is a derivative of Dibenzylcyclooctyne (DBCO) used in copper-free click chemistry.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>DBCO-NHS ester 3</p> <p style="text-align: right;">Cat. No.: HY-115545</p> <p>DBCO-NHS ester 3 (Compound 12) is a cleavable linker that is used for making antibody-drug conjugate (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>DBCO-PEG3 acetic-EVCit-PAB</p> <p style="text-align: right;">Cat. No.: HY-136096</p> <p>DBCO-PEG3 acetic-EVCit-PAB is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-PEG3-oxyamine</p> <p style="text-align: right;">Cat. No.: HY-133429</p> <p>DBCO-PEG3-oxyamine is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>DBCO-PEG3-propionic EVCit-PAB</p> <p style="text-align: right;">Cat. No.: HY-136141</p> <p>DBCO-PEG3-propionic EVCit-PAB is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-PEG3-SS-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133431</p> <p>DBCO-PEG3-SS-NHS ester is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥90.0% Clinical Data: Size: 25 mg</p>	<p>DBCO-PEG3-TCO</p> <p style="text-align: right;">Cat. No.: HY-133428</p> <p>DBCO-PEG3-TCO is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>DBCO-PEG4-acetic-Val-Cit-PAB</p> <p style="text-align: right;">Cat. No.: HY-136098</p> <p>DBCO-PEG4-acetic-Val-Cit-PAB is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>	<p>DBCO-PEG4-alkyne</p> <p style="text-align: right;">Cat. No.: HY-133430</p> <p>DBCO-PEG4-alkyne is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>DBCO-PEG4-amine</p> <p style="text-align: right;">Cat. No.: HY-130435</p> <p>DBCO-PEG4-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-PEG4-amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>	<p>DBCO-PEG4-DBCO</p> <p style="text-align: right;">Cat. No.: HY-130346</p> <p>DBCO-PEG4-DBCO is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-PEG4-DBCO is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>









<p>DBCO-PEG4-HyNic</p> <p>Cat. No.: HY-136067</p>	<p>DBCO-PEG4-Maleimide</p> <p>Cat. No.: HY-120770</p>
<p>DBCO-PEG4-HyNic is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 25 mg, 50 mg, 100 mg</p>	<p>DBCO-PEG4-Maleimide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.02% Clinical Data: No Development Reported Size: 100 mg</p>
<p>DBCO-PEG4-Propionic-Val-Cit-PAB</p> <p>Cat. No.: HY-136103</p>	<p>DBCO-PEG4-SS-TCO</p> <p>Cat. No.: HY-133432</p>
<p>DBCO-PEG4-Propionic-Val-Cit-PAB is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>DBCO-PEG4-SS-TCO is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>DBCO-PEG5-NHS ester</p> <p>Cat. No.: HY-126885</p>	<p>DBCO-S-S-acid</p> <p>Cat. No.: HY-138506</p>
<p>DBCO-PEG5-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-PEG5-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>DBCO-S-S-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-SS-aldehyde</p> <p>Cat. No.: HY-135977</p>	<p>DBCO-SS-amine</p> <p>Cat. No.: HY-135978</p>
<p>DBCO-SS-aldehyde is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>DBCO-SS-amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-SS-PEG4-Biotin</p> <p>Cat. No.: HY-135979</p>	<p>DBCO-Sulfo-Link-biotin</p> <p>Cat. No.: HY-130810</p>
<p>DBCO-SS-PEG4-Biotin is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>DBCO-Sulfo-Link-biotin is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg</p>

<p>DBCO-Sulfo-NHS ester sodium</p> <p>Cat. No.: HY-123687</p>	<p>DBCO-Val-Cit-OH</p> <p>Cat. No.: HY-130935</p>
<p>DBCO-Sulfo-NHS ester sodium is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 25 mg</p>	<p>DBCO-Val-Cit-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-Val-Cit-PABC-OH</p> <p>Cat. No.: HY-130936</p>	<p>DBCO-Val-Cit-PABC-PNP</p> <p>Cat. No.: HY-130937</p>
<p>DBCO-Val-Cit-PABC-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>DBCO-Val-Cit-PABC-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>DBCO-C6-acid</p> <p>Cat. No.: HY-121805</p>	<p>Diazo Biotin-PEG3-DBCO</p> <p>Cat. No.: HY-140930</p>
<p>DBCO-C6-acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). DBCO-C6-acid can be used in the synthesis of camphycin analogues.</p>  <p>Purity: 95.38% Clinical Data: No Development Reported Size: 25 mg</p>	<p>Diazo Biotin-PEG3-DBCO is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Dimethylamine-SPDB</p> <p>Cat. No.: HY-133542</p>	<p>DM21</p> <p>Cat. No.: HY-139441</p>
<p>Dimethylamine-SPDB is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>DM21 is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.67% Clinical Data: No Development Reported Size: 5 mg, 10 mg</p>
<p>DMAC-PDB</p> <p>Cat. No.: HY-126531</p>	<p>DMAC-SPDB</p> <p>Cat. No.: HY-133550</p>
<p>DMAC-PDB is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>DMAC-SPDB is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>DMAC-SPDB-sulfo</p> <p style="text-align: right;">Cat. No.: HY-131084</p> <p>DMAC-SPDB-sulfo is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>DMAC-SPP</p> <p style="text-align: right;">Cat. No.: HY-130111</p> <p>DMAC-SPP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Docosanedioic acid</p> <p style="text-align: right;">Cat. No.: HY-W034918</p> <p>Docosanedioic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Docosanedioic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>	<p>DOTA-NHS-ester</p> <p style="text-align: right;">Cat. No.: HY-128890</p> <p>DOTA-NHS-ester is a linker for affibody molecules and is applied in small animals PET, SPECT, and CT. DOTA-NHS-ester can be used to label radiotherapeutic agents or imaging probes for the detection of tumors.</p>  <p>Purity: ≥90.0% Clinical Data: No Development Reported Size: 100 mg</p>
<p>DSG Crosslinker</p> <p style="text-align: right;">Cat. No.: HY-114697</p> <p>DSG Crosslinker is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.39% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 10 mg, 50 mg, 100 mg</p>	<p>DSP Crosslinker</p> <p style="text-align: right;">Cat. No.: HY-118759</p> <p>DSP Crosslinker is a cleavable ADC linker, used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.73% Clinical Data: No Development Reported Size: 100 mg, 250 mg, 500 mg</p>
<p>DSS Crosslinker</p> <p style="text-align: right;">Cat. No.: HY-W019543</p> <p>DSS Crosslinker is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.73% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 100 mg</p>	<p>DTSSP Crosslinker</p> <p style="text-align: right;">Cat. No.: HY-126349</p> <p>DTSSP Crosslinker is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>EC089</p> <p style="text-align: right;">Cat. No.: HY-128940</p> <p>EC089 is a cleavable linker used in conjugates of tubulysins and folates, and extracted from patent WO2011069116A1.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Eicosanedioic acid</p> <p style="text-align: right;">Cat. No.: HY-W034595</p> <p>Eicosanedioic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Eicosanedioic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>Eicosanedioic acid-d4</p> <p style="text-align: right;">Cat. No.: HY-W034595S</p> <p>Eicosanedioic acid-d4 is the deuterium labeled Eicosanedioic acid. Eicosanedioic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>endo-BCN-PEG4-Val-Cit-PAB-MMAE</p> <p style="text-align: right;">Cat. No.: HY-141155</p> <p>endo-BCN-PEG4-Val-Cit-PAB-MMAE is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg, 10 mg</p>
<p>Ethyl azetidine-3-carboxylate hydrochloride</p> <p style="text-align: right;">Cat. No.: HY-W052600</p> <p>Ethyl azetidine-3-carboxylate hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Ethyl azetidine-3-carboxylate hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs^{4/5}.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg</p>	<p>Fluorescein-DBCO</p> <p style="text-align: right;">Cat. No.: HY-126851</p> <p>Fluorescein-DBCO is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Fmoc-3VVD-OH</p> <p style="text-align: right;">Cat. No.: HY-78921</p> <p>Fmoc-3VVD-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.61% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 10 mg, 25 mg, 50 mg, 100 mg</p>	<p>Fmoc-8-amino-3,6-dioxaoctanoic acid (Fmoc-NH-PEG2-CH2COOH)</p> <p style="text-align: right;">Cat. No.: HY-W007713</p> <p>Fmoc-8-amino-3,6-dioxaoctanoic acid (Fmoc-NH-PEG2-CH2COOH) is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-8-amino-3,6-dioxaoctanoic acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: 99.65% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>
<p>Fmoc-Ala-Ala-Asn(Trt)-OH</p> <p style="text-align: right;">Cat. No.: HY-130933</p> <p>Fmoc-Ala-Ala-Asn(Trt)-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 5 mg, 10 mg</p>	<p>Fmoc-Ala-Ala-Asn-PABC-PNP</p> <p style="text-align: right;">Cat. No.: HY-129361</p> <p>Fmoc-Ala-Ala-Asn-PABC-PNP is a peptide cleavable ADC linker.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Fmoc-aminoxy-PEG2-NH2</p> <p style="text-align: right;">Cat. No.: HY-131955</p> <p>Fmoc-aminoxy-PEG2-NH2 is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 95.05% Clinical Data: No Development Reported Size: 250 mg, 1 g</p>	<p>Fmoc-Asp-NH2</p> <p style="text-align: right;">Cat. No.: HY-135418</p> <p>Fmoc-Asp-NH2 is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.04% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 500 mg, 1 g, 2 g</p>

<p>Fmoc-azetidine-3-carboxylic acid</p> <p>Cat. No.: HY-W011277</p> <p>Fmoc-azetidine-3-carboxylic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-azetidine-3-carboxylic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p> 	<p>Fmoc-D-Trp(Boc)-OH</p> <p>Cat. No.: HY-79129</p> <p>Fmoc-D-Trp(Boc)-OH is a cleavable ADC linker that used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: 99.13% Clinical Data: No Development Reported Size: 5 g</p> 
<p>Fmoc-D-Val-Cit-PAB</p> <p>Cat. No.: HY-19318B</p> <p>Fmoc-D-Val-Cit-PAB is a cleavable linker for antibody-drug-conjugation (ADC).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg</p> 	<p>Fmoc-D-Val-D-Cit-PAB</p> <p>Cat. No.: HY-19318C</p> <p>Fmoc-D-Val-D-Cit-PAB is a cleavable linker for antibody-drug-conjugation (ADC).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg</p> 
<p>Fmoc-Glu-(Boc)-Val-Cit-PAB-PNP</p> <p>Cat. No.: HY-136154</p> <p>Fmoc-Glu-(Boc)-Val-Cit-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>Fmoc-Gly-Gly-D-Phe-OH</p> <p>Cat. No.: HY-131833A</p> <p>Fmoc-Gly-Gly-D-Phe-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-Gly-Gly-D-Phe-OH is the D-isomer of Fmoc-Gly-Gly-Phe-OH (HY-131833).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 250 mg</p> 
<p>Fmoc-Gly-Gly-D-Phe-OtBu</p> <p>Cat. No.: HY-44234A</p> <p>Fmoc-Gly-Gly-D-Phe-OtBu is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-Gly-Gly-D-Phe-OtBu is the R-isomer of Fmoc-Gly-Gly-Phe-OtBu (HY-44234).</p> <p>Purity: 98.28% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 25 mg, 50 mg, 100 mg</p> 	<p>Fmoc-Gly-Gly-OH</p> <p>Cat. No.: HY-W023121</p> <p>Fmoc-Gly-Gly-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: 99.81% Clinical Data: No Development Reported Size: 500 mg</p> 
<p>Fmoc-Gly-Gly-Phe-OH</p> <p>Cat. No.: HY-131833</p> <p>Fmoc-Gly-Gly-Phe-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: 99.30% Clinical Data: No Development Reported Size: 500 mg</p> 	<p>Fmoc-Gly-Gly-Phe-OtBu</p> <p>Cat. No.: HY-44234</p> <p>Fmoc-Gly-Gly-Phe-OtBu is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: 99.80% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 25 mg, 50 mg, 100 mg</p> 











<p>Fmoc-Gly3-Val-Cit-PAB</p> <p style="text-align: right;">Cat. No.: HY-136106</p> <p>Fmoc-Gly3-Val-Cit-PAB is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>Fmoc-Gly3-Val-Cit-PAB-PNP</p> <p style="text-align: right;">Cat. No.: HY-136108</p> <p>Fmoc-Gly3-Val-Cit-PAB-PNP is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Fmoc-Hyp(Bom)-OH</p> <p style="text-align: right;">Cat. No.: HY-79125</p> <p>Fmoc-Hyp(Bom)-OH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-Hyp(Bom)-OH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs <.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Fmoc-Lys(Pal-Glu-OtBu)-OH</p> <p style="text-align: right;">Cat. No.: HY-W045822</p> <p>Fmoc-Lys(Pal-Glu-OtBu)-OH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-Lys(Pal-Glu-OtBu)-OH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Fmoc-Lys-OH hydrochloride</p> <p style="text-align: right;">Cat. No.: HY-W010975</p> <p>Fmoc-Lys-OH hydrochloride is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-Lys-OH hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 1 g, 5 g</p>	<p>Fmoc-N-bis-PEG3-NH-Boc</p> <p style="text-align: right;">Cat. No.: HY-130941</p> <p>Fmoc-N-bis-PEG3-NH-Boc is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Fmoc-N-methyl-PEG3-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W035378</p> <p>Fmoc-N-methyl-PEG3-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-N-methyl-PEG3-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Fmoc-NH-Azide-PEG4-L-Lysine-PFP ester</p> <p style="text-align: right;">Cat. No.: HY-136155</p> <p>Fmoc-NH-Azide-PEG4-L-Lysine-PFP ester is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Fmoc-NH-ethyl-SS-propionic acid</p> <p style="text-align: right;">Cat. No.: HY-140118</p> <p>Fmoc-NH-ethyl-SS-propionic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Fmoc-NH-ethyl-SS-propionic NHS ester</p> <p style="text-align: right;">Cat. No.: HY-140119</p> <p>Fmoc-NH-ethyl-SS-propionic NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>







<p>Fmoc-NH-PEG1-CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W055861</p> <p>Fmoc-NH-PEG1-CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG1-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: 99.91% Clinical Data: No Development Reported Size: 500 mg</p>	<p>Fmoc-NH-PEG2-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W040238</p> <p>Fmoc-NH-PEG2-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG2-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 100 mg</p>
<p>Fmoc-NH-PEG3-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W040231</p> <p>Fmoc-NH-PEG3-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG3-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: 99.80% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 50 mg, 100 mg</p>	<p>Fmoc-NH-PEG4-CH2CH2COOH (Fmoc-15-amino-4,7,10,13-tetraoxapentadecanoic acid)</p> <p style="text-align: right;">Cat. No.: HY-W000434</p> <p>Fmoc-NH-PEG4-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG4-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: 99.92% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Fmoc-NH-PEG4-CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-130175</p> <p>Fmoc-NH-PEG4-CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG4-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Fmoc-NH-PEG5-CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-133062</p> <p>Fmoc-NH-PEG5-CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG5-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Fmoc-NH-PEG6-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W040246</p> <p>Fmoc-NH-PEG6-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.86% Clinical Data: No Development Reported Size: 100 mg, 500 mg, 1 g</p>	<p>Fmoc-NH-PEG6-CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-130364</p> <p>Fmoc-NH-PEG6-CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG6-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Fmoc-NH-PEG8-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W040135</p> <p>Fmoc-NH-PEG8-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Fmoc-NH-PEG8-CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-133063</p> <p>Fmoc-NH-PEG8-CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG8-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>











<p>Fmoc-NH-PEG9-CH2CH2COOH</p> <p>Cat. No.: HY-130167</p>	<p>Fmoc-PEA</p> <p>Cat. No.: HY-128929</p>
<p>Fmoc-NH-PEG9-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG9-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Fmoc-PEA (Example 1-2) is used as a cleavable linker for antibody-drug conjugates (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 25 mg, 50 mg, 100 mg</p>
<p>Fmoc-PEG3-Ala-Ala-Asn(Trt)-PAB</p> <p>Cat. No.: HY-141151</p>	<p>Fmoc-PEG4-Ala-Ala-Asn-PAB</p> <p>Cat. No.: HY-141149</p>
<p>Fmoc-PEG3-Ala-Ala-Asn(Trt)-PAB is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Fmoc-PEG4-Ala-Ala-Asn-PAB is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Fmoc-Phe-Lys(Boc)-PAB-PNP</p> <p>Cat. No.: HY-114430</p>	<p>Fmoc-Phe-Lys(Trt)-PAB</p> <p>Cat. No.: HY-136107</p>
<p>Fmoc-Phe-Lys(Boc)-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.67% Clinical Data: No Development Reported Size: 50 mg</p>	<p>Fmoc-Phe-Lys(Trt)-PAB is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Fmoc-Phe-Lys(Trt)-PAB-PNP</p> <p>Cat. No.: HY-129350</p>	<p>Fmoc-Val-Ala-PAB-OH</p> <p>Cat. No.: HY-126353</p>
<p>Fmoc-Phe-Lys(Trt)-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g</p>	<p>Fmoc-Val-Ala-PAB-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.01% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>Fmoc-Val-Ala-PAB-PNP</p> <p>Cat. No.: HY-136136</p>	<p>Fmoc-Val-Cit-PAB</p> <p>Cat. No.: HY-19318</p>
<p>Fmoc-Val-Ala-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Fmoc-Val-Cit-PAB is a cleavable linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: 97.15% Clinical Data: No Development Reported Size: 250 mg, 500 mg, 1 g</p>











<p>Fmoc-Val-Cit-PAB-PNP</p> <p>Cat. No.: HY-41189</p> <p>Fmoc-Val-Cit-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: 95.87% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 250 mg</p> 	<p>Fmoc-Val-D-Cit-PAB</p> <p>Cat. No.: HY-19318A</p> <p>Fmoc-D-Val-Cit-PAB is a cleavable linker for antibody-drug-conjugation (ADC).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg</p> 
<p>Folate-PEG3-amine</p> <p>Cat. No.: HY-138484</p> <p>Folate-PEG3-amine is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>Gly-Gly-Gly-PEG3-TCO</p> <p>Cat. No.: HY-141190</p> <p>Gly-Gly-Gly-PEG3-TCO is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p> 
<p>Gly-Gly-Gly-PEG4-azide</p> <p>Cat. No.: HY-145066</p> <p>Gly-Gly-Gly-PEG4-azide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p> 	<p>Gly-Gly-Gly-PEG4-DBCO</p> <p>Cat. No.: HY-140309</p> <p>Gly-Gly-Gly-PEG4-DBCO is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: Size: 5 mg, 10 mg</p> 
<p>Gly-Gly-Gly-PEG4-methyltetrazine</p> <p>Cat. No.: HY-141284</p> <p>Gly-Gly-Gly-PEG4-methyltetrazine is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p> 	<p>Gly-PEG3-amine</p> <p>Cat. No.: HY-140244</p> <p>Gly-PEG3-amine is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p> 
<p>H-cis-Hyp-OMe hydrochloride</p> <p>Cat. No.: HY-W016429</p> <p>H-cis-Hyp-OMe hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). H-cis-Hyp-OMe hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PR.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g, 5 g</p> <p>HCl</p> 	<p>H-Glu-OtBu</p> <p>Cat. No.: HY-W018154</p> <p>H-Glu-OtBu is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). H-Glu-OtBu is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs¹².</p> <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 1 g</p> 











<p>H-Hyp-OMe hydrochloride</p> <p>Cat. No.: HY-76043</p>	<p>Hydroxy-PEG1-acid</p> <p>Cat. No.: HY-116655</p>
<p>H-Hyp-OMe hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). H-Hyp-OMe hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs⁺.</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 g, 5 g</p>	<p>Hydroxy-PEG1-acid is a non-cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>
<p>Hydroxy-PEG10-acid (HO-PEG10-CH₂CH₂COOH)</p> <p>Cat. No.: HY-133307</p> <p>Hydroxy-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 100 mg, 250 mg</p>	<p>Hydroxy-PEG10-Boc</p> <p>Cat. No.: HY-W019939</p> <p>Hydroxy-PEG10-Boc is extracted from patent CN108707228 (example 0024). Hydroxy-PEG10-Boc is a uncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Hydroxy-PEG10-Boc can be conjugated to Paclitaxel (HY-B0015) or docetaxel (HY-B0011).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>
<p>Hydroxy-PEG2-(CH₂)₂-Boc</p> <p>Cat. No.: HY-W067061</p> <p>Hydroxy-PEG2-(CH₂)₂-Boc is a uncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Hydroxy-PEG2-(CH₂)₂-Boc is extracted from patent WO2004008101A2 (compound 196).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 10 mM × 1 mL, 50 mg, 100 mg</p>	<p>Hydroxy-PEG3-(CH₂)₂-Boc</p> <p>Cat. No.: HY-42488</p> <p>Hydroxy-PEG2-(CH₂)₂-Boc is a uncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Hydroxy-PEG2-(CH₂)₂-Boc is extracted from patent WO2004008101A2 (compound 196).</p> <p>Purity: ≥95.0%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 10 mM × 1 mL, 100 mg</p>
<p>Hydroxy-PEG3-SS-PEG3-alcohol</p> <p>Cat. No.: HY-130546</p> <p>Hydroxy-PEG3-SS-PEG3-alcohol is also a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>	<p>Hydroxy-PEG4-(CH₂)₂-Boc</p> <p>Cat. No.: HY-W039178</p> <p>Hydroxy-PEG4-(CH₂)₂-Boc is a uncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Hydroxy-PEG4-(CH₂)₂-Boc is extracted from patent WO2004008101A2 (compound 191).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 10 mM × 1 mL, 500 mg</p>
<p>Hydroxy-PEG4-acid</p> <p>Cat. No.: HY-117104</p> <p>Hydroxy-PEG4-acid is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Hydroxy-PEG4-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>	<p>Hynic-PEG3-N3</p> <p>Cat. No.: HY-130954</p> <p>Hynic-PEG3-N3 is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data:</p> <p>Size: 50 mg, 100 mg, 250 mg</p>

<p>HyNic-PEG4-alkyne</p> <p style="text-align: right;">Cat. No.: HY-136075</p> <p>HyNic-PEG4-alkyne is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>LC-PEG8-SPDP</p> <p style="text-align: right;">Cat. No.: HY-126497</p> <p>LC-PEG8-SPDP is a cleavable ADC linker used for the antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>m-C-tri(CH2-PEG1-NHS ester)</p> <p style="text-align: right;">Cat. No.: HY-44149</p> <p>m-C-tri(CH2-PEG1-NHS ester) is a non-cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>m-PEG10-acid</p> <p style="text-align: right;">Cat. No.: HY-140500</p> <p>m-PEG10-acid is a non-cleavable 10 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG10-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>m-PEG10-alcohol (Decaethylene glycol monomethyl ether)</p> <p style="text-align: right;">Cat. No.: HY-141218</p> <p>m-PEG10-alcohol (Decaethylene glycol monomethyl ether) is a non-cleavable 10 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG10-alcohol is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>m-PEG10-amine</p> <p style="text-align: right;">Cat. No.: HY-140226</p> <p>m-PEG10-amine is a non-cleavable 10 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG10-amine is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>m-PEG11-acid</p> <p style="text-align: right;">Cat. No.: HY-140501</p> <p>m-PEG11-acid is a non-cleavable 11 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG11-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>m-PEG11-Amine</p> <p style="text-align: right;">Cat. No.: HY-W040222</p> <p>m-PEG11-Amino is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG11-Amine is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>m-PEG12-amine</p> <p style="text-align: right;">Cat. No.: HY-140227</p> <p>m-PEG12-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. m-PEG12-amine is also a non-cleavable 12 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>m-PEG12-OH</p> <p style="text-align: right;">Cat. No.: HY-141220</p> <p>m-PEG12-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. m-PEG12-OH is also a non-cleavable 12 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>m-PEG2-Amine</p> <p style="text-align: right;">Cat. No.: HY-W008429</p> <p>m-PEG2-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG2-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.58% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 50 mg, 100 mg</p>	<p>m-PEG2-Tos</p> <p style="text-align: right;">Cat. No.: HY-42745</p> <p>m-PEG2-Tos is a uncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG2-Tos is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>
<p>m-PEG3-Amine</p> <p style="text-align: right;">Cat. No.: HY-W018174</p> <p>m-PEG3-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG3-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 97.63% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 50 mg, 100 mg</p>	<p>m-PEG3-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W067509</p> <p>m-PEG3-CH2CH2COOH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG3-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>m-PEG4-Amine</p> <p style="text-align: right;">Cat. No.: HY-W040214</p> <p>m-PEG4-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG4-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>m-PEG4-Boc</p> <p style="text-align: right;">Cat. No.: HY-141395</p> <p>m-PEG4-Boc is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG4-Boc is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>m-PEG4-Br</p> <p style="text-align: right;">Cat. No.: HY-130161</p> <p>m-PEG4-Br is a cleavable ADC linker used in the synthesis of antibody-drug conjugate (ADC) for Trastuzumab (HY-P9907).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>m-PEG4-Ms</p> <p style="text-align: right;">Cat. No.: HY-130457</p> <p>m-PEG4-Ms is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG4-Ms is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>m-PEG5-CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-120537</p> <p>m-PEG5-CH2COOH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG5-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>m-PEG5-Ms</p> <p style="text-align: right;">Cat. No.: HY-116186</p> <p>m-PEG5-Ms is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG5-Ms is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>m-PEG5-succinimidyl carbonate</p> <p style="text-align: right;">Cat. No.: HY-130150</p> <p>m-PEG5-succinimidyl carbonate is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG5-succinimidyl carbonate is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>m-PEG6-Amine</p> <p style="text-align: right;">Cat. No.: HY-130408</p> <p>m-PEG6-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG6-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>m-PEG6-azide</p> <p style="text-align: right;">Cat. No.: HY-115374</p> <p>m-PEG6-azide is a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>m-PEG6-CH2CH2CHO</p> <p style="text-align: right;">Cat. No.: HY-W035376</p> <p>m-PEG6-CH2CH2CHO is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG6-CH2CH2CHO is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>m-PEG6-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133066</p> <p>m-PEG6-NHS ester is a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG6-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>m-PEG6-SS-PEG6-methyl</p> <p style="text-align: right;">Cat. No.: HY-140121</p> <p>m-PEG6-SS-PEG6-methyl is a cleavable 12 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>m-PEG7-Amine</p> <p style="text-align: right;">Cat. No.: HY-120237</p> <p>m-PEG7-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG7-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>m-PEG7-CH2CH2CHO</p> <p style="text-align: right;">Cat. No.: HY-130185</p> <p>m-PEG7-CH2CH2CHO is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG7-CH2CH2CHO is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>m-PEG7-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-130151</p> <p>m-PEG7-CH2CH2COOH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG7-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>m-PEG7-Ms</p> <p style="text-align: right;">Cat. No.: HY-130528</p> <p>m-PEG7-Ms is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG7-Ms is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 

<p>m-PEG8-Amine</p> <p style="text-align: right;">Cat. No.: HY-W040236</p>	<p>m-PEG8-Ms</p> <p style="text-align: right;">Cat. No.: HY-117031</p>
<p>m-PEG8-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg</p>	<p>m-PEG8-Ms is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG8-Ms is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>m-PEG8-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-W019793</p>	<p>m-PEG9-Amine</p> <p style="text-align: right;">Cat. No.: HY-130571</p>
<p>m-PEG8-NHS ester is a non-cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>m-PEG9-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG9-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 50 mg</p>
<p>MAC glucuronide linker-1</p> <p style="text-align: right;">Cat. No.: HY-44221</p>	<p>MAC glucuronide linker-2</p> <p style="text-align: right;">Cat. No.: HY-44222</p>
<p>MAC glucuronide linker-1 is a cleavable ADC linker for antibody-drug-conjugations (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: 95.91% Clinical Data: No Development Reported Size: 100 mg, 250 mg, 500 mg</p>	<p>MAC glucuronide linker-2 is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: 99.38% Clinical Data: No Development Reported Size: 100 mg, 500 mg, 1 g</p>
<p>Mal-Ala-Ala-PAB-PNP</p> <p style="text-align: right;">Cat. No.: HY-139856</p>	<p>Mal-amido-(CH2COOH)2</p> <p style="text-align: right;">Cat. No.: HY-23642</p>
<p>Mal-Ala-Ala-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-amido-(CH2COOH)2, compound 7a, is a maleimidoethyl-containing intermediate for hydrophilic ADC linker.</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Mal-amido-PEG1-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126507</p>	<p>Mal-amido-PEG10-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126509</p>
<p>Mal-amido-PEG1-C2-NHS ester is a noncleavable ADC linker containing a maleimide group and an NHS ester. The NHS ester can be used to label the primary amines (-NH2) of proteins, amine-modified oligonucleotides, and other amine-containing molecules.</p> <p style="text-align: center;"></p> <p>Purity: 99.90% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Mal-amido-PEG10-C2-NHS ester is a noncleavable ADC linker containing a maleimide group and an NHS ester. The NHS ester can be used to label the primary amines (-NH2) of proteins, amine-modified oligonucleotides, and other amine-containing molecules.</p> <p style="text-align: center;"></p> <p>Purity: 95.23% Clinical Data: No Development Reported Size: 100 mg, 500 mg, 1 g</p>

<p>Mal-amido-PEG2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-W040289</p>	<p>Mal-amido-PEG2-Val-Cit-PAB-OH</p> <p style="text-align: right;">Cat. No.: HY-140146</p>
<p>Mal-amido-PEG2-NHS ester is a noncleavable ADC linker containing a maleimide group and an NHS ester. The NHS ester can be used to label the primary amines (-NH₂) of proteins, amine-modified oligonucleotides, and other amine-containing molecules.</p> <p style="text-align: center;"></p> <p>Purity: 98.14% Clinical Data: No Development Reported Size: 1 g, 10 g</p>	<p>Mal-amido-PEG2-Val-Cit-PAB-OH is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Mal-amido-PEG2-Val-Cit-PAB-PNP</p> <p style="text-align: right;">Cat. No.: HY-140147</p>	<p>Mal-amido-PEG3-C1-PFP ester</p> <p style="text-align: right;">Cat. No.: HY-133574</p>
<p>Mal-amido-PEG2-Val-Cit-PAB-PNP is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Mal-amido-PEG3-C1-PFP ester is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Mal-amido-PEG3-C1-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133582</p>	<p>Mal-amido-PEG5-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126508</p>
<p>Mal-amido-PEG3-C1-NHS ester is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-amido-PEG10-C2-NHS ester is a noncleavable ADC linker containing a maleimide group and an NHS ester. The NHS ester can be used to label the primary amines (-NH₂) of proteins, amine-modified oligonucleotides, and other amine-containing molecules.</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Mal-amido-PEG8-C2-acid</p> <p style="text-align: right;">Cat. No.: HY-101159</p>	<p>Mal-amido-PEG8-val-gly-PAB-OH</p> <p style="text-align: right;">Cat. No.: HY-141146</p>
<p>Mal-amido-PEG8-C2-acid (example 142) is a noncleavable ADC linker, extracted from patent US2018339985.</p> <p style="text-align: center;"></p> <p>Purity: 98.72% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 500 mg</p>	<p>Mal-amido-PEG8-val-gly-PAB-OH is a cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Mal-amido-PEG9-Val-Ala-PAB-SG3200</p> <p style="text-align: right;">Cat. No.: HY-139956</p>	<p>Mal-bis-PEG3-DBCO</p> <p style="text-align: right;">Cat. No.: HY-136087</p>
<p>Mal-amido-PEG9-Val-Ala-PAB-SG3200 is a cleavable ADC linker conjugate used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-bis-PEG3-DBCO is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>

<p>Mal-C2-Gly3-EDA</p> <p>Cat. No.: HY-126673</p>	<p>Mal-C2-NHS ester</p> <p>Cat. No.: HY-126502</p>
<p>Mal-C2-Gly3-EDA is a cleavable ADC linker containing a Maleimide group. Mal-C2-Gly3-EDA is used for making antibody-drug conjugate.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-C2-NHS ester is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 250 mg</p>
<p>Mal-CO-PEG5-NHS ester</p> <p>Cat. No.: HY-133544</p>	<p>Mal-NH-ethyl-SS-propionic acid</p> <p>Cat. No.: HY-140120</p>
<p>Mal-CO-PEG5-NHS ester is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-NH-ethyl-SS-propionic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.50% Clinical Data: Size: 25 mg, 50 mg, 100 mg</p>
<p>Mal-PEG1-acid</p> <p>Cat. No.: HY-126960</p>	<p>Mal-PEG1-NHS ester</p> <p>Cat. No.: HY-126886</p>
<p>Mal-PEG1-acid is a non-cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Mal-PEG1-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: 99.40% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>	<p>Mal-PEG1-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Mal-PEG1-NHS ester is PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: 98.41% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 250 mg</p>
<p>Mal-PEG1-Val-Cit-OH</p> <p>Cat. No.: HY-133459</p>	<p>Mal-PEG1-Val-Cit-PAB-PNP</p> <p>Cat. No.: HY-140144</p>
<p>Mal-PEG1-Val-Cit-OH is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Mal-PEG1-Val-Cit-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Mal-PEG1-Val-Cit-PABC-OH</p> <p>Cat. No.: HY-130944</p>	<p>Mal-PEG2-acid</p> <p>Cat. No.: HY-130442</p>
<p>Mal-PEG1-Val-Cit-PABC-OH is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-PEG2-acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Mal-PEG2-acid can be conjugated to Tubulysin (HY-128914) and its derivative cytotoxic molecule. Mal-PEG2-acid is also a PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>











<p>Mal-PEG2-bis-PEG3-BCN</p> <p style="text-align: right;">Cat. No.: HY-136060</p> <p>Mal-PEG2-bis-PEG3-BCN is a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Mal-PEG2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126504</p> <p>Mal-PEG2-NHS ester is a noncleavable ADC linker containing a Maleimide group, 2-unit PEG and an NHS ester.</p>  <p>Purity: 98.06% Clinical Data: No Development Reported Size: 250 mg</p>
<p>Mal-PEG2-Val-Cit-amido-PAB-OH</p> <p style="text-align: right;">Cat. No.: HY-130222</p> <p>Mal-PEG2-Val-Cit-amido-PAB-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Mal-PEG2-Val-Cit-amido-PAB-OH also can be used as a PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-PEG2-Val-Cit-PABA</p> <p style="text-align: right;">Cat. No.: HY-145489</p> <p>Mal-PEG2-Val-Cit-PABA is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Mal-PEG2-Val-Cit-PABA-PNP</p> <p style="text-align: right;">Cat. No.: HY-131156</p> <p>Mal-PEG2-Val-Cit-PABA-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-PEG2-VCP-NB</p> <p style="text-align: right;">Cat. No.: HY-130084</p> <p>Mal-PEG2-VCP-NB is a cleavable ADC linker containing a Maleimide group, 2-unit PEG and a VCP NB.</p>  <p>Purity: 95.38% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>
<p>Mal-PEG3-C1-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133581</p> <p>Mal-PEG3-C1-NHS ester is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-PEG3-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-129526</p> <p>Mal-PEG3-NHS ester is a noncleavable ADC linker containing a Maleimide group. Mal-PEG3-NHS ester is used for making antibody-drug conjugate.</p>  <p>Purity: 98.75% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>Mal-PEG4-(PEG3-DBCO)-(PEG3-TCO)</p> <p style="text-align: right;">Cat. No.: HY-136084</p> <p>Mal-PEG4-(PEG3-DBCO)-(PEG3-TCO) is a cleavable 10 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-PEG4-bis-PEG3-DBCO</p> <p style="text-align: right;">Cat. No.: HY-130971</p> <p>Mal-PEG4-bis-PEG3-DBCO is a cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>


<p>Mal-PEG4-bis-PEG3-methyltetrazine</p> <p>Cat. No.: HY-130953</p>	<p>Mal-PEG4-bis-PEG4-propargyl</p> <p>Cat. No.: HY-130973</p>
<p>Mal-PEG4-bis-PEG3-methyltetrazine is a cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Mal-PEG4-bis-PEG4-propargyl is a cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Mal-PEG4-PFP ester</p> <p>Cat. No.: HY-126506</p>	<p>Mal-PEG4-VA</p> <p>Cat. No.: HY-126669</p>
<p>Mal-PEG4-PFP ester is a noncleavable ADC linker containing a Maleimide group, 4-unit PEG and a PFP ester.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-PEG4-VA is a cleavable ADC linker containing a Maleimide group. Mal-PEG4-VA is used for making antibody-drug conjugate.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Mal-PEG4-Val-Cit-PAB</p> <p>Cat. No.: HY-126672</p>	<p>Mal-PEG4-Val-Cit-PAB-OH</p> <p>Cat. No.: HY-140143</p>
<p>Mal-PEG4-Val-Cit-PAB is a cleavable ADC linker containing a Maleimide group. Mal-PEG4-Val-Cit-PAB is used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mal-PEG4-Val-Cit-PAB-OH is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Mal-PEG4-Val-Cit-PAB-PNP</p> <p>Cat. No.: HY-140145</p>	<p>Mal-PEG4-VC-PAB-DMEA</p> <p>Cat. No.: HY-126668</p>
<p>Mal-PEG4-Val-Cit-PAB-PNP is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.05% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>	<p>Mal-PEG4-VC-PAB-DMEA is a cleavable ADC linker containing a Maleimide group. Mal-PEG4-VC-PAB-DMEA is used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 25 mg, 50 mg, 100 mg</p>
<p>Mal-PEG6-NHS ester</p> <p>Cat. No.: HY-130085</p>	<p>Mal-Ph-CONH-PEG4-NHS ester</p> <p>Cat. No.: HY-133545</p>
<p>Mal-PEG6-NHS ester is a noncleavable ADC linker containing a Maleimide group, 6-unit PEG and a NHS ester.</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>Mal-Ph-CONH-PEG4-NHS ester is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>











<p>Mal-Phe-C4-Val-Cit-PAB</p> <p>Cat. No.: HY-126671</p>	<p>Mal-Phe-C4-Val-Cit-PAB-DMEA</p> <p>Cat. No.: HY-126674</p>
<p>Mal-Phe-C4-Val-Cit-PAB is a cleavable ADC linker containing a Maleimide group. Mal-Phe-C4-Val-Cit-PAB is used for making antibody-drug conjugate.</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>	<p>Mal-Phe-C4-Val-Cit-PAB-DMEA is a cleavable ADC linker containing a Maleimide group. Mal-Phe-C4-Val-Cit-PAB-DMEA is used for making antibody-drug conjugate.</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>
<p>Mal-Sulfo-DBCO</p> <p>Cat. No.: HY-140306</p>	<p>Mal-PEG4-NHS ester</p> <p>Cat. No.: HY-126505</p>
<p>Mal-Sulfo-DBCO is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data:</p> <p>Size: 1 mg, 5 mg</p>	<p>Mal-PEG4-NHS ester is a non-cleavable ADC linker which links Quantum dots (QDs) with PEGylated liposomes.</p> <p>Purity: 99.10%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 50 mg, 100 mg, 250 mg</p>
<p>Maleimide-DOTA (Maleimido-mono-amide-DOTA)</p> <p>Cat. No.: HY-133540</p>	<p>Maleimide-PEG2-hydrazide TFA</p> <p>Cat. No.: HY-136097</p>
<p>Maleimide-DOTA is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: 99.17%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 50 mg, 100 mg</p>	<p>Maleimide-PEG2-hydrazide (TFA) is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data:</p> <p>Size: 1 mg, 5 mg</p>
<p>Maleimido-tri(ethylene glycol)-propionic acid (Mal-PEG3-acid)</p> <p>Cat. No.: HY-130426</p>	<p>MC(C5)-Val-Cit</p> <p>Cat. No.: HY-141143</p>
<p>Maleimido-tri(ethylene glycol)-propionic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: 99.14%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 100 mg, 250 mg</p>	<p>MC(C5)-Val-Cit is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>
<p>MC-AAA-NHCH2OCH2COOH</p> <p>Cat. No.: HY-132159</p>	<p>MC-Gly-Gly-Phe</p> <p>Cat. No.: HY-44235</p>
<p>MC-AAA-NHCH2OCH2COOH (compound 20) is a cleavable ADC linker that is used for making antibody-drug conjugate (ADC).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>	<p>MC-Gly-Gly-Phe is a cleavable linker used for antibody-drug conjugates (ADC).</p> <p>Purity: 96.57%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 10 mg, 50 mg, 100 mg, 500 mg</p>











<p>MC-Gly-Gly-Phe-Gly</p> <p>Cat. No.: HY-44246</p>	<p>MC-Gly-Gly-Phe-Gly-NH-CH₂-O-CH₂COOH</p> <p>Cat. No.: HY-131990</p>
<p>MC-Gly-Gly-Phe-Gly is a cleavable ADC linker used for antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>MC-Gly-Gly-Phe-Gly-NH-CH₂-O-CH₂COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 50 mg, 100 mg</p>
<p>Mc-Gly-Gly-Phe-Gly-PAB-OH (Mc-GGFG-PAB-OH)</p> <p>Cat. No.: HY-136432</p>	<p>Mc-Gly-Gly-Phe-Gly-PAB-OH TFA (Mc-GGFG-PAB-OH TFA)</p> <p>Cat. No.: HY-136432A</p>
<p>Mc-Gly-Gly-Phe-Gly-PAB-OH (Mc-GGFG-PAB-OH) is a cleavable ADC linker used for antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Mc-Gly-Gly-Phe-Gly-PAB-OH (Mc-GGFG-PAB-OH) TFA is a cleavable ADC linker used for antibody-drug conjugates (ADCs).</p>  <p>Purity: 96.81% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>Mc-Leu-Gly-Arg</p> <p>Cat. No.: HY-128927</p>	<p>Mc-O-Si(di-iso)-Cl</p> <p>Cat. No.: HY-130817</p>
<p>Mc-Leu-Gly-Arg is a cleavable ether linker for antibody-drug conjugates (ADC) design.</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Mc-O-Si(di-iso)-Cl is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs), such as Gemcitabine-O-Si(di-iso)-O-Mc (HY-130812).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>MC-PEG2-C2-NHS ester</p> <p>Cat. No.: HY-126510</p>	<p>MC-Val-Ala-OH</p> <p>Cat. No.: HY-101153</p>
<p>MC-PEG2-C2-NHS ester is a noncleavable 2-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 1 g</p>	<p>MC-Val-Ala-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.55% Clinical Data: No Development Reported Size: 100 mg</p>
<p>MC-Val-Ala-PAB-PNP</p> <p>Cat. No.: HY-135975</p>	<p>MC-Val-Cit-PAB</p> <p>Cat. No.: HY-78738</p>
<p>MC-Val-Ala-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.34% Clinical Data: No Development Reported Size: 10 mg, 50 mg</p>	<p>MC-Val-Cit-PAB is a cathepsin cleavable ADC linker that is used for making antibody-drug conjugate.</p>  <p>Purity: 99.33% Clinical Data: No Development Reported Size: 250 mg, 500 mg, 1 g, 2 g</p>

<p>Mc-Val-Cit-PAB-Cl</p> <p>Cat. No.: HY-112099</p>	<p>MC-Val-Cit-PAB-NH-C2-NH-Boc</p> <p>Cat. No.: HY-132973</p>
<p>Mc-Val-Cit-PAB-Cl is a cleavable ADC linker. Mc-Val-Cit-PAB-Cl can be used to conjugate MMAE and antibody to form antibody-MC-vc-MMAE (e.g., anti-CD22-MC-VC-PABC-MMAE with IC₅₀s of 3.3 and 0.95 nM for BJAB and WSU cell lines in cytotoxicity assay).</p> <p>Purity: ≥95.0%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 100 mg</p>	<p>MC-Val-Cit-PAB-NH-C2-NH-Boc is a cathepsin cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: ≥95.0%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 5 mg, 10 mg, 25 mg, 50 mg, 100 mg</p>
<p>Mc-Val-Cit-PABC-PNP</p> <p>Cat. No.: HY-20336</p>	<p>MC-VC-PAB-Azide</p> <p>Cat. No.: HY-136138</p>
<p>Mc-Val-Cit-PABC-PNP is a cathepsin cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: 98.80%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 10 mg, 100 mg, 500 mg, 1 g</p>	<p>MC-VC-PAB-Azide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 10 mg, 50 mg, 100 mg</p>
<p>MC-VC-PAB-NH2</p> <p>Cat. No.: HY-136132</p>	<p>Mc-Val-Ala-PAB</p> <p>Cat. No.: HY-126364</p>
<p>MC-VC-PAB-NH₂ is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>	<p>Mc-Val-Ala-PAB is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>
<p>MCC</p> <p>Cat. No.: HY-132251</p>	<p>mDPR(Boc)-Val-Cit-PAB</p> <p>Cat. No.: HY-126670</p>
<p>MCC is non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs), such as MCC-DM1.</p> <p>Purity: ≥95.0%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 5 mg, 10 mg, 50 mg, 100 mg</p>	<p>mDPR(Boc)-Val-Cit-PAB is a cleavable ADC linker used as a linker for antibody-drug conjugates (ADC).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>
<p>Me-triacetyl-β-D-glucopyranuronate-Ph-ald-NO2</p> <p>Cat. No.: HY-131086</p>	<p>Me-triacetyl-β-D-glucopyranuronate-Ph-CH2OH-Fmoc</p> <p>Cat. No.: HY-131087</p>
<p>Me-triacetyl-β-D-glucopyranuronate-Ph-ald-NO₂ is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>	<p>Me-triacetyl-β-D-glucopyranuronate-Ph-CH₂OH-Fmoc is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>

<p>Methyl 1-Boc-azetidine-3-carboxylate</p> <p>Cat. No.: HY-40151</p> <p>Methyl 1-Boc-azetidine-3-carboxylate is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Methyl 1-Boc-azetidine-3-carboxylate is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs¹.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg, 1 g</p> 	<p>Methyl 1-Cbz-azetidine-3-carboxylate</p> <p>Cat. No.: HY-W019226</p> <p>Methyl 1-Cbz-azetidine-3-carboxylate is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Methyl 1-Cbz-azetidine-3-carboxylate is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs¹.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>Methyl azetidine-3-carboxylate hydrochloride</p> <p>Cat. No.: HY-33615</p> <p>Methyl azetidine-3-carboxylate hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Methyl azetidine-3-carboxylate hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs¹.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 250 mg, 500 mg</p> 	<p>MethylCBI-azaindole-benzamide-MOM-Boc-ethylenediamine-D</p> <p>Cat. No.: HY-145488</p> <p>MethylCBI-azaindole-benzamide-MOM-Boc-ethylenediamine-D is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>Methylcyclopropene-PEG3-amine</p> <p>Cat. No.: HY-136047</p> <p>Methylcyclopropene-PEG3-amine is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>Methylcyclopropene-PEG4-NHS</p> <p>Cat. No.: HY-136048</p> <p>Methylcyclopropene-PEG4-NHS is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>Methyltetrazine-DBCO</p> <p>Cat. No.: HY-140313</p> <p>Methyltetrazine-DBCO is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>Methyltetrazine-Maleimide</p> <p>Cat. No.: HY-136104</p> <p>Methyltetrazine-Maleimide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>Methyltetrazine-PEG4-aldehyde</p> <p>Cat. No.: HY-136074</p> <p>Methyltetrazine-PEG4-aldehyde is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>Methyltetrazine-PEG4-hydrazone-DBCO</p> <p>Cat. No.: HY-136079</p> <p>Methyltetrazine-PEG4-hydrazone-DBCO is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 

<p>Methyltetrazine-PEG4-oxyamine</p> <p>Cat. No.: HY-136056</p>	<p>Methyltetrazine-PEG4-SS-NHS ester</p> <p>Cat. No.: HY-133466</p>
<p>Methyltetrazine-PEG4-oxyamine is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG4-SS-NHS ester is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Methyltetrazine-PEG4-SS-PEG4-methyltetrazine</p> <p>Cat. No.: HY-130943</p>	<p>Methyltetrazine-SS-NHS</p> <p>Cat. No.: HY-136033</p>
<p>Methyltetrazine-PEG4-SS-PEG4-methyltetrazine is a cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Methyltetrazine-SS-NHS is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Methyltetrazine-SS-PEG4-Biotin</p> <p>Cat. No.: HY-136035</p>	<p>MP-PEG4-VK(Boc)G-OSu</p> <p>Cat. No.: HY-132163</p>
<p>Methyltetrazine-SS-PEG4-Biotin is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>MP-PEG4-VK(Boc)G-OSu is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Mp-polymer ester</p> <p>Cat. No.: HY-128970</p>	<p>N,N-Bis(PEG2-alkyne)-N-amido-PEG2-thiol</p> <p>Cat. No.: HY-136130</p>
<p>Mp-polymer ester is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>N,N-Bis(PEG2-alkyne)-N-amido-PEG2-thiol is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>N,N-Bis(PEG2-N3)-N-amido-PEG2-thiol</p> <p>Cat. No.: HY-136129</p>	<p>N-(5-Hydroxypentyl)maleimide</p> <p>Cat. No.: HY-130818</p>
<p>N,N-Bis(PEG2-N3)-N-amido-PEG2-thiol is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>N-(5-Hydroxypentyl)maleimide is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs), such as Gemcitabine-O-Si(di-iso)-O-Mc (HY-130812).</p>  <p>Purity: 99.66% Clinical Data: No Development Reported Size: 100 mg, 500 mg, 1 g</p>


<p>N-(Iodoacetamido)-Doxorubicin</p> <p>Cat. No.: HY-141158</p> <p>N-(Iodoacetamido)-Doxorubicin is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>N-Boc-cis-4-Hydroxy-D-proline</p> <p>Cat. No.: HY-W002887</p> <p>N-Boc-cis-4-Hydroxy-D-proline is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-Boc-cis-4-Hydroxy-D-proline is also a alkyl chain-based PROTAC linker that can be used in the synth.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g, 5 g</p>
<p>N-Boc-cis-4-hydroxy-D-proline methyl ester</p> <p>Cat. No.: HY-W002680</p> <p>N-Boc-cis-4-hydroxy-D-proline methyl ester is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-Boc-cis-4-hydroxy-D-proline methyl ester is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs¹.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g</p>	<p>N-Boc-cis-4-hydroxy-L-proline</p> <p>Cat. No.: HY-W002886</p> <p>N-Boc-cis-4-hydroxy-L-proline is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-Boc-cis-4-hydroxy-L-proline is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs¹².</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 500 mg</p>
<p>N-Boc-cis-4-hydroxy-L-proline methyl ester</p> <p>Cat. No.: HY-Y0755</p> <p>N-Boc-cis-4-hydroxy-L-proline methyl ester is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-Boc-cis-4-hydroxy-L-proline methyl ester is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs¹².</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>	<p>N-Boc-diethanolamine</p> <p>Cat. No.: HY-W044078</p> <p>N-Boc-diethanolamine is an Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-diethanolamine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg</p>
<p>N-Boc-N-bis(PEG2-OH)</p> <p>Cat. No.: HY-117079</p> <p>N-Boc-N-bis(PEG2-OH) is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-N-bis(PEG2-OH) is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>N-Boc-N-bis(PEG4-OH)</p> <p>Cat. No.: HY-130449</p> <p>N-Boc-N-bis(PEG4-OH) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-N-bis(PEG4-OH) is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>N-Boc-PEG2-bromide</p> <p>Cat. No.: HY-130503</p> <p>N-Boc-PEG2-bromide is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG2-bromide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>N-Boc-PEG3-bromide</p> <p>Cat. No.: HY-W006445</p> <p>N-Boc-PEG3-bromide is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG3-bromide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>N-Boc-PEG4-bromide</p> <p style="text-align: right;">Cat. No.: HY-W046471</p> <p>N-Boc-PEG4-bromide is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG4-bromide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 250 mg</p>	<p>N-Boc-PEG5-bromide</p> <p style="text-align: right;">Cat. No.: HY-120702</p> <p>N-Boc-PEG5-bromide is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG5-bromide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>N-Boc-PEG6-alcohol</p> <p style="text-align: right;">Cat. No.: HY-W071584</p> <p>N-Boc-PEG6-alcohol is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG6-alcohol is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>	<p>N-Boc-PEG7-alcohol</p> <p style="text-align: right;">Cat. No.: HY-130505</p> <p>N-Boc-PEG7-alcohol is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG7-alcohol is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>N-Boc-PEG9-alcohol</p> <p style="text-align: right;">Cat. No.: HY-W071583</p> <p>N-Boc-PEG9-alcohol is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG9-alcohol is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>N-Boc-Val-Dil-Dap-Doe</p> <p style="text-align: right;">Cat. No.: HY-130976</p> <p>N-Boc-Val-Dil-Dap-Doe is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>N-Bromoacetyl-β-alanine</p> <p style="text-align: right;">Cat. No.: HY-141379</p> <p>N-Bromoacetyl-β-alanine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs. N-Bromoacetyl-β-alanine is also a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>	<p>N-Butanoyl-L-homoserine lactone (C4-HSL; N-Butyryl-L-homoserine lactone)</p> <p style="text-align: right;">Cat. No.: HY-114816</p> <p>N-Butanoyl-L-homoserine lactone (C4-HSL) is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-Butanoyl-L-homoserine lactone has antibacterial activity and is used in antibacterial biofilm.</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>
<p>N-butyryl-L-Homoserine lactone-d5</p> <p style="text-align: right;">Cat. No.: HY-114816S</p> <p>N-butyryl-L-Homoserine lactone-d5 is the deuterium labeled N-Butanoyl-L-homoserine lactone. N-Butanoyl-L-homoserine lactone (C4-HSL) is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>N-Hydroxysulfosuccinimide sodium</p> <p style="text-align: right;">Cat. No.: HY-W002213</p> <p>N-Hydroxysulfosuccinimide (sodium) is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 10 mM × 1 mL, 500 mg</p>

N-Succinimidyl 3-(Bromoacetamido)propionate (3-(2-Bromoacetamido)propanoic acid NHS ester)

Cat. No.: HY-141385

N-Succinimidyl 3-(Bromoacetamido)propionate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. N-Succinimidyl 3-(Bromoacetamido)propionate is also a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




Purity: >98%
Clinical Data: No Development Reported
Size: 1 mg, 5 mg

N-Succinimidyl oxycarbonylpropyl methanethiosulfonate (NHS-C4-MTS)

Cat. No.: HY-130112

N-Succinimidyl oxycarbonylpropyl methanethiosulfonate is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




Purity: >98%
Clinical Data: No Development Reported
Size: 1 mg, 5 mg

N-tert-Butoxycarbonyl-trans-4-hydroxy-D-proline

Cat. No.: HY-77593

N-tert-Butoxycarbonyl-trans-4-hydroxy-D-proline is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-tert-Butoxycarbonyl-trans-4-hydroxy-D-proline is also an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




Purity: >98%
Clinical Data: No Development Reported
Size: 1 g, 5 g

N-trifluoroacetyl-β-alanyl chloride

Cat. No.: HY-138322

N-trifluoroacetyl-β-alanyl chloride is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




Purity: >98%
Clinical Data: No Development Reported
Size: 1 mg, 5 mg

N3-C2-NHS ester

Cat. No.: HY-126520

N3-C2-NHS ester is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




Purity: >98%
Clinical Data: No Development Reported
Size: 100 mg, 500 mg, 1 g

N3-C3-NHS ester

Cat. No.: HY-126521

N3-C3-NHS ester is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




Purity: 99.46%
Clinical Data: No Development Reported
Size: 100 mg, 500 mg

N3-C4-NHS ester

Cat. No.: HY-126522

N3-C4-NHS ester is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




Purity: >98%
Clinical Data: No Development Reported
Size: 100 mg, 500 mg

N3-C5-NHS ester

Cat. No.: HY-126523

N3-C5-NHS ester is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




Purity: 98.04%
Clinical Data: No Development Reported
Size: 100 mg, 500 mg

N3-PEG2-C2-NHS ester

Cat. No.: HY-126526

N3-PEG2-C2-NHS ester is a noncleavable 2-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).



Purity: ≥98.0%
Clinical Data: No Development Reported
Size: 100 mg











N3-PEG2-C2-PFP ester











Cat. No.: HY-126527

N3-PEG2-C2-PFP ester is a noncleavable 2-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).













Purity: >98%
Clinical Data: No Development Reported
Size: 1 mg, 5 mg











<p>N3-PEG3-C2-NHS ester</p> <p>Cat. No.: HY-126528</p>	<p>N3-PEG3-C2-PFP ester</p> <p>Cat. No.: HY-126529</p>
<p>N3-PEG3-C2-NHS ester is a noncleavable 3-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 500 mg, 1 g</p>	<p>N3-PEG3-C2-PFP ester is a noncleavable 3-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>N3-PEG3-CH2CH2-Boc</p> <p>Cat. No.: HY-42489</p>	<p>N3-PEG3-CH2CH2COOH</p> <p>Cat. No.: HY-42490</p>
<p>N3-PEG3-CH2CH2-Boc is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N3-PEG3-CH2CH2-Boc is also a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>N3-PEG3-CH2CH2COOH a PEG-based PROTAC linker can be used in the synthesis of BI-3663 (HY-111546), BI-4216 and BI-0319. Azido-PEG3-acid is also a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 50 mg, 100 mg</p>
<p>N3-PEG4-amido-Lys(Fmoc)-acid</p> <p>Cat. No.: HY-136058</p>	<p>N3-PEG4-C2-NHS ester</p> <p>Cat. No.: HY-130109</p>
<p>N3-PEG4-amido-Lys(Fmoc)-acid is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>N3-PEG4-C2-NHS ester is a noncleavable 4-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 100 mg</p>
<p>N3-PEG4-C2-Pfp ester</p> <p>Cat. No.: HY-130108</p>	<p>N3-PEG5-aldehyde</p> <p>Cat. No.: HY-136054</p>
<p>N3-PEG4-C2-Pfp ester is a noncleavable 4-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>N3-PEG5-aldehyde is a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>N3-Ph-NHS ester</p> <p>Cat. No.: HY-126524</p>	<p>NH-bis(C1-Boc)</p> <p>Cat. No.: HY-23641</p>
<p>N3-Ph-NHS ester is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.55% Clinical Data: No Development Reported Size: 100 mg</p>	<p>NH-bis(C1-Boc) is a noncleavable linker used for antibody-drug conjugates (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>



<p>NH-bis-PEG2</p> <p style="text-align: right;">Cat. No.: HY-130328</p>	<p>NH2-C5-PEG4-N3-L-Lysine-PEG3-N3</p> <p style="text-align: right;">Cat. No.: HY-130946</p>
<p>NH-bis-PEG2 is a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). NH-bis-PEG2 is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>NH2-C5-PEG4-N3-L-Lysine-PEG3-N3 is a cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>NH2-PEG2-C2-Boc</p> <p style="text-align: right;">Cat. No.: HY-42149</p>	<p>NH2-PEG4-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W021787</p>
<p>NH2-PEG2-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. NH2-PEG2-C2-Boc is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg</p>	<p>NH2-PEG4-CH2CH2COOH is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). NH2-PEG4-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>
<p>NH2-PEG4-hydrazone-DBCO</p> <p style="text-align: right;">Cat. No.: HY-136131</p>	<p>NH2-PEG4-Lys(Boc)-NH-(m-PEG24)</p> <p style="text-align: right;">Cat. No.: HY-140242</p>
<p>NH2-PEG4-hydrazone-DBCO is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 25 mg, 50 mg</p>	<p>NH2-PEG4-Lys(Boc)-NH-(m-PEG24) is a cleavable 28 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>NH2-PEG5-OH</p> <p style="text-align: right;">Cat. No.: HY-129637</p>	<p>NH2-PEG6-Boc</p> <p style="text-align: right;">Cat. No.: HY-130486</p>
<p>NH2-PEG5-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. NH2-PEG5-OH is also a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>NH2-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. NH2-PEG6-Boc is also a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 10 mg, 50 mg, 100 mg</p>
<p>NH2-PEG6-CH2CH2COOH</p> <p style="text-align: right;">Cat. No.: HY-W040257</p>	<p>NH2-PEG9-acid</p> <p style="text-align: right;">Cat. No.: HY-W019798</p>
<p>NH2-PEG6-CH2CH2COOH is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). NH2-PEG6-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>NH2-PEG9-acid is a non-cleavable 9 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). NH2-PEG9-acid also is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>











<p>NHPI-PEG2-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130095</p>	<p>NHPI-PEG2-C2-Pfp ester</p> <p style="text-align: right;">Cat. No.: HY-130094</p>
<p>NHPI-PEG2-C2-NHS ester is a noncleavable 2-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>NHPI-PEG2-C2-Pfp ester is a noncleavable 2-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>
<p>NHPI-PEG3-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130093</p>	<p>NHPI-PEG3-C2-Pfp ester</p> <p style="text-align: right;">Cat. No.: HY-130092</p>
<p>NHPI-PEG3-C2-NHS ester is a noncleavable 3-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>	<p>NHPI-PEG3-C2-Pfp ester is a noncleavable 3-unit PEG linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>NHPI-PEG4-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130091</p>	<p>NHPI-PEG4-C2-Pfp ester</p> <p style="text-align: right;">Cat. No.: HY-130090</p>
<p>NHPI-PEG4-C2-NHS ester, example 40 (WO2014185985A1), is used as a linker for antibody-drug conjugates (ADC).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg</p>	<p>NHPI-PEG4-C2-Pfp ester is used as a linker for antibody-drug conjugates (ADC).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>NHS-PEG2-SS-PEG2-NHS</p> <p style="text-align: right;">Cat. No.: HY-136133</p>	<p>NHS-SS-biotin</p> <p style="text-align: right;">Cat. No.: HY-140129</p>
<p>NHS-PEG2-SS-PEG2-NHS is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>	<p>NHS-SS-biotin is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>NO2-SPDB-sulfo</p> <p style="text-align: right;">Cat. No.: HY-133548</p>	<p>NO2-SPDMV</p> <p style="text-align: right;">Cat. No.: HY-W071007</p>
<p>NO2-SPDB-sulfo is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>NO2-SPDMV is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>











<p>NO2-SPDMV-sulfo</p> <p>Cat. No.: HY-133549</p>	<p>NO2-SPP</p> <p>Cat. No.: HY-129367</p>
<p>NO2-SPDMV-sulfo is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>NO2-SPP is a cleavable linker that is used for making antibody-drug conjugate (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>NO2-SPP-sulfo</p> <p>Cat. No.: HY-133547</p>	<p>NO2-SPP-sulfo-Me</p> <p>Cat. No.: HY-129378</p>
<p>NO2-SPP-sulfo is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>NO2-SPP-sulfo-Me is a cleavable linker that is used for making antibody-drug conjugate (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Oleoyl-Gly-Lys-N-(m-PEG11)</p> <p>Cat. No.: HY-141292</p>	<p>OPSS-PEG36-acid</p> <p>Cat. No.: HY-141355</p>
<p>Oleoyl-Gly-Lys-N-(m-PEG11) is a cleavable 11 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>OPSS-PEG36-acid is a cleavable 36 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>OPSS-Val-Cit-PAB-OH</p> <p>Cat. No.: HY-141144</p>	<p>OPSS-Val-Cit-PAB-PNP</p> <p>Cat. No.: HY-141145</p>
<p>OPSS-Val-Cit-PAB-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>OPSS-Val-Cit-PAB-PNP is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>PC Alkyne-PEG4-NHS ester</p> <p>Cat. No.: HY-140139</p>	<p>PC Biotin-PEG3-alkyne</p> <p>Cat. No.: HY-140130</p>
<p>PC Alkyne-PEG4-NHS ester is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>PC Biotin-PEG3-alkyne is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>










<p>PC Biotin-PEG3-azide</p> <p style="text-align: right;">Cat. No.: HY-140132</p> <p>PC Biotin-PEG3-azide is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 5 mg, 10 mg, 50 mg, 100 mg</p>	<p>PC Biotin-PEG3-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-140134</p> <p>PC Biotin-PEG3-NHS ester is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>PC DBCO-PEG3-biotin</p> <p style="text-align: right;">Cat. No.: HY-140136</p> <p>PC DBCO-PEG3-biotin is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>PC Mal-NHS carbonate ester</p> <p style="text-align: right;">Cat. No.: HY-140140</p> <p>PC Mal-NHS carbonate ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>PC SPDP-NHS carbonate ester</p> <p style="text-align: right;">Cat. No.: HY-140138</p> <p>PC SPDP-NHS carbonate ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>PC-Biotin-PEG4-PEG3-azide</p> <p style="text-align: right;">Cat. No.: HY-140133</p> <p>PC-Biotin-PEG4-PEG3-azide is a cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>PDB-Pfp</p> <p style="text-align: right;">Cat. No.: HY-129366</p> <p>PDB-Pfp is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 1 g</p>	<p>PDdB-Pfp</p> <p style="text-align: right;">Cat. No.: HY-129372</p> <p>PDdB-Pfp is a cleavable ADC linker used for the agents that target for the extracellular loop 1 (ECL1) of TM4SF1 (transmembrane 4 L6 family member 1).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>PDdEC-NB</p> <p style="text-align: right;">Cat. No.: HY-126519</p> <p>PDdEC-NB is a disulfide cleavable linker used for the antibody-drug conjugate (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>PDEC-NB</p> <p style="text-align: right;">Cat. No.: HY-126498</p> <p>PDEC-NB is a disulfide cleavable linker used for the antibody-drug conjugate (ADC).</p>  <p>Purity: 98.04% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>











<p>PDP-C1-Ph-Val-Cit</p> <p>Cat. No.: HY-126533</p> <p>PDP-C1-Ph-Val-Cit is a cleavable ADC linker used for antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>PDP-Pfp</p> <p>Cat. No.: HY-129359</p> <p>PDP-Pfp is a reducible ADC linker used for the agents that target for the extracellular loop 1 (ECL1) of TM4SF1 (transmembrane 4 L6 family member 1).</p>  <p>Purity: 98.66% Clinical Data: No Development Reported Size: 500 mg, 1 g</p>
<p>PEG12-Tos</p> <p>Cat. No.: HY-117050</p> <p>Tos-PEG12 is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). PEG12-Tos is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>PEG4-SPDP</p> <p>Cat. No.: HY-126496</p> <p>PEG4-SPDP is a cleavable ADC linker used for the antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Phe-Lys(Fmoc)-PAB</p> <p>Cat. No.: HY-129362</p> <p>Phe-Lys(Fmoc)-PAB is a cathepsin cleavable ADC linker used for the antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥99.0% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Phe-Lys(Trt)-PAB</p> <p>Cat. No.: HY-129349</p> <p>Phe-Lys(Trt)-PAB is a cathepsin cleavable ADC linker used for the antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>PPA</p> <p>Cat. No.: HY-141664</p> <p>PPA is an ADC linker (US20060073528A1). PPA can be used for making antibody-drug conjugate.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>PPC-NB</p> <p>Cat. No.: HY-126530</p> <p>PPC-NB is a glutathione cleavable linker used for the antibody-drug conjugate (ADC).</p>  <p>Purity: 99.90% Clinical Data: No Development Reported Size: 100 mg</p>
<p>PPC-NHS ester (2,5-Dioxopyrrolidin-1-yl 3-(pyridin-2-yl)disulfanyl)butanoate</p> <p>Cat. No.: HY-W071006</p> <p>PPC-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 1 g</p>	<p>Propargyl-C1-NHS ester</p> <p>Cat. No.: HY-126511</p> <p>Propargyl-C1-NHS ester is a noncleavable linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: 95.12% Clinical Data: No Development Reported Size: 100 mg, 500 mg, 1 g</p>







<p>Propargyl-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126512</p> <p>Propargyl-C2-NHS ester is a noncleavable linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: 96.60% Clinical Data: No Development Reported Size: 100 mg</p>	<p>Propargyl-C8-amido-PEG2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133539</p> <p>Propargyl-C8-amido-PEG2-NHS ester is a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Propargyl-NH-PEG3-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130931</p> <p>Propargyl-NH-PEG3-C2-NHS ester is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-O-C1-amido-PEG2-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126514</p> <p>Propargyl-O-C1-amido-PEG2-C2-NHS ester is a noncleavable 2-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Propargyl-O-C1-amido-PEG3-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-133583</p> <p>Propargyl-O-C1-amido-PEG3-C2-NHS ester is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-O-C1-amido-PEG4-C2-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126515</p> <p>Propargyl-O-C1-amido-PEG4-C2-NHS ester is a noncleavable 4-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Propargyl-PEG1-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126513</p> <p>Propargyl-PEG1-NHS ester is a noncleavable 1-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>Propargyl-PEG1-SS-alcohol</p> <p style="text-align: right;">Cat. No.: HY-140108</p> <p>Propargyl-PEG1-SS-alcohol is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Propargyl-PEG1-SS-PEG1-acid</p> <p style="text-align: right;">Cat. No.: HY-140109</p> <p>Propargyl-PEG1-SS-PEG1-acid is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Propargyl-PEG1-SS-PEG1-C2-Boc</p> <p style="text-align: right;">Cat. No.: HY-130690</p> <p>Propargyl-PEG1-SS-PEG1-C2-Boc is a Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG1-SS-PEG1-C2-Boc is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>Propargyl-PEG1-SS-PEG1-PFP ester</p> <p>Cat. No.: HY-140110</p>	<p>Propargyl-PEG1-SS-PEG1-propargyl</p> <p>Cat. No.: HY-140111</p>
<p>Propargyl-PEG1-SS-PEG1-PFP ester is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-PEG1-SS-PEG1-propargyl is a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Propargyl-PEG2-acid</p> <p>Cat. No.: HY-118764</p>	<p>Propargyl-PEG2-amine</p> <p>Cat. No.: HY-W051634</p>
<p>Propargyl-PEG2-acid is a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG2-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg, 250 mg</p>	<p>Propargyl-PEG2-amine is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG2-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Propargyl-PEG2-NHBoc</p> <p>Cat. No.: HY-118808</p>	<p>Propargyl-PEG3-acid</p> <p>Cat. No.: HY-126975</p>
<p>Propargyl-PEG2-NHBoc is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG2-NHBoc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-PEG3-acid is a non-cleavable (3 unit PEG) ADC linker and also a PEG-based PROTAC linker that can be used to synthesis 6-OHDA-PEG3-yne. 6-OHDA-PEG3-yne contains 6-OHDA (HY-B1081, HY-B1081A) and Propargyl-PEG3-acid.</p>  <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>Propargyl-PEG3-NHS ester</p> <p>Cat. No.: HY-126974</p>	<p>Propargyl-PEG4-Br</p> <p>Cat. No.: HY-130591</p>
<p>Propargyl-PEG3-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG3-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-PEG4-Br is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG4-Br is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>Propargyl-PEG4-CH2CH2-Boc</p> <p>Cat. No.: HY-130293</p>	<p>Propargyl-PEG4-hydrazide</p> <p>Cat. No.: HY-133427</p>
<p>Propargyl-PEG4-CH2CH2-Boc is a non-cleavable ADC linker that can be used to synthesize ADC inhibitors of Galectin-3. Propargyl-PEG4-CH2CH2-Boc is a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-PEG4-hydrazide is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>Propargyl-PEG4-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-126516</p> <p>Propargyl-PEG4-NHS ester is a noncleavable 4-unit PEG linker for antibody-drug-conjugation (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-PEG4-thiol</p> <p style="text-align: right;">Cat. No.: HY-116427</p> <p>Propargyl-PEG4-thiol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG4-thiol is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Propargyl-PEG4-Tos</p> <p style="text-align: right;">Cat. No.: HY-130387</p> <p>Propargyl-PEG4-Tos is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG4-Tos is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-PEG5-acid</p> <p style="text-align: right;">Cat. No.: HY-101157</p> <p>Propargyl-PEG5-acid is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG5-acid can be used to synthesize ADC inhibitors of Galectin-3. Propargyl-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 10 mg, 25 mg, 50 mg, 100 mg</p>
<p>Propargyl-PEG5-amine</p> <p style="text-align: right;">Cat. No.: HY-126976</p> <p>Propargyl-PEG5-amine is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG5-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>Propargyl-PEG5-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130388</p> <p>Propargyl-PEG5-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG5-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>Propargyl-PEG6-acid</p> <p style="text-align: right;">Cat. No.: HY-130386</p> <p>Propargyl-PEG6-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG6-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-PEG6-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130385</p> <p>Propargyl-PEG6-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG6-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Propargyl-PEG7-acid</p> <p style="text-align: right;">Cat. No.: HY-130383</p> <p>Propargyl-PEG7-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG7-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Propargyl-PEG7-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130381</p> <p>Propargyl-PEG7-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG7-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>









<p>Propargyl-PEG8-acid</p> <p style="text-align: right;">Cat. No.: HY-130379</p> <p>Propargyl-PEG8-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG8-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). The ADCs can be used in bacterial infections caused by Gram-negative bacteria.</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>Propargyl-PEG8-bromide</p> <p style="text-align: right;">Cat. No.: HY-130377</p> <p>Propargyl-PEG8-bromide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG8-bromide is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>Propargyl-PEG8-NH2</p> <p style="text-align: right;">Cat. No.: HY-130182</p> <p>Propargyl-PEG8-NH2 (compound 3b) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG8-NH2 is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>Propargyl-PEG8-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-130376</p> <p>Propargyl-PEG8-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG8-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>Propargyl-PEG9-bromide</p> <p style="text-align: right;">Cat. No.: HY-130372</p> <p>Propargyl-PEG9-bromide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG9-bromide is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>Propargyl-Tos</p> <p style="text-align: right;">Cat. No.: HY-79584</p> <p>Propargyl-Tos is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>PTAD-PEG4-alkyne</p> <p style="text-align: right;">Cat. No.: HY-136046</p> <p>PTAD-PEG4-alkyne is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>PTAD-PEG4-amine</p> <p style="text-align: right;">Cat. No.: HY-135961</p> <p>PTAD-PEG4-amine is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 
<p>PTAD-PEG4-N3</p> <p style="text-align: right;">Cat. No.: HY-130940</p> <p>PTAD-PEG4-N3 is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 	<p>Py-ds-dmBut-amido-PEG4-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-136157</p> <p>Py-ds-dmBut-amido-PEG4-NHS ester is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p> 











<p>Py-ds-Prp-Osu</p> <p>Cat. No.: HY-136102</p>	<p>S-(1-Hydroxy-2-methylpropan-2-yl) methanesulfonothioate</p> <p>Cat. No.: HY-129942</p>
<p>Py-ds-Prp-Osu is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>S-(1-Hydroxy-2-methylpropan-2-yl) methanesulfonothioate is a glutathione cleavable ADC linker used for the antibody-drug conjugates (ADCs) and refers to the Alkyl-Chain composition.</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 250 mg</p>
<p>SC-Val-Cit-PAB</p> <p>Cat. No.: HY-126667</p>	<p>SIA Crosslinker</p> <p>Cat. No.: HY-W011541</p>
<p>SC-Val-Cit-PAB is a cleavable ADC linker for antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>SIA Crosslinker is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>SMCC</p> <p>Cat. No.: HY-42360</p>	<p>SMPT</p> <p>Cat. No.: HY-126405</p>
<p>SMCC is a protein crosslinker. SMCC-conjugated antigen coupled spleen cells to induce antigen-specific immune responses.</p>  <p>Purity: 99.25% Clinical Data: No Development Reported Size: 100 mg</p>	<p>SMPT is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 98.07% Clinical Data: No Development Reported Size: 100 mg</p>
<p>SNPB</p> <p>Cat. No.: HY-129365</p>	<p>SNPB-sulfo-Me</p> <p>Cat. No.: HY-129375</p>
<p>SNPB is a cleavable linker that is used for making antibody-drug conjugate (ADC).</p>  <p>Purity: 98.01% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>	<p>SNPB-sulfo-Me is a cleavable linker that is used for making antibody-drug conjugate (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g</p>
<p>SPDB</p> <p>Cat. No.: HY-12448</p>	<p>SPDB-sulfo</p> <p>Cat. No.: HY-129370</p>
<p>SPDB is a glutathione cleavable ADC linker used for the antibody-drug conjugate (ADCs).</p>  <p>Purity: 99.30% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 50 mg, 100 mg, 200 mg</p>	<p>SPDB-sulfo is a glutathione cleavable ADC linker used for the antibody-drug conjugate (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>










<p>SPDH</p> <p style="text-align: right;">Cat. No.: HY-129374</p> <p>SPDH is a cleavable ADC linker used for diagnosis and treatment of cancer or B cell proliferative diseases.</p>  <p>Purity: 98.57% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>SPDMB</p> <p style="text-align: right;">Cat. No.: HY-129369</p> <p>SPDMB is a glutathione cleavable ADC linker used for the antibody-drug conjugate (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg, 1 g</p>
<p>SPDMV</p> <p style="text-align: right;">Cat. No.: HY-129368</p> <p>SPDMV is a glutathione cleavable ADC linker used for the antibody-drug conjugate (ADCs).</p>  <p>Purity: 95.49% Clinical Data: No Development Reported Size: 100 mg, 500 mg, 1 g</p>	<p>SPDMV-sulfo</p> <p style="text-align: right;">Cat. No.: HY-129373</p> <p>SPDMV-sulfo is a glutathione cleavable ADC linker used for the antibody-drug conjugate (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg</p>
<p>SPDP (SPDP Crosslinker)</p> <p style="text-align: right;">Cat. No.: HY-100216</p> <p>SPDP (SPDP Crosslinker) is a short-chain crosslinker for amine-to-sulfhydryl conjugation via NHS-ester and pyridyldithiol reactive groups that form cleavable (reducible) disulfide bonds with cysteine sulfhydryls.</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 10 mg, 50 mg, 100 mg, 200 mg, 500 mg</p>	<p>SPDP-C6-Gly-Leu-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-141123</p> <p>SPDP-C6-Gly-Leu-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>SPDP-PEG12-acid</p> <p style="text-align: right;">Cat. No.: HY-141353</p> <p>SPDP-PEG12-acid is a cleavable 12 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>SPDP-PEG36-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-141358</p> <p>SPDP-PEG36-NHS ester is a cleavable 36 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>SPDP-sulfo</p> <p style="text-align: right;">Cat. No.: HY-133543</p> <p>SPDP-sulfo is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>SPDV</p> <p style="text-align: right;">Cat. No.: HY-129371</p> <p>SPDV is a cleavable ADC linker used for diagnosis and treatment of cancer or B cell proliferative diseases.</p>  <p>Purity: 98.03% Clinical Data: No Development Reported Size: 100 mg, 500 mg, 1 g</p>

<p>Sulfo-SNPB</p> <p>Cat. No.: HY-129376</p> <p>Sulfo-SNPB is a cleavable linker that is used for making antibody-drug conjugate (ADC).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 g</p>	<p>sulfo-SPDB</p> <p>Cat. No.: HY-101151</p> <p>sulfo-SPDB is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: ≥95.0% Clinical Data: No Development Reported Size: 5 mg, 10 mg, 25 mg</p>
<p>Sulfo-SPDP-C6-NHS sodium</p> <p>Cat. No.: HY-126495A</p> <p>Sulfo-SPDP-C6-NHS sodium is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Sulfo-SPP</p> <p>Cat. No.: HY-129377</p> <p>Sulfo-SPP is a heterobifunctional, thiol-cleavable and membrane impermeable crosslinker.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 25 mg, 50 mg, 100 mg</p>
<p>tans-4-Hydroxy-D-proline hydrochloride</p> <p>Cat. No.: HY-W003511</p> <p>tans-4-Hydroxy-D-proline hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). tans-4-Hydroxy-D-proline hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PR.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>tans-4-Hydroxy-D-proline methyl ester hydrochloride</p> <p>Cat. No.: HY-W006629</p> <p>tans-4-Hydroxy-D-proline methyl ester hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 250 mg, 500 mg</p>
<p>TCO-PEG1-Val-Cit-OH</p> <p>Cat. No.: HY-130934</p> <p>TCO-PEG1-Val-Cit-OH is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>TCO-PEG1-Val-Cit-PABC-OH</p> <p>Cat. No.: HY-130966</p> <p>TCO-PEG1-Val-Cit-PABC-OH is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>TCO-PEG1-Val-Cit-PABC-PNP</p> <p>Cat. No.: HY-136100</p> <p>TCO-PEG1-Val-Cit-PABC-PNP is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>TCO-PEG12-NHS ester</p> <p>Cat. No.: HY-141170</p> <p>TCO-PEG12-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACS. TCO-PEG12-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>TCO-PEG3-aldehyde</p> <p style="text-align: right;">Cat. No.: HY-136077</p> <p>TCO-PEG3-aldehyde is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>TCO-PEG3-Biotin</p> <p style="text-align: right;">Cat. No.: HY-136050</p> <p>TCO-PEG3-Biotin is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>TCO-PEG3-CH2-aldehyde</p> <p style="text-align: right;">Cat. No.: HY-136076</p> <p>TCO-PEG3-CH2-aldehyde is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>TCO-PEG4-DBCO</p> <p style="text-align: right;">Cat. No.: HY-140310</p> <p>TCO-PEG4-DBCO is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. TCO-PEG4-DBCO is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>TCO-PEG4-NHS ester</p> <p style="text-align: right;">Cat. No.: HY-141167</p> <p>TCO-PEG4-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. TCO-PEG4-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 99.58% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>TCO-SS-amine</p> <p style="text-align: right;">Cat. No.: HY-136039</p> <p>TCO-SS-amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Tetraethylene glycol monotosylate (Tos-PEG4)</p> <p style="text-align: right;">Cat. No.: HY-41541</p> <p>Tetraethylene glycol monotosylate is a cleavable and acylhydrazone-based ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Tetraethylene glycol monotosylate also can be used as a PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Tetrazine-biotin</p> <p style="text-align: right;">Cat. No.: HY-136095</p> <p>Tetrazine-biotin is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 5 mg, 10 mg, 50 mg, 100 mg</p>
<p>Tetrazine-diazo-PEG4-biotin</p> <p style="text-align: right;">Cat. No.: HY-136078</p> <p>Tetrazine-diazo-PEG4-biotin is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Tetrazine-PEG4-amine hydrochloride</p> <p style="text-align: right;">Cat. No.: HY-130977</p> <p>Tetrazine-PEG4-amine (hydrochloride) is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 95.15% Clinical Data: No Development Reported Size: 100 mg, 250 mg, 500 mg</p>

<p>Tetrazine-PEG4-biotin</p> <p style="text-align: right;">Cat. No.: HY-136053</p> <p>Tetrazine-PEG4-biotin is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Tetrazine-PEG4-oxyamine hydrochloride</p> <p style="text-align: right;">Cat. No.: HY-136052</p> <p>Tetrazine-PEG4-oxyamine (hydrochloride) is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Tetrazine-PEG4-SS-NHS</p> <p style="text-align: right;">Cat. No.: HY-136040</p> <p>Tetrazine-PEG4-SS-NHS is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: 90.21% Clinical Data: Size: 10 mg</p>	<p>Tetrazine-PEG4-SS-Py</p> <p style="text-align: right;">Cat. No.: HY-130947</p> <p>Tetrazine-PEG4-SS-Py is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Tetrazine-PEG5-SS-amine</p> <p style="text-align: right;">Cat. No.: HY-130945</p> <p>Tetrazine-PEG5-SS-amine is a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Tetrazine-PEG6-amine hydrochloride</p> <p style="text-align: right;">Cat. No.: HY-136086</p> <p>Tetrazine-PEG6-amine (hydrochloride) is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Tetrazine-Ph-OPSS</p> <p style="text-align: right;">Cat. No.: HY-130928</p> <p>Tetrazine-Ph-OPSS is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Tetrazine-Ph-SS-amine</p> <p style="text-align: right;">Cat. No.: HY-133504</p> <p>Tetrazine-Ph-SS-amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>Tetrazine-SS-Biotin</p> <p style="text-align: right;">Cat. No.: HY-136031</p> <p>Tetrazine-SS-Biotin is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Tetrazine-SS-NHS</p> <p style="text-align: right;">Cat. No.: HY-136032</p> <p>Tetrazine-SS-NHS is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>

<p>Tetrazine-SS-PEG4-Biotin</p> <p style="text-align: right;">Cat. No.: HY-136036</p> <p>Tetrazine-SS-PEG4-Biotin is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>THP-PEG6-OH</p> <p style="text-align: right;">Cat. No.: HY-126918</p> <p>THP-PEG6-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. THP-PEG6-OH is also a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>THP-SS-alcohol</p> <p style="text-align: right;">Cat. No.: HY-140122</p> <p>THP-SS-alcohol is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>THP-SS-PEG1-Boc</p> <p style="text-align: right;">Cat. No.: HY-140123</p> <p>THP-SS-PEG1-Boc is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>
<p>THP-SS-PEG1-Tos</p> <p style="text-align: right;">Cat. No.: HY-140124</p> <p>THP-SS-PEG1-Tos is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Tr-PEG2-OH</p> <p style="text-align: right;">Cat. No.: HY-114995</p> <p>Tr-PEG2-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Tr-PEG2-OH is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Tr-PEG3-OH</p> <p style="text-align: right;">Cat. No.: HY-120258</p> <p>Tr-PEG3-OH is a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Tr-PEG5-OH</p> <p style="text-align: right;">Cat. No.: HY-120845</p> <p>Tr-PEG5-OH is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Tr-PEG5-OH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>Tr-PEG6-OH</p> <p style="text-align: right;">Cat. No.: HY-129311</p> <p>Tr-PEG6-OH is a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Tr-PEG8-OH</p> <p style="text-align: right;">Cat. No.: HY-130165</p> <p>Tr-PEG8-OH is a non-cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Tr-PEG8-OH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>

<p>trans-Sulfo-SMCC</p> <p style="text-align: right;">Cat. No.: HY-126503</p> <p>trans-Sulfo-SMCC is a non-cleavable and membrane permeable ADC crosslinker.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 500 mg</p>	<p>Tris[[2-(tert-butoxycarbonyl)ethoxy]methyl]methylamine</p> <p style="text-align: right;">Cat. No.: HY-21577</p> <p>Tris[[2-(tert-butoxycarbonyl)ethoxy]methyl]methylamine is a cleavable PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-Tri-(t-butoxycarbonyloxyethyl)-methane is also a PEG/Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: ≥97.0% Clinical Data: No Development Reported Size: 50 mg</p>
<p>Val-Cit</p> <p style="text-align: right;">Cat. No.: HY-140014</p> <p>Val-Cit is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Val-cit-PAB-OH</p> <p style="text-align: right;">Cat. No.: HY-12362</p> <p>Val-cit-PAB-OH is a cleavable ADC linker.</p>  <p>Purity: 99.62% Clinical Data: No Development Reported Size: 500 mg, 1 g, 5 g</p>
<p>Vipivotide tetraxetan Linker (PSMA-617 Linker)</p> <p style="text-align: right;">Cat. No.: HY-43869</p> <p>Vipivotide tetraxetan Linker (PSMA-617 Linker) is a noncleavable peptide linker for synthesis of Vipivotide tetraxetan (PSMA-617).</p>  <p>Purity: 99.88% Clinical Data: No Development Reported Size: 50 mg, 100 mg, 250 mg</p>	<p>Zuvotolimod</p> <p style="text-align: right;">Cat. No.: HY-145620</p> <p>Zuvotolimod is a myeloid cell agonist compound-linker for antibody conjugate. Zuvotolimod can be used in the research of cancer and hepatitis.</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>β-D-glucuronide-pNP-carbonate</p> <p style="text-align: right;">Cat. No.: HY-136329</p> <p>β-D-glucuronide-pNP-carbonate is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>β-D-tetraacetylgalactopyranoside-PEG1-N3</p> <p style="text-align: right;">Cat. No.: HY-136318</p> <p>β-D-tetraacetylgalactopyranoside-PEG1-N3 is a cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p>β-Estradiol-6-CMO-PEG3-biotin</p> <p style="text-align: right;">Cat. No.: HY-130929</p> <p>β-Estradiol-6-CMO-PEG3-biotin is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p>Purity: >98% Clinical Data: Size: 1 mg, 5 mg</p>	