



[www.MedChemExpress.com](http://www.MedChemExpress.com)

Inhibitors, Screening Libraries, Proteins

# PROTAC Linkers

PROTACs (Proteolysis Targeting Chimeric Molecules) are heterobifunctional protein degraders and promising targeted therapeutics candidates for cancer. The PROTAC linker connects two functional motifs of a PROTAC, a target protein binder and an E3 ligase recruiter.

The linker plays an important role in a PROTAC. The features of the linker (type, length, attachment position) can affect the formation of ligase:PROTAC:target ternary complex, resulting in influencing the efficient ubiquitination of the target protein and its ultimate degradation. The optimal lengths of the PROTAC linkers are reported varying from 12-carbon to over 20-carbon, and the commonly used linkers in the development of PROTACs are PEGs, Alkyl-Chain and Alkyl/ether.


## PROTAC Linkers Chemicals

### (+)-Biotin-ONP

((+)-Biotin 4-nitrophenyl ester)

Cat. No.: HY-130888

(+)-Biotin-ONP is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### (+)-Biotin-PEG10-OH

((+)-Biotin-PEG10-alcohol)

Cat. No.: HY-130889

(+)-Biotin-PEG10-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### (+)-Biotin-PEG12-OH

Cat. No.: HY-130890

(+)-Biotin-PEG12-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### (+)-Biotin-PEG2-hydrazide

Cat. No.: HY-130892

(+)-Biotin-PEG2-hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### (+)-Biotin-PEG6-hydrazide

Cat. No.: HY-130891

(+)-Biotin-PEG6-hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### (+)-Biotin-SLC

Cat. No.: HY-130887

(+)-Biotin-SLC is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (10-Bromodecyl)phosphonic acid

Cat. No.: HY-140330

(10-Bromodecyl)phosphonic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### (2-Pyridyldithio)-PEG2-Boc

Cat. No.: HY-141361

(2-Pyridyldithio)-PEG2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (2-Pyridyldithio)-PEG4-alcohol

Cat. No.: HY-141359

(2-Pyridyldithio)-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (2-Pyridyldithio)-PEG4-propargyl

Cat. No.: HY-141360

(2-Pyridyldithio)-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### (4-Oxo-4H-quinazolin-3-yl)-acetic acid

Cat. No.: HY-W030545

(4-Oxo-4H-quinazolin-3-yl)-acetic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (5,6)TAMRA-PEG3-Azide-PEG3-Desthiobiotin

Cat. No.: HY-140948

(5,6)TAMRA-PEG3-Azide-PEG3-Desthiobiotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (E)-TCO-PEG4-NHS ester

Cat. No.: HY-141167A

(E)-TCO-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (R)-Azetidine-2-carboxylic acid

Cat. No.: HY-W017755

(R)-Azetidine-2-carboxylic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). (R)-Azetidine-2-carboxylic acid is also an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 100 mg

### (R)-TCO-OH

Cat. No.: HY-141186A

(R)-TCO-OH is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (S)-TCO-PEG3-amine

Cat. No.: HY-141178A

(S)-TCO-PEG3-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (S)-TCO-PEG4-acid

Cat. No.: HY-141159A

(S)-TCO-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (S,R,S)-AHPC-PEG2-acid

Cat. No.: HY-141016

(S,R,S)-AHPC-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### (±)15-HETE

Cat. No.: HY-113336B

(±)15-HETE is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** ≥99.0%  
**Clinical Data:** No Development Reported  
**Size:** 25 µg (312.04 µM \* 250 µL in Ethanol)











### 1,1,1-Trifluoroethyl-PEG2-azide











Cat. No.: HY-140845











1,1,1-Trifluoroethyl-PEG2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.













**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

<p><b>1,1,1-Trifluoroethyl-PEG2-propargyl</b></p> <p>Cat. No.: HY-140886</p>	<p><b>1,1,1-Trifluoroethyl-PEG4-alcohol</b></p> <p>Cat. No.: HY-141247</p>
<p>1,1,1-Trifluoroethyl-PEG2-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>1,1,1-Trifluoroethyl-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>1,1,1-Trifluoroethyl-PEG4-amine</b></p> <p>Cat. No.: HY-140243</p>	<p><b>1,1,1-Trifluoroethyl-PEG4-aminoxy</b></p> <p>Cat. No.: HY-140445</p>
<p>1,1,1-Trifluoroethyl-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>1,1,1-Trifluoroethyl-PEG4-aminoxy is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>1,1,1-Trifluoroethyl-PEG4-azide</b></p> <p>Cat. No.: HY-130543</p>	<p><b>1,1,1-Trifluoroethyl-PEG4-propargyl</b></p> <p>Cat. No.: HY-140887</p>
<p>1,1,1-Trifluoroethyl-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>1,1,1-Trifluoroethyl-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>1,1,1-Trifluoroethyl-PEG4-Tos</b></p> <p>Cat. No.: HY-140379</p>	<p><b>1,2-Bis(2-iodoethoxy)ethane</b></p> <p>Cat. No.: HY-133143</p>
<p>1,1,1-Trifluoroethyl-PEG4-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>1,2-Bis(2-iodoethoxy)ethane is a PEG-based PROTAC linker. 1,2-Bis(2-iodoethoxy)ethane can be used in the synthesis of MT802 (HY-122562) and SJF620 (HY-133137). MT-802 and SJF620 are potent PROTAC BTK degraders with <math>DC_{50}</math>s of 1 nM and 7.9 nM, respectively.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>1,3-bis(carboxyethoxy)-2,2-bis(carboxyethoxy)propane</b></p> <p>Cat. No.: HY-140527</p>	<p><b>1,3-Bis-aminoxy propane</b></p> <p>Cat. No.: HY-116924</p>
<p>1,3-bis(carboxyethoxy)-2,2-bis(carboxyethoxy)propane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>1,3-Bis-aminoxy propane is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>1,3-Dibromo-5,5-dimethylhydantoin</b></p> <p>Cat. No.: HY-Y0786</p>	<p><b>1,7-Bis-Boc-1,4,7-triazaheptane</b></p> <p>Cat. No.: HY-W017829</p>
<p>1,3-Dibromo-5,5-dimethylhydantoin is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 500 mg</p>	<p>1,7-Bis-Boc-1,4,7-triazaheptane is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> ≥97.0%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 100 mg</p>
	
<p><b>1-(Isopropylthio)-2,3,4,6-tetra-o-Ac-beta-D-glucosylpyranose</b></p> <p>Cat. No.: HY-141139</p>	<p><b>1-(t-Boc-Aminoxy)-3-aminoxy-propane</b></p> <p>Cat. No.: HY-140405</p>
<p>1-(Isopropylthio)-2,3,4,6-tetra-o-Ac-beta-D-glucosylpyranose is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>1-(t-Boc-Aminoxy)-3-aminoxy-propane is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
	
<p><b>1-Boc-azetidine-3-carboxylic acid</b></p> <p>Cat. No.: HY-40141</p>	<p><b>1-Boc-azetidine-3-yl-methanol</b></p> <p>Cat. No.: HY-40152</p>
<p>1-Boc-azetidine-3-carboxylic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-Boc-azetidine-3-carboxylic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>+/sup</sup>.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 500 mg, 1 g</p>	<p>1-Boc-azetidine-3-yl-methanol is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-Boc-azetidine-3-yl-methanol is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>+/sup</sup>.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 500 mg, 1 g</p>
	
<p><b>1-Bromo-6-chlorohexane</b></p> <p>Cat. No.: HY-W009787</p>	<p><b>1-Cbz-3-Hydroxyazetidine</b></p> <p>Cat. No.: HY-77475</p>
<p>1-Bromo-6-chlorohexane is a PROTAC linker can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>1-Cbz-3-Hydroxyazetidine is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-Cbz-3-Hydroxyazetidine is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 g, 5 g</p>
	
<p><b>1-Cbz-azetidine-3-carboxylic acid</b></p> <p>Cat. No.: HY-W004868</p>	<p><b>1-Isothiocyanato-PEG3-azide</b></p> <p>Cat. No.: HY-140843</p>
<p>1-Cbz-azetidine-3-carboxylic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-Cbz-azetidine-3-carboxylic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>+/sup</sup>.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 500 mg</p>	<p>1-Isothiocyanato-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
	











<b>1-Isothiocyanato-PEG4-alcohol</b> Cat. No.: HY-141246	<b>1-N-Boc-3-hydroxyazetidine</b> Cat. No.: HY-40142
<p>1-Isothiocyanato-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>1-N-Boc-3-hydroxyazetidine is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 1-N-Boc-3-hydroxyazetidine is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg, 1 g</p>
<b>11-Aminoundecanoic acid</b> Cat. No.: HY-W014831	<b>11-Maleimidoundecanoic acid</b> Cat. No.: HY-130908
<p>11-Aminoundecanoic acid is an alkyl chain based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p>11-Maleimidoundecanoic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<b>2,2-Oxybis(ethylamine)</b> Cat. No.: HY-140208	<b>2,3,4,6-Tetra-o-acetyl-alpha-galactosylpyranosyl bromide</b> Cat. No.: HY-141137
<p>2,2-Oxybis(ethylamine) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 250 mg, 500 mg</p>	<p>2,3,4,6-Tetra-o-acetyl-alpha-galactosylpyranosyl bromide is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<b>2-((Azido-PEG8-carbamoyl)methoxy)acetic acid</b> Cat. No.: HY-140459	<b>2-(2-(6-chlorohexyloxy)ethoxy)ethanamine hydrochloride</b> Cat. No.: HY-W096093
<p>2-((Azido-PEG8-carbamoyl)methoxy)acetic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg, 1 g</p>	<p>2-(2-(6-chlorohexyloxy)ethoxy)ethanamine (hydrochloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<b>2-(Azido-PEG2-amido)-1,3-propanediol</b> Cat. No.: HY-141248	<b>2-(Azido-PEG3-amido)-1,3-bis(carboxylethoxy)propane</b> Cat. No.: HY-140522
<p>2-(Azido-PEG2-amido)-1,3-propanediol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>2-Azido-PEG3-amido-1,3-bis(carboxylethoxy)propane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>


<p><b>2-(Azido-PEG3-amido)-1,3-bis(NHS ester)</b></p> <p>Cat. No.: HY-140865</p>	<p><b>2-(Benzyloxy)ethanol</b></p> <p>Cat. No.: HY-W007853</p>
<p>2-(Azido-PEG3-amido)-1,3-bis(NHS ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>2-(Benzyloxy)ethanol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.32%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>2-(Biotin-amido)-1,3-bis-(C1-PEG1-acid)</b></p> <p>Cat. No.: HY-125392</p>	<p><b>2-Amino-1,3-bis(carboxylethoxy)propane</b></p> <p>Cat. No.: HY-23212</p>
<p>2-(Biotin-amido)-13-bis(carboxylethoxy)propane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>2-Amino-13-bis(carboxylethoxy)propane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>2-Bromo-2,2-dimethyl-acetamido-PEG3-acid</b></p> <p>Cat. No.: HY-141384</p>	<p><b>2-Phthalimidehydroxy-acetic acid</b></p> <p>Cat. No.: HY-133425</p>
<p>2-Bromo-22-dimethyl-acetamido-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>2-Phthalimidehydroxy-acetic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>20-(tert-Butoxy)-20-oxoicosanoic acid</b></p> <p>Cat. No.: HY-W034597</p>	<p><b>22-(tert-Butoxy)-22-oxodocosanoic acid</b></p> <p>Cat. No.: HY-W046348</p>
<p>20-(tert-Butoxy)-20-oxoicosanoic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 20-(tert-Butoxy)-20-oxoicosanoic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs &lt;su.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg, 500 mg</p>	<p>22-(tert-Butoxy)-22-oxodocosanoic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). 22-(tert-Butoxy)-22-oxodocosanoic acid is also a alkyl chain-based PROTAC linker that can be used in t.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>3,4-Dibromo-Mal-PEG2-amine</b></p> <p>Cat. No.: HY-141004</p>	<p><b>3,4-Dibromo-Mal-PEG2-amine TFA</b></p> <p>Cat. No.: HY-141004A</p>
<p>3,4-Dibromo-Mal-PEG2-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>3,4-Dibromo-Mal-PEG2-amine TFA is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg</p>

<p><b>3,4-Dibromo-Mal-PEG2-N-Boc</b></p> <p>Cat. No.: HY-141005</p>	<p><b>3,4-Dibromo-Mal-PEG4-Boc</b></p> <p>Cat. No.: HY-141006</p>
<p>3,4-Dibromo-Mal-PEG2-N-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>3,4-Dibromo-Mal-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>3,4-Dibromo-Mal-PEG8-acid</b></p> <p>Cat. No.: HY-141003</p>	<p><b>3,4-Dibromo-Mal-PEG8-Boc</b></p> <p>Cat. No.: HY-141007</p>
<p>3,4-Dibromo-Mal-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>3,4-Dibromo-Mal-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>3,6,9-Trioxaundecanedioic Acid</b></p> <p>Cat. No.: HY-W067705</p>	<p><b>3,6-Dioxaoctanedioic acid</b></p> <p>Cat. No.: HY-W004945</p>
<p>369-Trioxaundecanedioic Acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 500 mg</p>	<p>36-Dioxaoctanedioic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 500 mg</p>
<p><b>3-(2-Pyridyldithio)propanoic Acid</b></p> <p>Cat. No.: HY-130157</p>	<p><b>3-Aminophenol-PEG4-methyl</b></p> <p>Cat. No.: HY-134742</p>
<p>3-(2-Pyridyldithio)propanoic Acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.36% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg</p>	<p>3-Aminophenol-PEG4-methyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>3-Maleimidopropionic acid</b></p> <p>Cat. No.: HY-42145</p>	<p><b>3-Mercaptopropionic acid NHS ester</b></p> <p>Cat. No.: HY-136159</p>
<p>3-Maleimidopropionic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.27% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 500 mg</p>	<p>3-Mercaptopropionyl-N-hydroxysuccinimide ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 500 mg</p>


















<p><b>4-(6-Methyl-1,2,4,5-tetrazin-3-yl)phenol</b></p> <p>Cat. No.: HY-141274</p>	<p><b>4-(N-Boc-amino)-1,6-heptanedioic acid</b></p> <p>Cat. No.: HY-140534</p>
<p>4-(6-Methyl-1,2,4,5-tetrazin-3-yl)phenol is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 250 mg, 500 mg</p>	<p>4-(N-Boc-amino)-1,6-heptanedioic acid is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>4-Boc-amino-2,2-dimethylbutyric acid</b></p> <p>Cat. No.: HY-W090446</p>	<p><b>4-Bromobutylphosphonic acid</b></p> <p>Cat. No.: HY-140328</p>
<p>4-Boc-amino-2,2-dimethylbutyric acid is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>4-Bromobutylphosphonic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>4-Maleimidobutyric acid</b></p> <p>Cat. No.: HY-W037355</p>	<p><b>5-(Biotinamido)pentylamine</b></p> <p>Cat. No.: HY-W076543</p>
<p>4-Maleimidobutyric acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 500 mg</p>	<p>5-Biotinamidopentylamine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.40%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 100 mg</p>
<p><b>5-(Biotinamido)pentylazide</b></p> <p>Cat. No.: HY-134695</p>	<p><b>5-endo-BCN-pentanoic acid</b></p> <p>Cat. No.: HY-140350</p>
<p>5-Biotinamidopentylazide is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>5-endo-BCN-pentanoic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>5-Ethynyl-2'-deoxyuridine</b></p> <p>Cat. No.: HY-118411</p>	<p><b>5-[Boc(methyl)amino]pentanal</b></p> <p>Cat. No.: HY-138520</p>
<p>5-Ethynyl-2'-deoxyuridine (EdU), a thymidine analogue, is incorporated into cellular DNA during DNA replication and the subsequent reaction of EdU with a fluorescent azide in a "Click" reaction.</p>  <p><b>Purity:</b> 99.77%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 10 mM × 1 mL, 50 mg, 100 mg, 250 mg, 500 mg</p>	<p>5-[Boc(methyl)amino]pentanal is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>

<p><b>6-Bromohexylphosphonic acid</b></p> <p>Cat. No.: HY-140329</p>	<p><b>6-Maleimidocaproic acid sulfo-NHS</b></p> <p>Cat. No.: HY-136158</p>
<p>6-Bromohexylphosphonic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p></p> <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>6-Maleimidocaproic acid sulfo-NHS is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p></p> <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>
<p><b>6-Maleimidocaproic acid-PFP ester</b></p> <p>Cat. No.: HY-140990</p>	<p><b>6-Maleimidocaproic acid</b></p> <p>Cat. No.: HY-77959</p>
<p>6-Maleimidocaproic acid-PFP ester is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p></p> <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>6-Maleimidocaproic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p></p> <p>Purity: &gt;98% Clinical Data: No Development Reported Size: 100 mg</p>
<p><b>7-O-(Amino-PEG4)-paclitaxel</b></p> <p>Cat. No.: HY-141147</p>	<p><b>7-O-(Cbz-N-amido-PEG4)-paclitaxel</b></p> <p>Cat. No.: HY-141148</p>
<p>7-O-(Amino-PEG4)-paclitaxel is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p></p> <p>Purity: 98.61% Clinical Data: No Development Reported Size: 25 mg</p>	<p>7-O-(Cbz-N-amido-PEG4)-paclitaxel is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p></p> <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>
<p><b>7-Octynoic acid</b></p> <p>Cat. No.: HY-69220</p>	<p><b>9-Decyn-1-ol</b></p> <p>Cat. No.: HY-130985</p>
<p>7-Octynoic acid (compound 42) is a PROTAC linker and can be used in the synthesis of a series of PROTACs. PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein.</p> <p></p> <p>Purity: ≥98.0% Clinical Data: No Development Reported Size: 100 mg, 500 mg</p>	<p>9-Decyn-1-ol is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs. 9-Decyn-1-ol can be used to conjugate GDC-0068 with Lenalidomide to generate INY-03-041. INY-03-041 is a potent, highly selective and PROTAC-based pan-Akt degrader.</p> <p></p> <p>Purity: &gt;98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p><b>Ac4GalNAI</b></p> <p>Cat. No.: HY-140343</p>	<p><b>Ac4GalNAz</b></p> <p>Cat. No.: HY-141128</p>
<p>Ac4GalNAI is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p></p> <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Ac4GalNAz is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p></p> <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>

<p><b>Ac4GlcNAk</b></p> <p>Cat. No.: HY-141136</p>	<p><b>Acetamido-PEG2-Br</b></p> <p>Cat. No.: HY-134701</p>
<p>Ac4GlcNAk is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Acetamido-PEG2-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acetamido-PEG3-Br</b></p> <p>Cat. No.: HY-134706</p>	<p><b>Acid-C1-PEG5-Boc</b></p> <p>Cat. No.: HY-140483</p>
<p>Acetamido-PEG3-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-C1-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-C2-PEG3-NHS ester</b></p> <p>Cat. No.: HY-140011</p>	<p><b>Acid-C2-PEG4-C2-NHS ester</b></p> <p>Cat. No.: HY-130436</p>
<p>Acid-C2-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-C2-PEG4-C2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG1-C2-Boc</b></p> <p>Cat. No.: HY-140479</p>	<p><b>Acid-PEG10-t-butyl ester</b></p> <p>Cat. No.: HY-143836</p>
<p>Acid-PEG1-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG10-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG12-CHO</b></p> <p>Cat. No.: HY-134703</p>	<p><b>Acid-PEG12-t-butyl ester</b></p> <p>Cat. No.: HY-143837</p>
<p>Acid-PEG12-CHO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG12-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Acid-PEG13-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141097</p>	<p><b>Acid-PEG14-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-143838</p>
<p>Acid-PEG13-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG14-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG2-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140480</p>	<p><b>Acid-PEG2-ethyl propionate</b></p> <p style="text-align: right;">Cat. No.: HY-W096144</p>
<p>Acid-PEG2-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 250 mg</p>	<p>Acid-PEG2-ethyl propionate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG25-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141098</p>	<p><b>Acid-PEG3-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130549</p>
<p>Acid-PEG25-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG3-C2-Boc is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs for the degradation of EGFR and inhibition of mTOR.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG3-mono-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140484</p>	<p><b>Acid-PEG3-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140511</p>
<p>Acid-PEG3-mono-methyl ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG3-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG3-SSPy</b></p> <p style="text-align: right;">Cat. No.: HY-132090</p>	<p><b>Acid-PEG4-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-23167</p>
<p>Acid-PEG3-SSPy is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG4-C2-Boc is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs for the inhibition of mTOR.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Acid-PEG4-mono-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-W039197</p>	<p><b>Acid-PEG4-S-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140589</p>
<p>Acid-PEG4-mono-methyl ester is a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG4-S-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG4-S-S-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140114</p>	<p><b>Acid-PEG5-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140481</p>
<p>Acid-PEG4-S-S-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG5-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG5-mono-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140485</p>	<p><b>Acid-PEG5-TEMPO</b></p> <p style="text-align: right;">Cat. No.: HY-140512</p>
<p>Acid-PEG5-mono-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG5-TEMPO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG6-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140482</p>	<p><b>Acid-PEG6-mono-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140486</p>
<p>Acid-PEG6-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG6-mono-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acid-PEG7-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-W190816</p>	<p><b>Acid-PEG8-S-S-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W096114</p>
<p>Acid-PEG7-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Acid-PEG8-S-S-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Acid-PEG9-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141096</p> <p>Acid-PEG9-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Acid-PEG9-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-143835</p> <p>Acid-PEG9-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acrylate-PEG-NH2 (MW 10000)</b></p> <p style="text-align: right;">Cat. No.: HY-140637</p> <p>Acrylate-PEG-NH2 (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 10000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Acrylate-PEG-NH2 (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140007</p> <p>Acrylate-PEG-NH2 (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 2000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acrylate-PEG-OH (MW 10000)</b></p> <p style="text-align: right;">Cat. No.: HY-140640</p> <p>Acrylate-PEG-OH (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 10000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Acrylate-PEG-OH (MW 3400)</b></p> <p style="text-align: right;">Cat. No.: HY-140638</p> <p>Acrylate-PEG-OH (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 3400</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Acrylate-PEG-OH (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140639</p> <p>Acrylate-PEG-OH (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Acryloyl-PEG4-OH</b></p> <p style="text-align: right;">Cat. No.: HY-138423</p> <p>Acryloyl-PEG4-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Active-mono-sulfone-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140513</p> <p>Active-mono-sulfone-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>AEEA-AEEA</b></p> <p style="text-align: right;">Cat. No.: HY-W125504</p> <p>AEEA-AEEA is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). AEEA-AEEA is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Aeide-C1-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140755</p>	<p><b>Ald-C2-PEG4-azide</b> (N3-PEG4-CH2CH2CHO)</p> <p style="text-align: right;">Cat. No.: HY-140633</p>
<p>Aeide-C1-NHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-C2-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-CH2-PEG10-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140631</p>	<p><b>Ald-CH2-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130144</p>
<p>Ald-CH2-PEG10-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-CH2-PEG3-azide is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Ald-CH2-PEG3-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-CH2-PEG3-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130770</p>	<p><b>Ald-CH2-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140630</p>
<p>Ald-CH2-PEG3-CH2-Boc is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of SGK3 kinase PROTAC degrader.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-CH2-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>ALD-CH2-PEG4-NHBOC</b></p> <p style="text-align: right;">Cat. No.: HY-138408</p>	<p><b>Ald-CH2-PEG5-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130330</p>
<p>ALD-CH2-PEG4-NHBOC is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-CH2-PEG5-Boc is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-CH2-PEG8-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140635</p>	<p><b>Ald-PEG1-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130758</p>
<p>Ald-CH2-PEG8-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-PEG1-C2-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Ald-Ph-amido-C2-PEG2-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140632</p>	<p><b>Ald-Ph-amido-C2-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130667</p>
<p>Ald-Ph-amido-C2-PEG2-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-Ph-amido-C2-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-Ph-amido-C2-PEG3-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130669</p>	<p><b>Ald-Ph-amido-PEG2-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130154</p>
<p>Ald-Ph-amido-C2-PEG3-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-Ph-amido-PEG2-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-Ph-amido-PEG2-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130202</p>	<p><b>Ald-Ph-amido-PEG24-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140622</p>
<p>Ald-Ph-amido-PEG2-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-Ph-amido-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-Ph-amido-PEG3-C2-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-130174</p>	<p><b>Ald-Ph-PEG12-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140624</p>
<p>Ald-Ph-amido-PEG3-C2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-Ph-PEG12-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-Ph-PEG2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140628</p>	<p><b>Ald-Ph-PEG2-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140636</p>
<p>Ald-Ph-PEG2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-Ph-PEG2-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>























<p><b>Ald-Ph-PEG24-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140623</p>	<p><b>Ald-Ph-PEG24-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140625</p>
<p>Ald-Ph-PEG24-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-Ph-PEG24-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-Ph-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140626</p>	<p><b>Ald-Ph-PEG4-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130527</p>
<p>Ald-Ph-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-Ph-PEG4-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-Ph-PEG5-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140629</p>	<p><b>Ald-Ph-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130496</p>
<p>Ald-Ph-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Ald-Ph-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ald-Ph-PEG6-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140627</p>	<p><b>Aldehyde-benzyl-PEG5-alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-136140</p>
<p>Ald-Ph-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Aldehyde-benzyl-PEG5-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Alkyne-ethyl-PEG1-Boc</b> (Alkyne-ethyl-PEG1-t-butyl ester)</p> <p style="text-align: right;">Cat. No.: HY-140032</p>	<p><b>Alkyne-PEG4-maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-133399</p>
<p>Alkyne-ethyl-PEG1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Alkyne-PEG4-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Alkynyl myristic acid</b></p> <p style="text-align: right;">Cat. No.: HY-140335</p>	<p><b>Alkynyl Palmitic Acid</b></p> <p style="text-align: right;">Cat. No.: HY-W040304</p>
<p>Alkynyl myristic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b>  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Alkynyl Palmitic Acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg, 10 mg</p>
<p><b>AM-Imidazole-PA-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-129968</p>	<p><b>Amine-PEG-amine (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140646</p>
<p>AM-Imidazole-PA-Boc is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTAC IRAK4 degrader-1 (HY-129966).</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>Amine-PEG-amine (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 2000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amine-PEG-amine (MW 20000)</b></p> <p style="text-align: right;">Cat. No.: HY-140649</p>	<p><b>Amine-PEG-amine (MW 3400)</b></p> <p style="text-align: right;">Cat. No.: HY-140647</p>
<p>Amine-PEG-amine (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 20000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amine-PEG-amine (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 3400</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amine-PEG-amine (MW 35000)</b></p> <p style="text-align: right;">Cat. No.: HY-140650</p>	<p><b>Amine-PEG-amine (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140648</p>
<p>Amine-PEG-amine (MW 35000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 35000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amine-PEG-amine (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 5000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 g</p>
<p><b>Amine-PEG-CH2COOH (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140643</p>	<p><b>Amine-PEG-CH2COOH (MW 3400)</b></p> <p style="text-align: right;">Cat. No.: HY-140644</p>
<p>Amine-PEG-CH2COOH (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 2000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amine-PEG-CH2COOH (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 3400</p> </div> <p><b>Purity:</b> ≥99.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>

<p><b>Amine-PEG-CH<sub>2</sub>COOH (MW 5000)</b></p> <p>Cat. No.: HY-140645</p>	<p><b>Amine-PEG-thiol (MW 2000)</b></p> <p>Cat. No.: HY-140651</p>
<p>Amine-PEG-CH<sub>2</sub>COOH (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amine-PEG-thiol (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 2000</p> <p><b>Purity:</b> ≥93.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Amine-PEG-thiol (MW 3400)</b></p> <p>Cat. No.: HY-140652</p>	<p><b>Amine-PEG-thiol (MW 5000)</b></p> <p>Cat. No.: HY-140653</p>
<p>Amine-PEG-thiol (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 3400</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Amine-PEG-thiol (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amine-PEG3-Desthiobiotin</b></p> <p>Cat. No.: HY-134721</p>	<p><b>Amine-PEG4-Desthiobiotin</b></p> <p>Cat. No.: HY-134720</p>
<p>Amine-PEG3-Desthiobiotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Amine-PEG4-Desthiobiotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amine-PEG6-thiol</b></p> <p>Cat. No.: HY-134711</p>	<p><b>Amino-PEG1-C2-acid</b></p> <p>Cat. No.: HY-140002</p>
<p>Amine-PEG6-thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG1-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg, 500 mg</p>
<p><b>Amino-PEG10-acid</b></p> <p>Cat. No.: HY-140176</p>	<p><b>Amino-PEG10-amine</b></p> <p>Cat. No.: HY-W040270</p>
<p>Amino-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG10-amine, a PEG-based PROTAC linker used to combine two mono diethylstilbestrol (DES)-based ligands, provides an alternative strategy for preparing more selective and active ER antagonists for endocrine therapy of breast cancer.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Amino-PEG10-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140196</p>	<p><b>Amino-PEG10-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133335</p>
<p>Amino-PEG10-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG10-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG10-OH</b></p> <p style="text-align: right;">Cat. No.: HY-120761</p>	<p><b>Amino-PEG11-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W190901</p>
<p>Amino-PEG10-OH is non-cleavable 10 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG10-OH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG11-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG11-amine</b></p> <p style="text-align: right;">Cat. No.: HY-130411</p>	<p><b>Amino-PEG11-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133332</p>
<p>Amino-PEG11-amine, a PEG-based (12 units) PROTAC linker used to combine two mono diethylstilbestrol (DES)-based ligands, provides an alternative strategy for preparing more selective and active ER antagonists for endocrine therapy of breast cancer.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>Amino-PEG11-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 50 mg, 100 mg</p>
<p><b>Amino-PEG11-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130298</p>	<p><b>Amino-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140177</p>
<p>Amino-PEG11-OH is non-cleavable 11 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG11-OH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG12-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140205</p>	<p><b>Amino-PEG12-amine</b> (H2N-PEG12-CH2CH2NH2)</p> <p style="text-align: right;">Cat. No.: HY-133327</p>
<p>Amino-PEG12-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG12-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>





<p><b>Amino-PEG12-Boc</b></p> <p>Cat. No.: HY-140197</p>	<p><b>Amino-PEG12-C2-azide</b></p> <p>Cat. No.: HY-138425</p>
<p>Amino-PEG12-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> <b>Size:</b> 100 mg</p>	<p>Amino-PEG12-C2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG12-CH2-Boc</b></p> <p>Cat. No.: HY-133336</p>	<p><b>Amino-PEG12-CH2COOH</b></p> <p>Cat. No.: HY-133333</p>
<p>Amino-PEG12-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG12-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG13-amine</b> (H2N-PEG13-CH2CH2NH2)</p> <p>Cat. No.: HY-133328</p>	<p><b>Amino-PEG14-acid</b></p> <p>Cat. No.: HY-140178</p>
<p>Amino-PEG13-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg</p>	<p>Amino-PEG14-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG14-alcohol</b> (H2N-PEG14-OH)</p> <p>Cat. No.: HY-133298</p>	<p><b>Amino-PEG15-amine</b> (H2N-PEG15-CH2CH2NH2)</p> <p>Cat. No.: HY-133329</p>
<p>Amino-PEG14-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG15-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG16-acid</b></p> <p>Cat. No.: HY-140179</p>	<p><b>Amino-PEG2-(CH2)3COOH</b></p> <p>Cat. No.: HY-140190</p>
<p>Amino-PEG16-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG2-(CH2)3COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>









<p><b>Amino-PEG2-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W040168</p>	<p><b>Amino-PEG2-C2-hydrazide-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140222</p>
<p>Amino-PEG2-C2-acid is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG2-C2-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Amino-PEG2-C2-hydrazide-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG2-CH2CH2-SH hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-132120</p>	<p><b>Amino-PEG2-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-W008279</p>
<p>Amino-PEG2-CH2CH2-SH (hydrochloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>H-Cl</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG2-NH-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Amino-PEG20-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140180</p>	<p><b>Amino-PEG20-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140198</p>
<p>Amino-PEG20-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG20-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG23-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140181</p>	<p><b>Amino-PEG23-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140211</p>
<p>Amino-PEG23-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG23-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG24-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140182</p>	<p><b>Amino-PEG24-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140206</p>
<p>Amino-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG24-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Amino-PEG24-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140199</p> <p>Amino-PEG24-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG24-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133337</p> <p>Amino-PEG24-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG25-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140183</p> <p>Amino-PEG25-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG27-amine</b> (H2N-PEG27-CH2CH2NH2)</p> <p style="text-align: right;">Cat. No.: HY-133330</p> <p>Amino-PEG27-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG28-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140184</p> <p>Amino-PEG28-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG3-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W040165</p> <p>Amino-PEG3-C2-acid is a cleavable PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG3-C2-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 250 mg, 500 mg</p>
<p><b>Amino-PEG3-C2-Amine</b></p> <p style="text-align: right;">Cat. No.: HY-W015088</p> <p>Amino-PEG3-C2-Amine is a PEG-based (3 units) PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.32% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 100 mg, 500 mg</p>	<p><b>Amino-PEG3-C2-Azido</b></p> <p style="text-align: right;">Cat. No.: HY-W021401</p> <p>Amino-PEG3-C2-Azido is a PEG-based PROTAC linker can be used in the synthesis of the PARP1 degrader iRucaparib-TP3 (HY-130645).</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Amino-PEG3-C2-sulfonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-140001</p> <p>Amino-PEG3-C2-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG3-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140189</p> <p>Amino-PEG3-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Amino-PEG32-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140185</p> <p>Amino-PEG32-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG36-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140186</p> <p>Amino-PEG36-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG36-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140207</p> <p>Amino-PEG36-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG36-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140200</p> <p>Amino-PEG36-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG36-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133338</p> <p>Amino-PEG36-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG36-CONH-PEG36-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140187</p> <p>Amino-PEG36-CONH-PEG36-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG4-(CH2)3CO2H</b></p> <p style="text-align: right;">Cat. No.: HY-140191</p> <p>Amino-PEG4-(CH2)3CO2H is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG4-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W008005</p> <p>Amino-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Amino-PEG4-alcohol is also a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>Amino-PEG4-benzyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140239</p> <p>Amino-PEG4-benzyl ester is a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140193</p> <p>Amino-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>



<p><b>Amino-PEG4-C1-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-42619</p>	<p><b>Amino-PEG4-C2-amine</b></p> <p style="text-align: right;">Cat. No.: HY-22335</p>
<p>Amino-PEG4-C1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg, 1 g</p>	<p>Amino-PEG4-C2-amine is a PEG-based (4 units) PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg, 250 mg</p>
<p><b>Amino-PEG4-C2-SH hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-138446</p>	<p><b>Amino-PEG4-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130524</p>
<p>Amino-PEG4-C2-SH hydrochloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Amino-PEG4-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Amino-PEG4-CH2COOH is also a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG4-hydrazide-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140223</p>	<p><b>Amino-PEG5-amine</b></p> <p style="text-align: right;">Cat. No.: HY-130447</p>
<p>Amino-PEG4-hydrazide-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG5-amine is a PEG-based (5 units) PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG5-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140194</p>	<p><b>Amino-PEG5-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-115384</p>
<p>Amino-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG5-C2-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Amino-PEG5-C2-acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG5-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130577</p>	<p><b>Amino-PEG6-acetic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W190960</p>
<p>Amino-PEG5-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>Amino-PEG6-acetic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Amino-PEG6-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-126942</p> <p>Amino-PEG6-alcohol is a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG6-alcohol is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG6-amido-C16-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140201</p> <p>Amino-PEG6-amido-C16-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG6-amido-C16-COOH</b>  (17-(Amino-PEG6-ethylcarbamoyl)heptadecanoic acid)</p> <p style="text-align: right;">Cat. No.: HY-140188</p> <p>Amino-PEG6-amido-C16-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG6-amine</b></p> <p style="text-align: right;">Cat. No.: HY-23105</p> <p>Amino-PEG6-amine is a PEG-based (6 units) PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG6-Thalidomide</b></p> <p style="text-align: right;">Cat. No.: HY-140240</p> <p>Amino-PEG6-Thalidomide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG7-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133334</p> <p>Amino-PEG7-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG7-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140209</p> <p>Amino-PEG7-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG7-C2-SH hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-138451</p> <p>Amino-PEG7-C2-SH hydrochloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG7-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132101</p> <p>Amino-PEG7-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Amino-PEG8-amine</b></p> <p style="text-align: right;">Cat. No.: HY-130659</p> <p>Amino-PEG8-amine is a PEG-based (8 units) PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Amino-PEG8-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-W019799</p>	<p><b>Amino-PEG8-hydrazide-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140224</p>
<p>Amino-PEG8-Boc is a cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-PEG8-Boc is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p>Amino-PEG8-hydrazide-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG9-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130166</p>	<p><b>Amino-PEG9-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140204</p>
<p>Amino-PEG9-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Amino-PEG9-acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mg, 25 mg, 50 mg, 100 mg</p>	<p>Amino-PEG9-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG9-amido-C16-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140202</p>	<p><b>Amino-PEG9-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140210</p>
<p>Amino-PEG9-amido-C16-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-PEG9-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-PEG9-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140195</p>	<p><b>Amino-Tri-(carboxyethoxymethyl)-methane</b></p> <p style="text-align: right;">Cat. No.: HY-117519</p>
<p>Amino-PEG9-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Amino-Tri-(carboxyethoxymethyl)-methane is a cleavable PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-Tri-(carboxyethoxymethyl)-methane is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Amino-Tri-(m-PEG4-ethoxymethyl)-methane</b></p> <p style="text-align: right;">Cat. No.: HY-140262</p>	<p><b>Aminoxy-PEG-OH (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140642</p>
<p>Amino-Tri-(m-PEG4-ethoxymethyl)-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Aminoxy-PEG-OH (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Aminoxy-PEG1-azide</b></p> <p style="text-align: right;">Cat. No.: HY-126948</p>	<p><b>Aminoxy-PEG1-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140056</p>
<p>Aminoxy-PEG1-azide is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Aminoxy-PEG1-propargyl is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Aminoxy-PEG2-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-126951</p>	<p><b>Aminoxy-PEG2-azide</b></p> <p style="text-align: right;">Cat. No.: HY-113931</p>
<p>Aminoxy-PEG2-alcohol is a non-cleavable 2 unit PEG <b>ADC linker</b> used in the synthesis of antibody-drug conjugates (ADCs). Aminoxy-PEG2-alcohol is also a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Aminoxy-PEG2-azide is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs. Aminoxy-PEG2-azide is also a non-cleavable 2 unit PEG <b>ADC linker</b> used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Aminoxy-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140005</p>	<p><b>Aminoxy-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-126949</p>
<p>Aminoxy-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Aminoxy-PEG3-azide is a non-cleavable 3 unit PEG <b>ADC linker</b> used in the synthesis of antibody-drug conjugates (ADCs). Aminoxy-C2-PEG3-azide is also a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Aminoxy-PEG3-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-140396</p>	<p><b>Aminoxy-PEG3-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140401</p>
<p>Aminoxy-PEG3-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Aminoxy-PEG3-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Aminoxy-PEG3-C2-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140406</p>	<p><b>Aminoxy-PEG3-C2-thiol</b></p> <p style="text-align: right;">Cat. No.: HY-134512</p>
<p>Aminoxy-PEG3-C2-NH-Boc is a PEG- and Alkyl/ether-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Aminoxy-PEG3-C2-thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>











<p><b>Aminoxy-PEG3-methyl ester</b></p> <p>Cat. No.: HY-140404</p>	<p><b>Aminoxy-PEG3-propargyl</b></p> <p>Cat. No.: HY-140057</p>
<p>Aminoxy-PEG3-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data:</p> <p>Size: 1 mg, 5 mg</p>	<p>Aminoxy-PEG3-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>
<p><b>Aminoxy-PEG4-acid</b></p> <p>Cat. No.: HY-140394</p>	<p><b>Aminoxy-PEG4-alcohol</b></p> <p>Cat. No.: HY-124123</p>
<p>Aminoxy-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data:</p> <p>Size: 1 mg, 5 mg</p>	<p>Aminoxy-PEG4-alcohol is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Aminoxy-PEG4-alcohol is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>
<p><b>Aminoxy-PEG4-azide</b></p> <p>Cat. No.: HY-126950</p>	<p><b>Aminoxy-PEG4-C2-Boc</b></p> <p>Cat. No.: HY-140402</p>
<p>Aminoxy-PEG4-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>	<p>Aminoxy-PEG4-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data:</p> <p>Size: 1 mg, 5 mg</p>
<p><b>Aminoxy-PEG4-CH2-Boc</b></p> <p>Cat. No.: HY-140403</p>	<p><b>Aminoxy-PEG4-propargyl</b></p> <p>Cat. No.: HY-140058</p>
<p>Aminoxy-PEG4-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data:</p> <p>Size: 1 mg, 5 mg</p>	<p>Aminoxy-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data:</p> <p>Size: 1 mg, 5 mg</p>
<p><b>Aminoxy-PEG5-azide</b></p> <p>Cat. No.: HY-130507</p>	<p><b>Aminoxy-PEG7-methane</b></p> <p>Cat. No.: HY-140399</p>
<p>Aminoxy-PEG5-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data: No Development Reported</p> <p>Size: 1 mg, 5 mg</p>	<p>Aminoxy-PEG7-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98%</p> <p>Clinical Data:</p> <p>Size: 1 mg, 5 mg</p>

<p><b>Aminoxy-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140395</p>	<p><b>Aminoxy-PEG8-methane</b></p> <p style="text-align: right;">Cat. No.: HY-140400</p>
<p>Aminoxy-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Aminoxy-PEG8-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>ANB-NOS</b></p> <p style="text-align: right;">Cat. No.: HY-140339</p>	<p><b>APN-C3-NH-Boc (tert-Butyl 3-(4-(2-cyanoethynyl)phenylcarbamoyl)propylcarbamate)</b></p> <p style="text-align: right;">Cat. No.: HY-140347</p>
<p>ANB-NOS is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>APN-C3-NH-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>APN-C3-PEG4-alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-116025</p>	<p><b>APN-C3-PEG4-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140841</p>
<p>APN-C3-PEG4-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>APN-C3-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg</p>
<p><b>APN-PEG4-PFP</b></p> <p style="text-align: right;">Cat. No.: HY-136030</p>	<p><b>APN-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140348</p>
<p>APN-PEG4-PFP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>APN-NH2 is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>ATP-PEG8-Biotin</b></p> <p style="text-align: right;">Cat. No.: HY-145249</p>	<p><b>AZD-CO-C2-Ph-amido-Ph-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140349</p>
<p>ATP-PEG8-Biotin is a PEG-based linker that incorporates ATP. ATP is a central component of energy storage and metabolism in vivo. ATP provides the metabolic energy to drive metabolic pumps and serves as a coenzyme in cells.</p>  <p><b>Purity:</b> ≥93.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>AZD-CO-C2-Ph-amido-Ph-azide is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Azetidin-3-ol hydrochloride</b></p> <p>Cat. No.: HY-40144</p>	<p><b>Azetidine-3-carboxylic acid</b></p> <p>Cat. No.: HY-Y0530</p>
<p>Azetidin-3-ol hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azetid-3-ol hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the syntheses PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 500 mg, 1 g</p>	<p>Azetidine-3-carboxylic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azetid-3-carboxylic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>12</sup>.</p> <p><b>Purity:</b> ≥97.0%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG-alcohol (MW 2000)</b></p> <p>Cat. No.: HY-140660</p>	<p><b>Azide-PEG-amine (MW 2000)</b></p> <p>Cat. No.: HY-140661</p>
<p>Azide-PEG-alcohol (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Azide-PEG-amine (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG-amine (MW 3500)</b></p> <p>Cat. No.: HY-140662</p>	<p><b>Azide-PEG-amine (MW 5000)</b></p> <p>Cat. No.: HY-140663</p>
<p>Azide-PEG-amine (MW 3500) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Azide-PEG-amine (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG-azide (MW 10000)</b></p> <p>Cat. No.: HY-140666</p>	<p><b>Azide-PEG-azide (MW 2000)</b></p> <p>Cat. No.: HY-140664</p>
<p>Azide-PEG-azide (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Azide-PEG-azide (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG-azide (MW 20000)</b></p> <p>Cat. No.: HY-140667</p>	<p><b>Azide-PEG-azide (MW 5000)</b></p> <p>Cat. No.: HY-140665</p>
<p>Azide-PEG-azide (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Azide-PEG-azide (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>











<p><b>Azide-PEG12-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140803</p> <p>Azide-PEG12-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azide-PEG12-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140355</p> <p>Azide-PEG12-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG16-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140804</p> <p>Azide-PEG16-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azide-PEG2-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-140381</p> <p>Azide-PEG2-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG3-C1-Ala</b></p> <p style="text-align: right;">Cat. No.: HY-140850</p> <p>Azide-PEG3-C1-Ala is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azide-PEG3-Desthiobiotin</b></p> <p style="text-align: right;">Cat. No.: HY-W096120</p> <p>Azide-PEG3-Desthiobiotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Azide-PEG3-L-alanine-Fmoc</b></p> <p style="text-align: right;">Cat. No.: HY-140848</p> <p>Azide-PEG3-L-alanine-Fmoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azide-PEG3-Sulfone-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140603</p> <p>Azide-PEG3-Sulfone-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG3-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140004</p> <p>Azide-PEG3-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Azide-PEG3-Tos is also a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azide-PEG4-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140351</p> <p>Azide-PEG4-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>













<p><b>Azide-PEG5-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140778</p>	<p><b>Azide-PEG6-amido-C16-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140787</p>
<p>Azide-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Azide-PEG6-amido-C16-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG6-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140353</p>	<p><b>Azide-PEG7-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140354</p>
<p>Azide-PEG6-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Azide-PEG7-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG8-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140799</p>	<p><b>Azide-PEG9-amido-C12-Boc</b> (13-(Azide-PEG9-ethylcarbamoyl)tridecanoic t-butyl ester) Cat. No.: HY-140790</p>
<p>Azide-PEG8-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Azide-PEG9-amido-C12-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG9-amido-C16-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140791</p>	<p><b>Azide-PEG9-amido-C4-Boc</b> (5-(Azide-PEG9-ethylcarbamoyl)pentanoic t-butyl ester) Cat. No.: HY-140788</p>
<p>Azide-PEG9-amido-C16-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Azide-PEG9-amido-C4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azide-PEG9-amido-C8-Boc</b> (9-(Azide-PEG9-ethylcarbamoyl)nonanoic t-butyl ester) Cat. No.: HY-140789</p>	<p><b>Azido-C1-PEG3-C3-NH2</b> Cat. No.: HY-134692</p>
<p>Azide-PEG9-amido-C8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-C1-PEG3-C3-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>




<p><b>Azido-C1-PEG4-C3-NH2</b></p> <p>Cat. No.: HY-134693</p>	<p><b>Azido-C3-UV-biotin</b> (UV Cleavable Biotin-PEG2-alkyne)</p> <p>Cat. No.: HY-140927</p>
<p>Azido-C1-PEG4-C3-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-C3-UV-biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG1</b></p> <p>Cat. No.: HY-138461</p>	<p><b>Azido-PEG1-amine</b></p> <p>Cat. No.: HY-140212</p>
<p>Azido-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.06% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 500 mg</p>	<p>Azido-PEG1-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG1-azide</b></p> <p>Cat. No.: HY-133390</p>	<p><b>Azido-PEG1-Boc</b></p> <p>Cat. No.: HY-140775</p>
<p>Azido-PEG1-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg</p>	<p>Azido-PEG1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG1-C1-Boc</b></p> <p>Cat. No.: HY-140792</p>	<p><b>Azido-PEG1-C2-acid</b></p> <p>Cat. No.: HY-140009</p>
<p>Azido-PEG1-C1-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG1-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Azido-PEG1-C2-methylamine</b></p> <p>Cat. No.: HY-140221</p>	<p><b>Azido-PEG1-CH2CO2-NHS</b></p> <p>Cat. No.: HY-140763</p>
<p>Azido-PEG1-C2-methylamine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG1-CH2CO2-NHS is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Azido-PEG1-CH2CO2H</b></p> <p style="text-align: right;">Cat. No.: HY-108369</p>	<p><b>Azido-PEG1-CH2COO-Cl</b></p> <p style="text-align: right;">Cat. No.: HY-130984</p>
<p>Azido-PEG1-CH2CO2H is a PROTAC linker, which refers to the alkyl/ether composition. Azido-PEG1-CH2CO2H can be used in the synthesis of PROTAC BRD4 Degradator-1.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG1-CH2COO-Cl (compound 43a) is an alkyl/ether-based PROTAC linker. Azido-PEG1-CH2COO-Cl can be used in the synthesis of PROTAC BRD4 Degradator-1 (HY-133131).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG1-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140851</p>	<p><b>Azido-PEG1-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140756</p>
<p>Azido-PEG1-methyl ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG1-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG1-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140771</p>	<p><b>Azido-PEG10-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140455</p>
<p>Azido-PEG1-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG10-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140801</p>	<p><b>Azido-PEG10-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140217</p>
<p>Azido-PEG10-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p>Azido-PEG10-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG10-azide</b></p> <p style="text-align: right;">Cat. No.: HY-132076</p>	<p><b>Azido-PEG10-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140781</p>
<p>Azido-PEG10-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG10-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Azido-PEG10-CH2CO2-NHS</b></p> <p style="text-align: right;">Cat. No.: HY-138350</p>	<p><b>Azido-PEG10-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138416</p>
<p>Azido-PEG10-CH2CO2-NHS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG10-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG10-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140759</p>	<p><b>Azido-PEG10-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-138739</p>
<p>Azido-PEG10-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG10-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG11-acid</b> (N3-PEG11-CH2CH2COOH)</p> <p style="text-align: right;">Cat. No.: HY-133317</p>	<p><b>Azido-PEG11-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140802</p>
<p>Azido-PEG11-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG11-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG11-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140218</p>	<p><b>Azido-PEG11-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133396</p>
<p>Azido-PEG11-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG11-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG11-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138415</p>	<p><b>Azido-PEG11-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132062</p>
<p>Azido-PEG11-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG11-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Azido-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140456</p>	<p><b>Azido-PEG12-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-138769</p>
<p>Azido-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG12-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG12-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140782</p>	<p><b>Azido-PEG12-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140760</p>
<p>Azido-PEG12-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG12-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-138741</p>	<p><b>Azido-PEG12-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138374</p>
<p>Azido-PEG12-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG12-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG13-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138435</p>	<p><b>Azido-PEG13-azide</b></p> <p style="text-align: right;">Cat. No.: HY-132059</p>
<p>Azido-PEG13-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG13-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG14-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132061</p>	<p><b>Azido-PEG15-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133397</p>
<p>Azido-PEG14-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG15-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Azido-PEG15-t-butyl ester</b></p> <p>Cat. No.: HY-140783</p>	<p><b>Azido-PEG16-acid</b> (N3-PEG16-CH2CH2COOH)</p> <p>Cat. No.: HY-133318</p>
<p>Azido-PEG15-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG16-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG16-Boc</b></p> <p>Cat. No.: HY-140784</p>	<p><b>Azido-PEG16-NHS ester</b></p> <p>Cat. No.: HY-140761</p>
<p>Azido-PEG16-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG16-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG2-alcohol</b></p> <p>Cat. No.: HY-140797</p>	<p><b>Azido-PEG2-azide</b></p> <p>Cat. No.: HY-W044155</p>
<p>Azido-PEG2-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 97.23% <b>Clinical Data:</b> <b>Size:</b> 50 mg, 100 mg</p>	<p>Azido-PEG2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.08% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 500 mg</p>
<p><b>Azido-PEG2-C1-Boc</b></p> <p>Cat. No.: HY-140793</p>	<p><b>Azido-PEG2-C2-acid</b></p> <p>Cat. No.: HY-140452</p>
<p>Azido-PEG2-C1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG2-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 100 mg, 250 mg, 500 mg</p>
<p><b>Azido-PEG2-C2-amine</b> (N3-PEG2-CH2CH2NH2)</p> <p>Cat. No.: HY-140213</p>	<p><b>Azido-PEG2-C2-Boc</b></p> <p>Cat. No.: HY-140776</p>
<p>Azido-PEG2-C2-amine (N3-PEG2-CH2CH2NH2) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Azido-PEG2-C2-amine is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 250 mg, 500 mg</p>	<p>Azido-PEG2-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Azido-PEG2-C2-sulfonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-140164</p>	<p><b>Azido-PEG2-C6-Cl</b></p> <p style="text-align: right;">Cat. No.: HY-138470</p>
<p>Azido-PEG2-C2-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG2-C6-Cl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG2-C6-OH</b></p> <p style="text-align: right;">Cat. No.: HY-138471</p>	<p><b>Azido-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-108368</p>
<p>Azido-PEG2-C6-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG2-hydrazide-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140816</p>	<p><b>Azido-PEG2-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-138740</p>
<p>Azido-PEG2-hydrazide-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG2-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG20-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140805</p>	<p><b>Azido-PEG20-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138457</p>
<p>Azido-PEG20-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG20-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG23-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140219</p>	<p><b>Azido-PEG23-C2-azide</b></p> <p style="text-align: right;">Cat. No.: HY-138358</p>
<p>Azido-PEG23-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG23-C2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Azido-PEG24-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140457</p> <p>Azido-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG24-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140806</p> <p>Azido-PEG24-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG24-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133319</p> <p>Azido-PEG24-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG24-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140762</p> <p>Azido-PEG24-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b>  <b>Size:</b> 25 mg</p>
<p><b>Azido-PEG3-Ala-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140849</p> <p>Azido-PEG3-Ala-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG3-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W016735</p> <p>Azido-PEG3-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥96.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>Azido-PEG3-aldehyde</b></p> <p style="text-align: right;">Cat. No.: HY-138518</p> <p>Azido-PEG3-aldehyde is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG3-amide-C3-triethoxysilane</b></p> <p style="text-align: right;">Cat. No.: HY-140019</p> <p>Azido-PEG3-amide-C3-triethoxysilane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG3-amino-OPSS</b></p> <p style="text-align: right;">Cat. No.: HY-138438</p> <p>Azido-PEG3-amino-OPSS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG3-aminoacetic acid-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140770</p> <p>Azido-PEG3-aminoacetic acid-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>




<p><b>Azido-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133391</p>	<p><b>Azido-PEG3-C-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140794</p>
<p>Azido-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 99.11% <b>Clinical Data:</b> <b>Size:</b> 100 mg, 500 mg</p>	<p>Azido-PEG3-C-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG3-C3-OH</b></p> <p style="text-align: right;">Cat. No.: HY-140808</p>	<p><b>Azido-PEG3-C6-Cl</b></p> <p style="text-align: right;">Cat. No.: HY-138466</p>
<p>Azido-PEG3-C3-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG3-C6-Cl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg</p>
<p><b>Azido-PEG3-CH2CO2Me</b></p> <p style="text-align: right;">Cat. No.: HY-140853</p>	<p><b>Azido-PEG3-chloroacetamide</b></p> <p style="text-align: right;">Cat. No.: HY-132085</p>
<p>Azido-PEG3-CH2CO2Me is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG3-chloroacetamide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG3-maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-140811</p>	<p><b>Azido-PEG3-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140852</p>
<p>Azido-PEG3-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Azido-PEG3-maleimide is also a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 25 mg</p>	<p>Azido-PEG3-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG3-MS</b></p> <p style="text-align: right;">Cat. No.: HY-138343</p>	<p><b>Azido-PEG3-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140764</p>
<p>Azido-PEG3-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Azido-PEG3-O-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140768</p> <p>Azido-PEG3-O-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG3-phosphonic acid ethyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140813</p> <p>Azido-PEG3-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG3-S-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140590</p> <p>Azido-PEG3-S-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG3-S-PEG4-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140591</p> <p>Azido-PEG3-S-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG3-S-PEG4-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140592</p> <p>Azido-PEG3-S-PEG4-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG3-SS-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140106</p> <p>Azido-PEG3-SS-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Azido-PEG3-Sulfone-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140602</p> <p>Azido-PEG3-Sulfone-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG3-Sulfone-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140604</p> <p>Azido-PEG3-Sulfone-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG3-Val-Cit-PAB-PNP</b></p> <p style="text-align: right;">Cat. No.: HY-140150</p> <p>Azido-PEG3-Val-Cit-PAB-PNP is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG3-Val-Cit-PAB-PNP is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg</p>	<p><b>Azido-PEG35-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140220</p> <p>Azido-PEG35-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Azido-PEG36-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140458</p>	<p><b>Azido-PEG36-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140807</p>
<p>Azido-PEG36-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG36-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG36-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140785</p>	<p><b>Azido-PEG36-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-138463</p>
<p>Azido-PEG36-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG36-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-(CH2)3-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140854</p>	<p><b>Azido-PEG4-(CH2)3OH</b></p> <p style="text-align: right;">Cat. No.: HY-140809</p>
<p>Azido-PEG4-(CH2)3-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG4-(CH2)3OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-4-nitrophenyl carbonate</b></p> <p style="text-align: right;">Cat. No.: HY-140858</p>	<p><b>Azido-PEG4-acyl chloride</b></p> <p style="text-align: right;">Cat. No.: HY-132096</p>
<p>Azido-PEG4-4-nitrophenyl carbonate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG4-acyl chloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-22340</p>	<p><b>Azido-PEG4-alpha-D-mannose</b></p> <p style="text-align: right;">Cat. No.: HY-141126</p>
<p>Azido-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p>Azido-PEG4-alpha-D-mannose is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg</p>

<p><b>Azido-PEG4-amido-PEG4-Boc</b></p> <p>Cat. No.: HY-140786</p>	<p><b>Azido-PEG4-amido-tri-(carboxyethoxymethyl)-methane</b></p> <p>Cat. No.: HY-140523</p>
<p>Azido-PEG4-amido-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG4-amido-tri-(carboxyethoxymethyl)-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-Amido-tri-(t-butoxycarbonylethoxymethyl)-methane</b></p> <p>Cat. No.: HY-140874</p>	<p><b>Azido-PEG4-Amido-Tris</b></p> <p>Cat. No.: HY-141249</p>
<p>Azido-PEG4-Amido-tri-(t-butoxycarbonylethoxymethyl)-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG4-Amido-Tris is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-azide</b></p> <p>Cat. No.: HY-138692</p>	<p><b>Azido-PEG4-beta-D-glucose</b></p> <p>Cat. No.: HY-140012</p>
<p>Azido-PEG4-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG4-beta-D-glucose is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-Boc</b></p> <p>Cat. No.: HY-140777</p>	<p><b>Azido-PEG4-C2-acid</b></p> <p>Cat. No.: HY-130653</p>
<p>Azido-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG4-C2-acid a PEG-based PROTAC linker can be used in the synthesis of vRucaparib-TP4. Azido-PEG4-C2-acid is also a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> ≥98.0%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 100 mg, 250 mg, 500 mg</p>
<p><b>Azido-PEG4-CH2-Boc</b></p> <p>Cat. No.: HY-42618</p>	<p><b>Azido-PEG4-formylhydrazine-Boc</b></p> <p>Cat. No.: HY-138388</p>
<p>Azido-PEG4-CH2-Boc is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG4-CH2-Boc is also a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Azido-PEG4-formylhydrazine-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>











<p><b>Azido-PEG4-hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-140814</p>	<p><b>Azido-PEG4-hydrazide-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140817</p>
<p>Azido-PEG4-hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b>  <b>Size:</b> 50 mg, 100 mg</p>	<p>Azido-PEG4-hydrazide-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-NHS-ester</b></p> <p style="text-align: right;">Cat. No.: HY-140765</p>	<p><b>Azido-PEG4-nitrile</b></p> <p style="text-align: right;">Cat. No.: HY-140842</p>
<p>Azido-PEG4-NHS-ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg, 25 mg, 50 mg, 100 mg</p>	<p>Azido-PEG4-nitrile is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-132050</p>	<p><b>Azido-PEG4-tetra-Ac-beta-D-glucose</b></p> <p style="text-align: right;">Cat. No.: HY-141127</p>
<p>Azido-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG4-tetra-Ac-beta-D-glucose is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-Thiol</b></p> <p style="text-align: right;">Cat. No.: HY-138525</p>	<p><b>Azido-PEG4-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138472</p>
<p>Azido-PEG4-Thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG4-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG4-Val-Cit-PAB-OH</b></p> <p style="text-align: right;">Cat. No.: HY-140149</p>	<p><b>Azido-PEG5-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130572</p>
<p>Azido-PEG4-Val-Cit-PAB-OH is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG4-Val-Cit-PAB-OH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>Azido-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs, such as the conjugate CPT-APO (CPT: Camptothecin (HY-16560)). Azido-PEG5-acid is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>




<p><b>Azido-PEG5-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-130211</p>	<p><b>Azido-PEG5-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140214</p>
<p>Azido-PEG5-alcohol is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG5-alcohol is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG5-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133392</p>	<p><b>Azido-PEG5-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140795</p>
<p>Azido-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b>  <b>Size:</b> 50 mg, 100 mg</p>	<p>Azido-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG5-CH2CO2-NHS</b></p> <p style="text-align: right;">Cat. No.: HY-140766</p>	<p><b>Azido-PEG5-CH2CO2-PFP</b></p> <p style="text-align: right;">Cat. No.: HY-130693</p>
<p>Azido-PEG5-CH2CO2-NHS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG5-CH2CO2-PFP is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG5-CH2CO2H</b></p> <p style="text-align: right;">Cat. No.: HY-130194</p>	<p><b>Azido-PEG5-maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-133398</p>
<p>Azido-PEG5-CH2CO2H is a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG5-CH2CO2H is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 99.60%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Azido-PEG5-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG5-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140757</p>	<p><b>Azido-PEG5-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-130420</p>
<p>Azido-PEG5-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG5-PFP ester is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Azido-PEG5-S-methyl ethanethioate</b></p> <p style="text-align: right;">Cat. No.: HY-132093</p>	<p><b>Azido-PEG5-succinimidyl carbonate</b></p> <p style="text-align: right;">Cat. No.: HY-140769</p>
<p>Azido-PEG5-S-methyl ethanethioate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG5-succinimidyl carbonate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG5-triethoxysilane</b></p> <p style="text-align: right;">Cat. No.: HY-140856</p>	<p><b>Azido-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140453</p>
<p>Azido-PEG5-triethoxysilane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg</p>	<p>Azido-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG6-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-130537</p>	<p><b>Azido-PEG6-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140215</p>
<p>Azido-PEG6-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS. Azido-PEG6-alcohol is also a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG6-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS. Azido-PEG6-amine is also a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 200 mg, 500 mg</p>
<p><b>Azido-PEG6-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133393</p>	<p><b>Azido-PEG6-C1-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140796</p>
<p>Azido-PEG6-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG6-C1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG6-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140779</p>	<p><b>Azido-PEG6-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-134696</p>
<p>Azido-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG6-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Azido-PEG6-MS</b></p> <p style="text-align: right;">Cat. No.: HY-138363</p>	<p><b>Azido-PEG6-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130474</p>
<p>Azido-PEG6-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG6-NHS ester is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG6-NHS ester is also a PEG- and Alkyl/ether based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 98.85%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Azido-PEG6-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140772</p>	<p><b>Azido-PEG6-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138325</p>
<p>Azido-PEG6-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG6-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG7-acid</b></p> <p style="text-align: right;">Cat. No.: HY-132078</p>	<p><b>Azido-PEG7-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140798</p>
<p>Azido-PEG7-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG7-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG7-amine</b></p> <p style="text-align: right;">Cat. No.: HY-130324</p>	<p><b>Azido-PEG7-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133394</p>
<p>Azido-PEG7-amine is a non-cleavable 7 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG7-amine is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p>Azido-PEG7-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG7-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138418</p>	<p><b>Azido-PEG7-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-138707</p>
<p>Azido-PEG7-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Azido-PEG7-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>



<p><b>Azido-PEG7-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-138333</p> <p>Azido-PEG7-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG7-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132060</p> <p>Azido-PEG7-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140454</p> <p>Azido-PEG8-acid is a non-cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG8-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p><b>Azido-PEG8-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140216</p> <p>Azido-PEG8-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Azido-PEG8-azide</b></p> <p style="text-align: right;">Cat. No.: HY-138706</p> <p>Azido-PEG8-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG8-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130742</p> <p>Azido-PEG8-Boc is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 500 mg</p>
<p><b>Azido-PEG8-C-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138377</p> <p>Azido-PEG8-C-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Azido-PEG8-C1-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140767</p> <p>Azido-PEG8-C1-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG8-CH2COO-PFP</b></p> <p style="text-align: right;">Cat. No.: HY-140774</p> <p>Azido-PEG8-CH2COO-PFP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>Azido-PEG8-hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-140815</p> <p>Azido-PEG8-hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Azido-PEG8-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-138499</p>	<p><b>Azido-PEG8-NHBoc</b></p> <p style="text-align: right;">Cat. No.: HY-138708</p>
<p>Azido-PEG8-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG8-NHBoc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG8-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130184</p>	<p><b>Azido-PEG8-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140773</p>
<p>Azido-PEG8-NHS ester is a cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG8-NHS ester is also a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Azido-PEG8-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 97.12%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Azido-PEG8-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-138391</p>	<p><b>Azido-PEG8-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-138437</p>
<p>Azido-PEG8-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG8-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG8-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132055</p>	<p><b>Azido-PEG9-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130475</p>
<p>Azido-PEG8-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG9-acid is a non-cleavable 9 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG9-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG9-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140800</p>	<p><b>Azido-PEG9-amine</b></p> <p style="text-align: right;">Cat. No.: HY-130169</p>
<p>Azido-PEG9-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG9-amine is a non-cleavable 9 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Azido-PEG9-amine is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Azido-PEG9-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133395</p>	<p><b>Azido-PEG9-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140780</p>
<p>Azido-PEG9-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG9-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG9-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138417</p>	<p><b>Azido-PEG9-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140758</p>
<p>Azido-PEG9-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azido-PEG9-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Azido-PEG9-S-methyl ethanethioate</b></p> <p style="text-align: right;">Cat. No.: HY-132094</p>	<p><b>Azidoethyl-SS-PEG2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140107</p>
<p>Azido-PEG9-S-methyl ethanethioate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Azidoethyl-SS-PEG2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>BCN-PEG3-Val-Cit</b></p> <p style="text-align: right;">Cat. No.: HY-140151</p>	<p><b>BCN-PEG4-alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-133437</p>
<p>BCN-PEG3-Val-Cit is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. BCN-PEG3-Val-Cit is also a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>BCN-PEG4-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>BCN-PEG4-hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133438</p>	<p><b>BCN-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-133439</p>
<p>BCN-PEG4-hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>BCN-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>BCN-exo-PEG3-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-133402</p>	<p><b>BDP FL-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-135083</p>
<p>BCN-exo-PEG3-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>BDP FL-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>BDP FL-PEG4-TCO</b></p> <p style="text-align: right;">Cat. No.: HY-141088</p>	<p><b>BDP FL-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-141086</p>
<p>BDP FL-PEG4-TCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>BDP FL-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>BDP FL-PEG5-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-141087</p>	<p><b>Benzaldehyde-PEG4-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133455</p>
<p>BDP FL-PEG5-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzaldehyde-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzenedimethanamine-diethylamine</b></p> <p style="text-align: right;">Cat. No.: HY-140338</p>	<p><b>Benzyl-N-bis(PEG3-Boc)</b> (N-Benzyl-N-bis(PEG3-t-butyl ester))</p> <p style="text-align: right;">Cat. No.: HY-140588</p>
<p>Benzenedimethanamine-diethylamine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-N-bis(PEG3-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG1-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-138342</p>	<p><b>Benzyl-PEG1-propanol</b></p> <p style="text-align: right;">Cat. No.: HY-138433</p>
<p>Benzyl-PEG1-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Benzyl-PEG1-propanol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>

<p><b>Benzyl-PEG1-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-114843</p> <p>Benzyl-PEG1-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG10-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W096081</p> <p>Benzyl-PEG10-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG10-Ots</b></p> <p style="text-align: right;">Cat. No.: HY-138713</p> <p>Benzyl-PEG10-Ots is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG10-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132081</p> <p>Benzyl-PEG10-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG10-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138341</p> <p>Benzyl-PEG10-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG11-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-132074</p> <p>Benzyl-PEG11-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG11-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138345</p> <p>Benzyl-PEG11-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG11-MS</b></p> <p style="text-align: right;">Cat. No.: HY-132023</p> <p>Benzyl-PEG11-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG12-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-132052</p> <p>Benzyl-PEG12-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG12-MS</b></p> <p style="text-align: right;">Cat. No.: HY-138376</p> <p>Benzyl-PEG12-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Benzyl-PEG12-Ots</b></p> <p style="text-align: right;">Cat. No.: HY-138714</p> <p>Benzyl-PEG12-Ots is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG13-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-132035</p> <p>Benzyl-PEG13-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG13-azide</b></p> <p style="text-align: right;">Cat. No.: HY-132082</p> <p>Benzyl-PEG13-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG13-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138480</p> <p>Benzyl-PEG13-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG13-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132011</p> <p>Benzyl-PEG13-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG14-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-132051</p> <p>Benzyl-PEG14-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG14-t-butyl-ester</b></p> <p style="text-align: right;">Cat. No.: HY-132073</p> <p>Benzyl-PEG14-t-butyl-ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG15-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-134697</p> <p>Benzyl-PEG15-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG16-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-132053</p> <p>Benzyl-PEG16-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG16-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138442</p> <p>Benzyl-PEG16-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Benzyl-PEG17-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132054</p>	<p><b>Benzyl-PEG18-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-138444</p>
<p>Benzyl-PEG17-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG18-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG18-MS</b></p> <p style="text-align: right;">Cat. No.: HY-138328</p>	<p><b>Benzyl-PEG18-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138360</p>
<p>Benzyl-PEG18-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG18-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W096142</p>	<p><b>Benzyl-PEG2-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140742</p>
<p>Benzyl-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG2-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG2-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130188</p>	<p><b>Benzyl-PEG2-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130172</p>
<p>Benzyl-PEG2-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG2-CH2-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-135339</p>	<p><b>Benzyl-PEG2-ethanol</b></p> <p style="text-align: right;">Cat. No.: HY-138479</p>
<p>Benzyl-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG2-ethanol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Benzyl-PEG2-ethoxyethane-PEG2</b></p> <p style="text-align: right;">Cat. No.: HY-138476</p>	<p><b>Benzyl-PEG2-MS</b></p> <p style="text-align: right;">Cat. No.: HY-W096072</p>
<p>Benzyl-PEG2-ethoxyethane-PEG2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG2-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG2-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-W043841</p>	<p><b>Benzyl-PEG20-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-138456</p>
<p>Benzyl-PEG2-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG20-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG24-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-134710</p>	<p><b>Benzyl-PEG24-azide</b></p> <p style="text-align: right;">Cat. No.: HY-134717</p>
<p>Benzyl-PEG24-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG24-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG24-MS</b></p> <p style="text-align: right;">Cat. No.: HY-138332</p>	<p><b>Benzyl-PEG24-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138477</p>
<p>Benzyl-PEG24-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG24-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG25-amine</b></p> <p style="text-align: right;">Cat. No.: HY-138319</p>	<p><b>Benzyl-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W096089</p>
<p>Benzyl-PEG25-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>







<p><b>Benzyl-PEG3-amine</b></p> <p style="text-align: right;">Cat. No.: HY-W190795</p>	<p><b>Benzyl-PEG3-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130738</p>
<p>Benzyl-PEG3-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG3-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG3-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-138367</p>	<p><b>Benzyl-PEG3-MS</b></p> <p style="text-align: right;">Cat. No.: HY-132108</p>
<p>Benzyl-PEG3-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG3-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG36-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-138323</p>	<p><b>Benzyl-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130595</p>
<p>Benzyl-PEG36-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG4-acyl chloride</b></p> <p style="text-align: right;">Cat. No.: HY-138513</p>	<p><b>Benzyl-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-W096073</p>
<p>Benzyl-PEG4-acyl chloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG4-Azido</b></p> <p style="text-align: right;">Cat. No.: HY-138715</p>	<p><b>Benzyl-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138346</p>
<p>Benzyl-PEG4-Azido is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Benzyl-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Benzyl-PEG4-MS</b></p> <p style="text-align: right;">Cat. No.: HY-W096090</p> <p>Benzyl-PEG4-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG4-Ots</b></p> <p style="text-align: right;">Cat. No.: HY-W108219</p> <p>Benzyl-PEG4-Ots is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG4-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132008</p> <p>Benzyl-PEG4-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG5-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138487</p> <p>Benzyl-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG5-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140743</p> <p>Benzyl-PEG5-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140744</p> <p>Benzyl-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG5-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-140751</p> <p>Benzyl-PEG5-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG5-NHBoc</b></p> <p style="text-align: right;">Cat. No.: HY-138378</p> <p>Benzyl-PEG5-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG5-Ots</b></p> <p style="text-align: right;">Cat. No.: HY-W108220</p> <p>Benzyl-PEG5-Ots is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG5-THP</b></p> <p style="text-align: right;">Cat. No.: HY-W096075</p> <p>Benzyl-PEG5-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>





<p><b>Benzyl-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138393</p> <p>Benzyl-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG6-amine</b></p> <p style="text-align: right;">Cat. No.: HY-132029</p> <p>Benzyl-PEG6-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG6-azide</b></p> <p style="text-align: right;">Cat. No.: HY-W096070</p> <p>Benzyl-PEG6-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG6-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-132028</p> <p>Benzyl-PEG6-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG6-MS</b></p> <p style="text-align: right;">Cat. No.: HY-132026</p> <p>Benzyl-PEG6-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG6-NHBoc</b></p> <p style="text-align: right;">Cat. No.: HY-132030</p> <p>Benzyl-PEG6-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG6-Ots</b></p> <p style="text-align: right;">Cat. No.: HY-138709</p> <p>Benzyl-PEG6-Ots is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG6-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132034</p> <p>Benzyl-PEG6-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG6-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132105</p> <p>Benzyl-PEG6-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG7-acid</b></p> <p style="text-align: right;">Cat. No.: HY-132027</p> <p>Benzyl-PEG7-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Benzyl-PEG7-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W096067</p> <p>Benzyl-PEG7-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>Benzyl-PEG7-amine</b></p> <p style="text-align: right;">Cat. No.: HY-W096077</p> <p>Benzyl-PEG7-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG7-azide</b></p> <p style="text-align: right;">Cat. No.: HY-132049</p> <p>Benzyl-PEG7-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG7-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-132009</p> <p>Benzyl-PEG7-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG7-MS</b></p> <p style="text-align: right;">Cat. No.: HY-132048</p> <p>Benzyl-PEG7-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG7-NHBoc</b></p> <p style="text-align: right;">Cat. No.: HY-132021</p> <p>Benzyl-PEG7-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG7-Ots</b></p> <p style="text-align: right;">Cat. No.: HY-138710</p> <p>Benzyl-PEG7-Ots is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG7-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132025</p> <p>Benzyl-PEG7-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG7-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132106</p> <p>Benzyl-PEG7-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-132015</p> <p>Benzyl-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Benzyl-PEG8-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W096107</p> <p>Benzyl-PEG8-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG8-amine</b></p> <p style="text-align: right;">Cat. No.: HY-132013</p> <p>Benzyl-PEG8-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG8-azide</b></p> <p style="text-align: right;">Cat. No.: HY-132010</p> <p>Benzyl-PEG8-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG8-Br</b></p> <p style="text-align: right;">Cat. No.: HY-138339</p> <p>Benzyl-PEG8-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG8-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-134741</p> <p>Benzyl-PEG8-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG8-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-132016</p> <p>Benzyl-PEG8-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG8-Ots</b></p> <p style="text-align: right;">Cat. No.: HY-138711</p> <p>Benzyl-PEG8-Ots is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG8-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132044</p> <p>Benzyl-PEG8-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG8-THP</b></p> <p style="text-align: right;">Cat. No.: HY-W096066</p> <p>Benzyl-PEG8-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG9-acid</b></p> <p style="text-align: right;">Cat. No.: HY-132012</p> <p>Benzyl-PEG9-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Benzyl-PEG9-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W093833</p> <p>Benzyl-PEG9-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG9-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-134752</p> <p>Benzyl-PEG9-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzyl-PEG9-Ots</b></p> <p style="text-align: right;">Cat. No.: HY-138712</p> <p>Benzyl-PEG9-Ots is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzyl-PEG9-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138338</p> <p>Benzyl-PEG9-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzoyloxy carbonyl-PEG3-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140745</p> <p>Benzoyloxy carbonyl-PEG3-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzoyloxy carbonyl-PEG3-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140750</p> <p>Benzoyloxy carbonyl-PEG3-C2-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Benzoyloxy carbonyl-PEG3-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140746</p> <p>Benzoyloxy carbonyl-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Benzoyloxy-C5-PEG1</b></p> <p style="text-align: right;">Cat. No.: HY-W096134</p> <p>Benzoyloxy-C5-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-138749</p> <p>Biotin alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.03%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mg, 25 mg, 50 mg</p>	<p><b>Biotin-amido-PEG4-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140937</p> <p>Biotin-amido-PEG4-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Biotin-bis-amido-SS-NHS</b></p> <p style="text-align: right;">Cat. No.: HY-130490</p>	<p><b>Biotin-C1-PEG3-C3-amine TFA</b></p> <p style="text-align: right;">Cat. No.: HY-133403A</p>
<p>Biotin-bis-amido-SS-NHS is an Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-C1-PEG3-C3-amine (TFA) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-C10-NHS Ester</b></p> <p style="text-align: right;">Cat. No.: HY-135920</p>	<p><b>Biotin-C5-amino-C5-amino</b></p> <p style="text-align: right;">Cat. No.: HY-135916</p>
<p>Biotin-C10-NHS Ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-C5-amino-C5-amino is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-C5-NHS Ester</b></p> <p style="text-align: right;">Cat. No.: HY-135915</p>	<p><b>Biotin-EDA</b></p> <p style="text-align: right;">Cat. No.: HY-130893</p>
<p>Biotin-C5-NHS Ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-EDA is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG-amine (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140654</p>	<p><b>Biotin-PEG-amine (MW 3400)</b></p> <p style="text-align: right;">Cat. No.: HY-140655</p>
<p>Biotin-PEG-amine (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG-amine (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG-Biotin (MW 1000)</b></p> <p style="text-align: right;">Cat. No.: HY-140656</p>	<p><b>Biotin-PEG-triethoxysilane (MW 1000)</b></p> <p style="text-align: right;">Cat. No.: HY-140657</p>
<p>Biotin-PEG-Biotin (MW 1000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG-triethoxysilane (MW 1000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Biotin-PEG-triethoxysilane (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140658</p>	<p><b>Biotin-PEG-triethoxysilane (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140659</p>
<p>Biotin-PEG-triethoxysilane (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG-triethoxysilane (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG10-amine</b></p> <p style="text-align: right;">Cat. No.: HY-143827</p>	<p><b>Biotin-PEG10-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-143855</p>
<p>Biotin-PEG10-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG10-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG11-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140900</p>	<p><b>Biotin-PEG11-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140912</p>
<p>Biotin-PEG11-amine is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG11-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG11-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-140909</p>	<p><b>Biotin-PEG11-oxyamine</b></p> <p style="text-align: right;">Cat. No.: HY-140939</p>
<p>Biotin-PEG11-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG11-oxyamine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140493</p>	<p><b>Biotin-PEG12-hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-140933</p>
<p>Biotin-PEG12-acid is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG12-hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>























<p><b>Biotin-PEG12-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-133379</p>	<p><b>Biotin-PEG12-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140892</p>
<p>Biotin-PEG12-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg</p>
<p><b>Biotin-PEG12-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140904</p>	<p><b>Biotin-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-126958</p>
<p>Biotin-PEG12-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG2-acid is a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Biotin-PEG2-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 96.14% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg, 250 mg</p>
<p><b>Biotin-PEG2-aldehyde</b></p> <p style="text-align: right;">Cat. No.: HY-133456</p>	<p><b>Biotin-PEG2-azide</b></p> <p style="text-align: right;">Cat. No.: HY-126957</p>
<p>Biotin-PEG2-aldehyde is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Biotin-PEG2-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG2-C1-aldehyde</b></p> <p style="text-align: right;">Cat. No.: HY-133457</p>	<p><b>Biotin-PEG2-C2-iodoacetamide</b></p> <p style="text-align: right;">Cat. No.: HY-140941</p>
<p>Biotin-PEG2-C1-aldehyde is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG2-C2-iodoacetamide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Biotin-PEG2-C4-alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-113828</p>	<p><b>Biotin-PEG2-C6-azide</b></p> <p style="text-align: right;">Cat. No.: HY-117042</p>
<p>Biotin-PEG2-C4-alkyne is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG2-C6-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Biotin-PEG2-CH2CH2N3</b></p> <p style="text-align: right;">Cat. No.: HY-135912</p>	<p><b>Biotin-PEG2-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-W010764</p>
<p>Biotin-PEG2-CH2CH2N3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG2-Mal is a PEG-based PROTAC linker can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Biotin-PEG2-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140934</p>	<p><b>Biotin-PEG2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140888</p>
<p>Biotin-PEG2-NH-Boc is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG2-NHS ester is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG2-OH</b></p> <p style="text-align: right;">Cat. No.: HY-135911</p>	<p><b>Biotin-PEG2-SH</b></p> <p style="text-align: right;">Cat. No.: HY-132112</p>
<p>Biotin-PEG2-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG2-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Biotin-PEG23-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140901</p>	<p><b>Biotin-PEG23-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140913</p>
<p>Biotin-PEG23-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG23-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG24-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133377</p>	<p><b>Biotin-PEG24-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-133378</p>
<p>Biotin-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG24-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Biotin-PEG24-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140905</p>	<p><b>Biotin-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-116027</p>
<p>Biotin-PEG24-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG3-acid is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg</p>
<p><b>Biotin-PEG3-alcohol</b>            ((+)-Biotin-PEG3-OH)</p> <p style="text-align: right;">Cat. No.: HY-135179</p>	<p><b>Biotin-PEG3-amido-SS-amido-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133458</p>
<p>Biotin-PEG3-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG3-amido-SS-amido-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130143</p>	<p><b>Biotin-PEG3-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W096098</p>
<p>Biotin-PEG3-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 250 mg</p>	<p>Biotin-PEG3-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG3-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-140907</p>	<p><b>Biotin-PEG3-methyl ethanethioate</b></p> <p style="text-align: right;">Cat. No.: HY-132116</p>
<p>Biotin-PEG3-Mal is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Biotin-PEG3-methyl ethanethioate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG3-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130291</p>	<p><b>Biotin-PEG3-OH</b></p> <p style="text-align: right;">Cat. No.: HY-135924</p>
<p>Biotin-PEG3-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG3-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Biotin-PEG3-oxyamine</b></p> <p style="text-align: right;">Cat. No.: HY-140938</p> <p>Biotin-PEG3-oxyamine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Biotin-PEG3-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-138503</p> <p>Biotin-PEG3-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG3-propionic hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-130560</p> <p>Biotin-PEG3-propionic hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Biotin-PEG3-pyridinrthiol</b></p> <p style="text-align: right;">Cat. No.: HY-132114</p> <p>Biotin-PEG3-pyridinrthiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG3-SH</b></p> <p style="text-align: right;">Cat. No.: HY-132115</p> <p>Biotin-PEG3-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Biotin-PEG36-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140494</p> <p>Biotin-PEG36-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG36-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140906</p> <p>Biotin-PEG36-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Biotin-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-126959</p> <p>Biotin-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG4-alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-140922</p> <p>Biotin-PEG4-alkyne is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.88% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mg, 50 mg, 100 mg</p>	<p><b>Biotin-PEG4-allyl</b></p> <p style="text-align: right;">Cat. No.: HY-138426</p> <p>Biotin-PEG4-allyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Biotin-PEG4-amide-Alkyne</b></p> <p>Cat. No.: HY-140923</p>	<p><b>Biotin-PEG4-Amide-C6-Azide</b></p> <p>Cat. No.: HY-140914</p>
<p>Biotin-PEG4-amide-Alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG4-Amide-C6-Azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.37%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 5 mg, 10 mg</p>
<p><b>Biotin-PEG4-amine</b></p> <p>Cat. No.: HY-140895</p>	<p><b>Biotin-PEG4-amino-t-Bu-DADPS-C3-alkyne</b></p> <p>Cat. No.: HY-140928</p>
<p>Biotin-PEG4-amine is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 10 mM × 1 mL, 5 mg, 10 mg, 50 mg</p>	<p>Biotin-PEG4-amino-t-Bu-DADPS-C3-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG4-amino-t-Bu-DADPS-C6-azide</b></p> <p>Cat. No.: HY-140921</p>	<p><b>Biotin-PEG4-azide</b></p> <p>Cat. No.: HY-140910</p>
<p>Biotin-PEG4-amino-t-Bu-DADPS-C6-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 5 mg, 10 mg</p>	<p>Biotin-PEG4-azide is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Biotin-PEG4-hydrazide</b></p> <p>Cat. No.: HY-140932</p>	<p><b>Biotin-PEG4-hydrazide TFA</b></p> <p>Cat. No.: HY-140932A</p>
<p>Biotin-PEG4-hydrazide is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG4-hydrazide (TFA) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG4-methyltetrazine</b></p> <p>Cat. No.: HY-140940</p>	<p><b>Biotin-PEG4-NHS ester</b></p> <p>Cat. No.: HY-140889</p>
<p>Biotin-PEG4-methyltetrazine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.04%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 5 mg, 10 mg</p>	<p>Biotin-PEG4-NHS ester is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.08%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 50 mg, 100 mg, 250 mg</p>

<p><b>Biotin-PEG4-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130290</p>	<p><b>Biotin-PEG4-PC-PEG4-alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-140131</p>
<p>Biotin-PEG4-OH is a PEG-based PROTAC linker can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG4-PC-PEG4-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG4-Picolyl azide</b></p> <p style="text-align: right;">Cat. No.: HY-140915</p>	<p><b>Biotin-PEG4-SH</b></p> <p style="text-align: right;">Cat. No.: HY-134704</p>
<p>Biotin-PEG4-Picolyl azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 5 mg, 10 mg</p>	<p>Biotin-PEG4-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG4-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140903</p>	<p><b>Biotin-PEG5-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140896</p>
<p>Biotin-PEG4-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG5-amine is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133174</p>	<p><b>Biotin-PEG5-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140935</p>
<p>Biotin-PEG5-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>Biotin-PEG5-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133176</p>	<p><b>Biotin-PEG6-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140893</p>
<p>Biotin-PEG6-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG6-alcohol is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Biotin-PEG6-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140897</p>	<p><b>Biotin-PEG6-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140911</p>
<p>Biotin-PEG6-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG6-azide is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG6-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140936</p>	<p><b>Biotin-PEG6-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-140908</p>
<p>Biotin-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG6-Mal is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Biotin-PEG6-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-W190945</p>	<p><b>Biotin-PEG6-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140890</p>
<p>Biotin-PEG6-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG6-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG6-Silane</b></p> <p style="text-align: right;">Cat. No.: HY-138419</p>	<p><b>Biotin-PEG6-Thalidomide</b></p> <p style="text-align: right;">Cat. No.: HY-140942</p>
<p>Biotin-PEG6-Silane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG6-Thalidomide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b>  <b>Size:</b> 5 mg, 10 mg, 25 mg</p>
<p><b>Biotin-PEG7-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140898</p>	<p><b>Biotin-PEG7-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133175</p>
<p>Biotin-PEG7-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG7-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>





<p><b>Biotin-PEG7-C2-NH-Vidarabine-S-CH3</b></p> <p style="text-align: right;">Cat. No.: HY-145248</p>	<p><b>Biotin-PEG7-C2-S-Vidarabine</b></p> <p style="text-align: right;">Cat. No.: HY-145247</p>
<p>Biotin-PEG7-C2-NH-Vidarabine-S-CH3 is a PEG-based linker that incorporates adenosine analog Vidarabine. Vidarabine is an antiviral agent which is active against herpes simplex and varicella zoster viruses.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG7-C2-S-Vidarabine is a PEG-based linker that incorporates adenosine analog Vidarabine. Vidarabine is an antiviral agent which is active against herpes simplex and varicella zoster viruses.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG7-thiourea</b></p> <p style="text-align: right;">Cat. No.: HY-140943</p>	<p><b>Biotin-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140492</p>
<p>Biotin-PEG7-thiourea is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG8-acid is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG8-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140894</p>	<p><b>Biotin-PEG8-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140899</p>
<p>Biotin-PEG8-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG8-amine is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG8-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140891</p>	<p><b>Biotin-PEG8-Vidarabine</b></p> <p style="text-align: right;">Cat. No.: HY-145246</p>
<p>Biotin-PEG8-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG8-Vidarabine is a PEG-based linker that incorporates adenosine analog Vidarabine. Vidarabine is an antiviral agent which is active against herpes simplex and varicella zoster viruses.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-PEG9-amine</b></p> <p style="text-align: right;">Cat. No.: HY-W190970</p>	<p><b>Biotin-PEG9-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-135937</p>
<p>Biotin-PEG9-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Biotin-PEG9-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>























<p><b>Biotin-PEG9-NHS Ester</b></p> <p style="text-align: right;">Cat. No.: HY-135941</p> <p>Biotin-PEG9-NHS Ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Biotin-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-119343</p> <p>Biotin-PFP ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Biotin-Thalidomide</b></p> <p style="text-align: right;">Cat. No.: HY-145152</p> <p>Biotin-Thalidomide is a cereblon affinity probe for PROTAC and targeted protein degradation research.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Biotinyl-NH-PEG3-C3-amido-C3-COOH (DIPEA)</b></p> <p style="text-align: right;">Cat. No.: HY-130142</p> <p>Biotinyl-NH-PEG3-C3-amido-C3-COOH (DIPEA) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis(2-bromoethyl) ether</b></p> <p style="text-align: right;">Cat. No.: HY-W013458</p> <p>Bis(2-bromoethyl) ether is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.35%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>	<p><b>Bis(m-PEG4)-N-OH</b></p> <p style="text-align: right;">Cat. No.: HY-140580</p> <p>Bis(m-PEG4)-N-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-(m-PEG4)-amido-hexanoic acid</b></p> <p style="text-align: right;">Cat. No.: HY-140016</p> <p>Bis-(m-PEG4)-amido-hexanoic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bis-(m-PEG8-amido)-hexanoic acid</b></p> <p style="text-align: right;">Cat. No.: HY-141290</p> <p>Bis-(m-PEG8-amido)-hexanoic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-(Mal-PEG3)-PH-N-succinimidyl acetate</b></p> <p style="text-align: right;">Cat. No.: HY-138519</p> <p>Bis-(Mal-PEG3)-PH-N-succinimidyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bis-(N,N'-carboxyl-PEG4)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141042</p> <p>Bis-(N,N'-carboxyl-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Bis-(N,N'-PEG4-NHS ester)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141047</p>	<p><b>Bis-(N,N'-amine-PEG3)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141057</p>
<p>Bis-(N,N'-PEG4-NHS ester)-Cy5 is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-(N,N'-amine-PEG3)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-acrylate-PEG5</b></p> <p style="text-align: right;">Cat. No.: HY-W096160</p>	<p><b>Bis-acrylate-PEG6</b></p> <p style="text-align: right;">Cat. No.: HY-W096166</p>
<p>Bis-acrylate-PEG5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-acrylate-PEG6 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-aminoxy-PEG1</b></p> <p style="text-align: right;">Cat. No.: HY-140407</p>	<p><b>Bis-aminoxy-PEG2</b></p> <p style="text-align: right;">Cat. No.: HY-140408</p>
<p>Bis-aminoxy-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg, 500 mg</p>	<p>Bis-aminoxy-PEG2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-aminoxy-PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-140409</p>	<p><b>Bis-aminoxy-PEG4</b></p> <p style="text-align: right;">Cat. No.: HY-140410</p>
<p>Bis-aminoxy-PEG3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-aminoxy-PEG4 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-aminoxy-PEG7</b></p> <p style="text-align: right;">Cat. No.: HY-140411</p>	<p><b>Bis-BCN-PEG1-diamide</b></p> <p style="text-align: right;">Cat. No.: HY-133510</p>
<p>Bis-aminoxy-PEG7 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-BCN-PEG1-diamide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Bis-Biotin-PEG23</b></p> <p style="text-align: right;">Cat. No.: HY-140929</p>	<p><b>Bis-Bromoacetamido-PEG11</b></p> <p style="text-align: right;">Cat. No.: HY-141393</p>
<p>Bis-Biotin-PEG23 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-Bromoacetamido-PEG11 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-CH2-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-132102</p>	<p><b>Bis-isopropyl-PEG1</b></p> <p style="text-align: right;">Cat. No.: HY-138455</p>
<p>Bis-CH2-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-isopropyl-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-Mal-Lysine-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141001</p>	<p><b>Bis-Mal-Lysine-PEG4-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-141002</p>
<p>Bis-Mal-Lysine-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg, 250 mg</p>	<p>Bis-Mal-Lysine-PEG4-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-Mal-PEG11</b></p> <p style="text-align: right;">Cat. No.: HY-140995</p>	<p><b>Bis-Mal-PEG19</b></p> <p style="text-align: right;">Cat. No.: HY-140996</p>
<p>Bis-Mal-PEG11 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-Mal-PEG19 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-Mal-PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-140993</p>	<p><b>Bis-Mal-PEG5</b></p> <p style="text-align: right;">Cat. No.: HY-130895</p>
<p>Bis-Mal-PEG3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-Mal-PEG5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Bis-Mal-PEG6</b></p> <p style="text-align: right;">Cat. No.: HY-140994</p> <p>Bis-Mal-PEG6 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bis-Mal-PEG7</b> (Mal-NH-PEG7-NH-Mal)</p> <p style="text-align: right;">Cat. No.: HY-130896</p> <p>Bis-Mal-PEG7 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-methacrylate-PEG5</b></p> <p style="text-align: right;">Cat. No.: HY-W096161</p> <p>Bis-methacrylate-PEG5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bis-NH2-C1-PEG3</b> (PROTAC Linker 24)</p> <p style="text-align: right;">Cat. No.: HY-128844</p> <p>Bis-NH2-C1-PEG3 (PROTAC Linker 24) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-NH2-PEG2</b> (PROTAC Linker 19)</p> <p style="text-align: right;">Cat. No.: HY-128833</p> <p>Bis-NH2-PEG2 (PROTAC Linker 19) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p><b>Bis-PEG-TFP ester (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140724</p> <p>Bis-PEG-TFP ester (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG1-acid</b></p> <p style="text-align: right;">Cat. No.: HY-23166</p> <p>Bis-PEG1-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p><b>Bis-PEG1-C-PEG1-CH2COOH</b> (PROTAC Linker 26)</p> <p style="text-align: right;">Cat. No.: HY-128847</p> <p>Bis-PEG1-C-PEG1-CH2COOH (PROTAC Linker 26) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg</p>
<p><b>Bis-PEG10-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133231</p> <p>Bis-PEG10-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bis-PEG10-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130824</p> <p>Bis-PEG10-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG10-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Bis-PEG10-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132057</p>	<p><b>Bis-PEG11-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141019</p>
<p>Bis-PEG10-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG11-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG11-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141100</p>	<p><b>Bis-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141020</p>
<p>Bis-PEG11-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG12-endo-BCN</b></p> <p style="text-align: right;">Cat. No.: HY-139060</p>	<p><b>Bis-PEG13-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141021</p>
<p>Bis-PEG12-endo-BCN is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG13-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG13-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130825</p>	<p><b>Bis-PEG13-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-141256</p>
<p>Bis-PEG13-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG13-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG13-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG14-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141022</p>	<p><b>Bis-PEG15-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141023</p>
<p>Bis-PEG14-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG15-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Bis-PEG17-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141024</p>	<p><b>Bis-PEG17-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130826</p>
<p>Bis-PEG17-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG17-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG17-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG18-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138326</p>	<p><b>Bis-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-112559</p>
<p>Bis-PEG18-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG2-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 500 mg</p>
<p><b>Bis-PEG2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138504</p>	<p><b>Bis-PEG2-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-112560</p>
<p>Bis-PEG2-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG2-PFP ester is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Bis-PEG2-PFP ester is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG21-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141025</p>	<p><b>Bis-PEG21-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130827</p>
<p>Bis-PEG21-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG21-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG21-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG23-endo-BCN</b></p> <p style="text-align: right;">Cat. No.: HY-140079</p>	<p><b>Bis-PEG25-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141026</p>
<p>Bis-PEG23-endo-BCN is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> <b>Size:</b> 50 mg</p>	<p>Bis-PEG25-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Bis-PEG25-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130828</p>	<p><b>Bis-PEG25-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-141258</p>
<p>Bis-PEG25-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG25-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG25-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG29-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141027</p>	<p><b>Bis-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-126891</p>
<p>Bis-PEG29-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG3-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Bis-PEG3-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-126997</p>	<p><b>Bis-PEG3-sulfonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-140165</p>
<p>Bis-PEG3-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG3-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG3-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132066</p>	<p><b>Bis-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-119429</p>
<p>Bis-PEG3-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG4-acid is a PEG PROTAC linker.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>Bis-PEG4-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-126998</p>	<p><b>Bis-PEG4-sulfonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-140166</p>
<p>Bis-PEG4-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG4-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Bis-PEG4-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132067</p>	<p><b>Bis-PEG4-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-W190728</p>
<p>Bis-PEG4-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG4-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG5-acid</b> (PROTAC Linker 36)</p> <p style="text-align: right;">Cat. No.: HY-116006</p>	<p><b>Bis-PEG5-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-126889</p>
<p>Bis-PEG5-acid (PROTAC Linker 36) is a PROTAC linker, which belongs to a polyethylene glycol (PEG) linker. Bis-PEG5-acid (PROTAC Linker 36) can be used in the synthesis of the CP5V. CP5V is a PROTAC, and specifically degrades Cdc20.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>	<p>Bis-PEG5-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG5-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG5-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-126999</p>	<p><b>Bis-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-120587</p>
<p>Bis-PEG5-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG6-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>
<p><b>Bis-PEG6-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130410</p>	<p><b>Bis-PEG6-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-W096076</p>
<p>Bis-PEG6-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG6-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p>Bis-PEG6-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG7-acid</b></p> <p style="text-align: right;">Cat. No.: HY-126892</p>	<p><b>Bis-PEG7-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-126890</p>
<p>Bis-PEG7-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG6-propionic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG7-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG7-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>














<p><b>Bis-PEG7-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-133000</p>	<p><b>Bis-PEG7-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132022</p>
<p>Bis-PEG7-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG7-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-126893</p>	<p><b>Bis-PEG8-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132080</p>
<p>Bis-PEG8-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG8-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG8-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-PEG9-acid</b></p> <p style="text-align: right;">Cat. No.: HY-126894</p>	<p><b>Bis-PEG9-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-117009</p>
<p>Bis-PEG9-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG9-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-PEG9-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Bis-PEG9-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Bis-PEG9-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-141255</p>	<p><b>Bis-propargyl-O-PEG17</b></p> <p style="text-align: right;">Cat. No.: HY-140042</p>
<p>Bis-PEG9-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-propargyl-O-PEG17 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-propargyl-PEG1</b></p> <p style="text-align: right;">Cat. No.: HY-140038</p>	<p><b>Bis-propargyl-PEG10</b></p> <p style="text-align: right;">Cat. No.: HY-132098</p>
<p>Bis-propargyl-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>	<p>Bis-propargyl-PEG10 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Bis-propargyl-PEG11</b></p> <p style="text-align: right;">Cat. No.: HY-140039</p> <p>Bis-propargyl-PEG11 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bis-propargyl-PEG12</b></p> <p style="text-align: right;">Cat. No.: HY-140040</p> <p>Bis-propargyl-PEG12 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-propargyl-PEG13</b></p> <p style="text-align: right;">Cat. No.: HY-140041</p> <p>Bis-propargyl-PEG13 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bis-propargyl-PEG2</b></p> <p style="text-align: right;">Cat. No.: HY-133191</p> <p>Bis-propargyl-PEG2 is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Bis-propargyl-PEG2 is used for the synthesis of demethylvancomycin dimers.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-propargyl-PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-133192</p> <p>Bis-propargyl-PEG3 is a PEG-based PROTAC linker used in the synthesis of PROTACs. Bis-propargyl-PEG3 is used in the synthesis of zinc-dipicolylamine (ZnDPA) complexes with antiparasmodial activity .</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 1 g</p>	<p><b>Bis-propargyl-PEG4</b></p> <p style="text-align: right;">Cat. No.: HY-120397</p> <p>Bis-propargyl-PEG4 is a PEG-based PROTAC linker used in the synthesis of PROTACs. Bis-propargyl-PEG4 is used for the synthesis of demethylvancomycin dimers.</p>  <p><b>Purity:</b> 95.64%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 250 mg</p>
<p><b>Bis-propargyl-PEG5</b></p> <p style="text-align: right;">Cat. No.: HY-133193</p> <p>Bis-propargyl-PEG5 is a PEG-based PROTAC linker used in the synthesis of PROTACs. Bis-propargyl-PEG5 is used for the synthesis of carbohydrate receptors (SCRs) with anti-Zika activity.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p><b>Bis-propargyl-PEG6</b></p> <p style="text-align: right;">Cat. No.: HY-117186</p> <p>Bis-propargyl-PEG6 is a PEG-based PROTAC linker used in the synthesis of PROTACs. Bis-propargyl-PEG6 can be used to synthesize the polymer linked multimers of guanosine-3', 5'-cyclic monophosphates.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bis-propargyl-PEG7</b></p> <p style="text-align: right;">Cat. No.: HY-133190</p> <p>Bis-propargyl-PEG7 is a PEG-based PROTAC linker used in the synthesis of PROTACs. Bis-propargyl-PEG7 can be used to synthesize the polymer linked multimers of guanosine-3', 5'-cyclic monophosphates.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg</p>	<p><b>Bis-propargyl-PEG8</b></p> <p style="text-align: right;">Cat. No.: HY-115414</p> <p>Bis-propargyl-PEG8 (compound 16e) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Bis-propargyl-PEG9</b></p> <p style="text-align: right;">Cat. No.: HY-133189</p>	<p><b>Bis-sulfone-PEG4-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-138753</p>
<p>Bis-propargyl-PEG9 is a PEG-based PROTAC linker used in the synthesis of PROTACs. Bis-propargyl-PEG9 can be used to synthesize the bivalent estrogen receptor ligands.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-sulfone-PEG4-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg, 25 mg, 50 mg</p>
<p><b>Bis-sulfone-PEG4-Tetrazine</b></p> <p style="text-align: right;">Cat. No.: HY-138747</p>	<p><b>Bis-Tos-PEG4 (PROTAC Linker 16)</b></p> <p style="text-align: right;">Cat. No.: HY-W004816</p>
<p>Bis-sulfone-PEG4-Tetrazine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-Tos-PEG4 (PROTAC Linker 16) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.61%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>Bis-Tos-PEG6</b></p> <p style="text-align: right;">Cat. No.: HY-140367</p>	<p><b>Bis-Tos-PEG7</b></p> <p style="text-align: right;">Cat. No.: HY-140368</p>
<p>Bis-Tos-PEG6 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bis-Tos-PEG7 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>BM-PEG3</b> (1,11-Bis-maleimidotetraethyleneglycol)</p> <p style="text-align: right;">Cat. No.: HY-130898</p>	<p><b>BnO-PEG1-CH2CO2tBu</b></p> <p style="text-align: right;">Cat. No.: HY-140747</p>
<p>BM-PEG3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>BnO-PEG1-CH2CO2tBu is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>BnO-PEG1-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-121957</p>	<p><b>BnO-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140748</p>
<p>BnO-PEG1-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>	<p>BnO-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>BnO-PEG4-OH</b></p> <p style="text-align: right;">Cat. No.: HY-W040228</p> <p>BnO-PEG4-OH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>BnO-PEG5-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140749</p> <p>BnO-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>BnO-PEG5-OH</b></p> <p style="text-align: right;">Cat. No.: HY-W042714</p> <p>BnO-PEG5-OH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>BnO-PEG6-OH</b></p> <p style="text-align: right;">Cat. No.: HY-W042654</p> <p>BnO-PEG6-OH is a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). BnO-PEG6-OH is also a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.88%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>BnOH-NH-bis-(C2-S)-propane-O-isoprene ester</b> (PROTAC Linker 29)</p> <p style="text-align: right;">Cat. No.: HY-126313</p> <p>BnOH-NH-bis-(C2-S)-propane-O-isoprene ester (PROTAC Linker 29) is an alkyl ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-11-aminoundecanoic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W013253</p> <p>Boc-11-aminoundecanoic acid is an Alkyl/ether-based PROTAC linker can be used in the synthesis of MS432 (HY-130602).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-6-aminohexanoic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W007529</p> <p>Boc-6-aminohexanoic acid is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 g</p>	<p><b>Boc-amido-PEG9-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140233</p> <p>Boc-amido-PEG9-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-Aminoxy-PEG1-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140430</p> <p>Boc-Aminoxy-PEG1-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-Aminoxy-PEG1-C2-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140425</p> <p>Boc-Aminoxy-PEG1-C2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>









<p><b>Boc-aminoxy-PEG1-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140049</p> <p>Boc-aminoxy-PEG1-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-Aminoxy-PEG2</b></p> <p style="text-align: right;">Cat. No.: HY-140421</p> <p>Boc-Aminoxy-PEG2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-Aminoxy-PEG2-C2-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140426</p> <p>Boc-Aminoxy-PEG2-C2-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-Aminoxy-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140415</p> <p>Boc-Aminoxy-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-aminoxy-PEG2-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140050</p> <p>Boc-aminoxy-PEG2-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-Aminoxy-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140412</p> <p>Boc-Aminoxy-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-Aminoxy-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140432</p> <p>Boc-Aminoxy-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-Aminoxy-PEG3-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-140435</p> <p>Boc-Aminoxy-PEG3-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-Aminoxy-PEG3-C2-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140427</p> <p>Boc-Aminoxy-PEG3-C2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-Aminoxy-PEG3-thiol</b></p> <p style="text-align: right;">Cat. No.: HY-140439</p> <p>Boc-Aminoxy-PEG3-thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Boc-Aminoxy-PEG4-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140433</p>	<p><b>Boc-Aminoxy-PEG4-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140420</p>
<p>Boc-Aminoxy-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-Aminoxy-PEG4-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-Aminoxy-PEG4-CH2CO2H</b></p> <p style="text-align: right;">Cat. No.: HY-140416</p>	<p><b>Boc-Aminoxy-PEG4-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140438</p>
<p>Boc-Aminoxy-PEG4-CH2CO2H is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-Aminoxy-PEG4-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-aminoxy-PEG4-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140052</p>	<p><b>Boc-Aminoxy-PEG4-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140377</p>
<p>Boc-aminoxy-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-Aminoxy-PEG4-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-Aminoxy-PEG5-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140428</p>	<p><b>Boc-aminoxy-PEG5-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140053</p>
<p>Boc-Aminoxy-PEG5-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-aminoxy-PEG5-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-aminoxy-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140413</p>	<p><b>Boc-Aminoxy-PEG4-OH</b></p> <p style="text-align: right;">Cat. No.: HY-140423</p>
<p>Boc-aminoxy-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-Aminoxy-PEG4-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Boc-bipiperidine-ethynylbenzoic acid</b></p> <p>Cat. No.: HY-139661</p>	<p><b>Boc-C1-PEG2-C4-Cl</b> (PROTAC Linker 1)</p> <p>Cat. No.: HY-108371</p>
<p>Boc-bipiperidine-ethynylbenzoic acid is an Alkyl/ether-based <b>PROTAC linker</b> can be used in the synthesis of ARD-61.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-C1-PEG2-C4-Cl (PROTAC Linker 1) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>Boc-C1-PEG3-C4-OBn</b></p> <p>Cat. No.: HY-130619</p>	<p><b>Boc-C1-PEG3-C4-OH</b></p> <p>Cat. No.: HY-130618</p>
<p>Boc-C1-PEG3-C4-OBn (PROTAC Linker 15) is a <b>PROTAC linker</b>, which refers to the PEG composition. Boc-C1-PEG3-C4-OBn can be used in the synthesis of a series of PROTACs, such as PROTAC SGK3 degrader-1 (HY-125878).</p>  <p><b>Purity:</b> 95.77% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 500 mg</p>	<p>Boc-C1-PEG3-C4-OH is a <b>PROTAC linker</b>, which refers to the Alkyl/ether composition. Boc-C1-PEG3-C4-OH can be used in the synthesis of a series of PROTACs.</p>  <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 250 mg</p>
<p><b>Boc-C14-COOH</b></p> <p>Cat. No.: HY-W034599</p>	<p><b>Boc-C16-COOH</b></p> <p>Cat. No.: HY-W045598</p>
<p>Boc-C14-COOH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-C14-COOH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-C16-COOH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-C16-COOH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs &lt;su.</p>  <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg, 250 mg</p>
<p><b>Boc-C16-NHS ester</b></p> <p>Cat. No.: HY-140342</p>	<p><b>Boc-C2-NH2</b></p> <p>Cat. No.: HY-W017970</p>
<p>Boc-C16-NHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-C2-NH2 is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 250 mg</p>
<p><b>Boc-C5-O-C5-O-C6-Cl</b> (PROTAC Linker 2)</p> <p>Cat. No.: HY-108372</p>	<p><b>Boc-GABA-OH</b></p> <p>Cat. No.: HY-W004697</p>
<p>Boc-C5-O-C5-O-C6-Cl (PROTAC Linker 2) is a <b>PROTAC linker</b> utilized to connect the respective tyrosine kinase inhibitor (TKI) to the E3 recruiting ligand.</p>  <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p>Boc-GABA-OH is a <b>PROTAC linker</b> which can be used to synthesis UNC6852, an EED-targeted PROTAC.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 100 mg</p>











<p><b>Boc-Gly-amido-C-PEG3-C3-amine</b></p> <p>Cat. No.: HY-140237</p>	<p><b>Boc-gly-PEG3-endo-BCN</b></p> <p>Cat. No.: HY-140081</p>
<p>Boc-Gly-amido-C-PEG3-C3-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-gly-PEG3-endo-BCN is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Boc-gly-PEG3-endo-BCN is also a cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-HyNic-PEG1-mal</b></p> <p>Cat. No.: HY-133499</p>	<p><b>Boc-HyNic-PEG2-alkyne</b></p> <p>Cat. No.: HY-133500</p>
<p>Boc-HyNic-PEG1-mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-HyNic-PEG2-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-HyNic-PEG2-DBCO</b></p> <p>Cat. No.: HY-133502</p>	<p><b>Boc-HyNic-PEG2-N3</b></p> <p>Cat. No.: HY-133501</p>
<p>Boc-HyNic-PEG2-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Boc-HyNic-PEG2-N3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-Hyp-OH</b></p> <p>Cat. No.: HY-10781</p>	<p><b>Boc-Hyp-OMe</b></p> <p>Cat. No.: HY-65039</p>
<p>Boc-Hyp-OH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-Hyp-OH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>1/5</sup>.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 g, 5 g</p>	<p>Boc-Hyp-OMe is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-Hyp-OMe is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 g, 5 g</p>
<p><b>Boc-N-Amido-PEG2-C2-azide</b></p> <p>Cat. No.: HY-140834</p>	<p><b>Boc-N-amido-PEG3-acid</b></p> <p>Cat. No.: HY-140468</p>
<p>Boc-N-Amido-PEG2-C2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-N-amido-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 500 mg</p>




<p><b>Boc-N-Amido-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140835</p> <p>Boc-N-Amido-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-N-Amido-PEG4-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140879</p> <p>Boc-N-Amido-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-N-Amido-PEG5-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-140392</p> <p>Boc-N-Amido-PEG5-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-N-PEG1-C2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141116</p> <p>Boc-N-PEG1-C2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-N-PEG2-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-140390</p> <p>Boc-N-PEG2-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-N-PEG5-C2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-126381</p> <p>Boc-N-PEG5-C2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-C12-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-133388</p> <p>Boc-NH-C12-NH2 is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-C4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W014099</p> <p>Boc-NH-C4-acid is a PROTAC linker, which belongs to a Alkyl/ether linker. Boc-NH-C4-acid can be used in the synthesis of the compound PROTAC1, and specifically degrades EED, EZH2, and SUZ12 in the PRC2 Complex.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>Boc-NH-C4-Br</b></p> <p style="text-align: right;">Cat. No.: HY-W007803</p> <p>Boc-NH-C4-Br is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p><b>Boc-NH-C6-amido-C4-acid (PROTAC Linker 32)</b></p> <p style="text-align: right;">Cat. No.: HY-125888</p> <p>Boc-NH-C6-amido-C4-acid (PROTAC Linker 32) is an alkyl ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Boc-NH-O-C1-NHS ester</b></p> <p>Cat. No.: HY-133405</p>	<p><b>Boc-NH-PEG-amine (MW 2000)</b></p> <p>Cat. No.: HY-140725</p>
<p>Boc-NH-O-C1-NHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>	<p>Boc-NH-PEG-amine (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG-amine (MW 3400)</b></p> <p>Cat. No.: HY-140726</p>	<p><b>Boc-NH-PEG-amine (MW 5000)</b></p> <p>Cat. No.: HY-140727</p>
<p>Boc-NH-PEG-amine (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG-amine (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG-Thiol (MW 10000)</b></p> <p>Cat. No.: HY-138311</p>	<p><b>Boc-NH-PEG1-C5-OH</b></p> <p>Cat. No.: HY-134737</p>
<p>Boc-NH-PEG-Thiol (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG1-C5-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG1-CH2CH2COOH</b></p> <p>Cat. No.: HY-120775</p>	<p><b>Boc-NH-PEG1-CH2COOH</b></p> <p>Cat. No.: HY-140476</p>
<p>Boc-NH-PEG1-CH2CH2COOH is a cleavable (1 unit PEG) ADC linker and also a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of antibody-drug conjugates (ADCs) or PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG1-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>
<p><b>Boc-NH-PEG1-OH</b></p> <p>Cat. No.: HY-W007322</p>	<p><b>Boc-NH-PEG1-Ph-O-CH2COOH</b></p> <p>Cat. No.: HY-130637</p>
<p>Boc-NH-PEG1-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>	<p>Boc-NH-PEG1-Ph-O-CH2COOH is a PROTAC Linker which is used for the EED-targeted PROTAC.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>







<p><b>Boc-NH-PEG10-CH2CH2COOH</b></p> <p>Cat. No.: HY-140471</p>	<p><b>Boc-NH-PEG10-CH2CH2NH2</b></p> <p>Cat. No.: HY-135930</p>
<p>Boc-NH-PEG10-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG10-CH2CH2NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG10-NHS ester</b></p> <p>Cat. No.: HY-141120</p>	<p><b>Boc-NH-PEG11-C2-acid</b></p> <p>Cat. No.: HY-138410</p>
<p>Boc-NH-PEG10-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG11-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG11-CH2CH2N3</b></p> <p>Cat. No.: HY-130897</p>	<p><b>Boc-NH-PEG11-NH2</b></p> <p>Cat. No.: HY-140234</p>
<p>Boc-NH-PEG11-CH2CH2N3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG11-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG11-NHS ester</b></p> <p>Cat. No.: HY-138441</p>	<p><b>Boc-NH-PEG11-OH</b></p> <p>Cat. No.: HY-135947</p>
<p>Boc-NH-PEG11-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG11-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG12-CH2CH2COOH</b></p> <p>Cat. No.: HY-140472</p>	<p><b>Boc-NH-PEG12-NH-Boc</b></p> <p>Cat. No.: HY-138494</p>
<p>Boc-NH-PEG12-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 100 mg, 250 mg</p>	<p>Boc-NH-PEG12-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>

<p><b>Boc-NH-PEG12-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-133355</p> <p>Boc-NH-PEG12-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG12-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140885</p> <p>Boc-NH-PEG12-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG15-azide</b></p> <p style="text-align: right;">Cat. No.: HY-133356</p> <p>Boc-NH-PEG15-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG15-C1-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138402</p> <p>Boc-NH-PEG15-C1-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG15-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138397</p> <p>Boc-NH-PEG15-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG15-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140235</p> <p>Boc-NH-PEG15-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG2</b> (PROTAC Linker 11)</p> <p style="text-align: right;">Cat. No.: HY-W004896</p> <p>Boc-NH-PEG2 (PROTAC Linker 11) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.82%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p><b>Boc-NH-PEG2-C2-amido-C4-acid</b> (PROTAC Linker 30)</p> <p style="text-align: right;">Cat. No.: HY-125887</p> <p>Boc-NH-PEG2-C2-amido-C4-acid (PROTAC Linker 30) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG2-C2-NH2</b> (PROTAC Linker 13)</p> <p style="text-align: right;">Cat. No.: HY-W008474</p> <p>Boc-NH-PEG2-C2-NH2 (PROTAC Linker 13) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.01%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>Boc-NH-PEG2-C2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141117</p> <p>Boc-NH-PEG2-C2-NHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Boc-NH-PEG2-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W025812</p> <p>Boc-NH-PEG2-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>Boc-NH-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W005056</p> <p>Boc-NH-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.87%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Boc-NH-PEG2-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141198</p> <p>Boc-NH-PEG2-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>BOC-NH-PEG2-propene</b></p> <p style="text-align: right;">Cat. No.: HY-138398</p> <p>BOC-NH-PEG2-propene is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG20-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140473</p> <p>Boc-NH-PEG20-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG22-C2-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-138399</p> <p>Boc-NH-PEG22-C2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG23-CH2CH2N3</b></p> <p style="text-align: right;">Cat. No.: HY-130899</p> <p>Boc-NH-PEG23-CH2CH2N3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG23-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140236</p> <p>Boc-NH-PEG23-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG24-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140474</p> <p>Boc-NH-PEG24-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg</p>	<p><b>Boc-NH-PEG26-C2-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-138432</p> <p>Boc-NH-PEG26-C2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>






<p><b>Boc-NH-PEG3</b> (PROTAC Linker 10)</p> <p>Cat. No.: HY-W017772</p> <p>Boc-NH-PEG3 (PROTAC Linker 10) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p><b>Boc-NH-PEG3-C2-triazole-DBCO-PEG4-VC-PAB-DMEA</b></p> <p>Cat. No.: HY-126677</p> <p>Boc-NH-PEG3-C2-triazole-DBCO-PEG4-VC-PAB-DMEA is a double cleavable 3-unit and 4-unit PEG linker for antibody-drug-conjugation (ADC). Boc-NH-PEG3-C2-triazole-DBCO-PEG4-VC-PAB-DMEA also is a PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG3-CH2COOH</b></p> <p>Cat. No.: HY-33366</p> <p>Boc-NH-PEG3-CH2COOH is a PEG-based PROTAC linker can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>	<p><b>Boc-NH-PEG3-NHS ester</b></p> <p>Cat. No.: HY-141118</p> <p>Boc-NH-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG3-propargyl</b></p> <p>Cat. No.: HY-140878</p> <p>Boc-NH-PEG3-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG3-sulfonic acid</b></p> <p>Cat. No.: HY-140172</p> <p>Boc-NH-PEG3-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG36-CH2CH2COOH</b></p> <p>Cat. No.: HY-140475</p> <p>Boc-NH-PEG36-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG4</b> (PROTAC Linker 12)</p> <p>Cat. No.: HY-W025896</p> <p>Boc-NH-PEG4 (PROTAC Linker 12) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>Boc-NH-PEG4-azide</b></p> <p>Cat. No.: HY-140836</p> <p>Boc-NH-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG4-C2-Boc</b></p> <p>Cat. No.: HY-138370</p> <p>Boc-NH-PEG4-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Boc-NH-PEG4-C3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140478</p>	<p><b>Boc-NH-PEG4-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W040132</p>
<p>Boc-N-amido-PEG4-C3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG4-CH2CH2COOH is a PEG-based PROTAC linker can be used in the synthesis of PROTAC. Boc-NH-PEG4-CH2CH2COOH is also a cleavable ADC linker used as a linker for antibody-drug conjugates (ADC).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Boc-NH-PEG4-CH2CH2NH2</b></p> <p style="text-align: right;">Cat. No.: HY-W008352</p>	<p><b>Boc-NH-PEG4-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-42640</p>
<p>Boc-NH-PEG4-CH2CH2NH2 a cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-NH-PEG4-CH2CH2NH2 is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Boc-NH-PEG4-CH2COOH is a cleavable ADC linker used as a linker for antibody-drug conjugates (ADC). Boc-NH-PEG4-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Boc-NH-PEG4-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-140391</p>	<p><b>Boc-NH-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141119</p>
<p>Boc-NH-PEG4-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140837</p>	<p><b>Boc-NH-PEG5-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W022232</p>
<p>Boc-NH-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG5-CH2CH2COOH is a PEG-based PROTAC linker can be used in the synthesis of PROTAC.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG5-Cl</b></p> <p style="text-align: right;">Cat. No.: HY-134709</p>	<p><b>Boc-NH-PEG5-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140880</p>
<p>Boc-NH-PEG5-Cl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG5-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Boc-NH-PEG6-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140231</p> <p>Boc-NH-PEG6-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG6-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140838</p> <p>Boc-NH-PEG6-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG6-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W040244</p> <p>Boc-NH-PEG6-CH2CH2COOH is a cleavable ADC linker used as a linker for antibody-drug conjugates (ADC). Boc-NH-PEG6-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG6-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140477</p> <p>Boc-NH-PEG6-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG6-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140881</p> <p>Boc-NH-PEG6-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG7-acetic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W190963</p> <p>Boc-NH-PEG7-acetic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG7-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140469</p> <p>Boc-NH-PEG7-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Boc-NH-PEG7-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140839</p> <p>Boc-NH-PEG7-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG7-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140232</p> <p>Boc-NH-PEG7-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 5 mg, 10 mg, 50 mg, 100 mg</p>	<p><b>Boc-NH-PEG7-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140882</p> <p>Boc-NH-PEG7-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>













<p><b>Boc-NH-PEG7-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140376</p>	<p><b>Boc-NH-PEG8-C2-Br</b></p> <p style="text-align: right;">Cat. No.: HY-138406</p>
<p>Boc-NH-PEG7-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG8-C2-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG8-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140470</p>	<p><b>Boc-NH-PEG8-CH2CH2NH2</b></p> <p style="text-align: right;">Cat. No.: HY-135933</p>
<p>Boc-NH-PEG8-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG8-CH2CH2NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>Boc-NH-PEG8-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140883</p>	<p><b>Boc-NH-PEG9-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140840</p>
<p>Boc-NH-PEG8-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PEG9-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NH-PEG9-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140884</p>	<p><b>Boc-NH-PPG2</b></p> <p style="text-align: right;">Cat. No.: HY-138492</p>
<p>Boc-NH-PEG9-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-NH-PPG2 is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-NHCH2CH2-PEG1-azide</b></p> <p style="text-align: right;">Cat. No.: HY-132119</p>	<p><b>Boc-PEG1-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138473</p>
<p>Boc-NHCH2CH2-PEG1-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-PEG1-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Boc-PEG1-PPG2-C2-azido</b></p> <p style="text-align: right;">Cat. No.: HY-138474</p>	<p><b>Boc-PEG1-PPG2-C2-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-138485</p>
<p>Boc-PEG1-PPG2-C2-azido is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-PEG1-PPG2-C2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-PEG2-ethoxyethane-PEG2-benzyl</b></p> <p style="text-align: right;">Cat. No.: HY-138475</p>	<p><b>Boc-PEG2-sulfonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-140173</p>
<p>Boc-PEG2-ethoxyethane-PEG2-benzyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-PEG2-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-PEG25-benzyl</b></p> <p style="text-align: right;">Cat. No.: HY-138468</p>	<p><b>Boc-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138379</p>
<p>Boc-PEG25-benzyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-PEG4-C2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141122</p>	<p><b>Boc-PEG4-methyl propionate</b></p> <p style="text-align: right;">Cat. No.: HY-138481</p>
<p>Boc-PEG4-C2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-PEG4-methyl propionate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-PEG4-phosphonic acid ethyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141304</p>	<p><b>Boc-PEG4-sulfone-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140613</p>
<p>Boc-PEG4-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-PEG4-sulfone-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>








<p><b>Boc-PEG4-sulfonic acid</b></p> <p>Cat. No.: HY-140174</p>	<p><b>Boc-PEG5-methyl ester</b></p> <p>Cat. No.: HY-138373</p>
<p>Boc-PEG4-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-PEG5-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-PEG8-Boc</b></p> <p>Cat. No.: HY-138335</p>	<p><b>Boc-Pip-alkyne-Ph-COOH</b></p> <p>Cat. No.: HY-133044</p>
<p>Boc-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Boc-Pip-alkyne-Ph-COOH is a PROTAC linker, which refers to the alkyl/ether composition. Boc-Pip-alkyne-Ph-COOH can be used in the synthesis of a series of PROTACs, such as ARD-266 (HY-133020).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-piperazine-benzoic acid</b></p> <p>Cat. No.: HY-W013249</p>	<p><b>Boc-Pyrrolidine-PEG2-COOH</b></p> <p>Cat. No.: HY-134744</p>
<p>Boc-piperazine-benzoic acid is a PROTAC linker and can be used in the synthesis of PROTACs, such as PROTAC androgen receptor (AR) degrader ARD-2128 (HY-13229).</p>  <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 500 mg, 1 g</p>	<p>Boc-Pyrrolidine-PEG2-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Boc-trans-D-Hyp-OMe</b></p> <p>Cat. No.: HY-W017882</p>	<p><b>BocNH-PEG5-CH2CH2Br</b></p> <p>Cat. No.: HY-W096092</p>
<p>Boc-trans-D-Hyp-OMe is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Boc-trans-D-Hyp-OMe is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 250 mg</p>	<p>BocNH-PEG5-CH2CH2Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Br-Boc-C2-azido</b></p> <p>Cat. No.: HY-138464</p>	<p><b>Br-C10-methyl ester</b></p> <p>Cat. No.: HY-130641</p>
<p>Br-Boc-C2-azido is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Br-C10-methyl ester is a PROTAC linker, which refers to the alkyl/ether composition. Br-C10-methyl ester is used in the synthesis of a series of PROTACs (MS432).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>

<p><b>Br-C3-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-W004701</p> <p>Br-C3-methyl ester is a Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTAC PD-1/PD-L1 degrader-1 (HY-131183).</p>  <p><b>Purity:</b> 96.86%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>	<p><b>Br-PEG2-oxazolidin-2-one</b></p> <p style="text-align: right;">Cat. No.: HY-134740</p> <p>Br-PEG2-oxazolidin-2-one is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Br-PEG3-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130483</p> <p>Br-PEG3-C2-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>Br-PEG3-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130500</p> <p>Br-PEG3-CH2COOH (compound 28) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Br-PEG3-ethyl acetate</b></p> <p style="text-align: right;">Cat. No.: HY-W096139</p> <p>Br-PEG3-ethyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Br-PEG3-MS</b></p> <p style="text-align: right;">Cat. No.: HY-138336</p> <p>Br-PEG3-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Br-PEG3-OH</b></p> <p style="text-align: right;">Cat. No.: HY-135142</p> <p>Br-PEG3-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Br-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138486</p> <p>Br-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Br-PEG4-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130315</p> <p>Br-PEG4-C2-Boc is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 250 mg</p>	<p><b>Br-PEG4-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130536</p> <p>Br-PEG4-CH2-Boc is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Br-PEG4-methyl acetate</b></p> <p style="text-align: right;">Cat. No.: HY-138450</p>	<p><b>Br-PEG4-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130512</p>
<p>Br-PEG4-methyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Br-PEG4-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Br-PEG4-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138372</p>	<p><b>Br-PEG6-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138362</p>
<p>Br-PEG4-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Br-PEG6-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Br-PEG6-C2-NHBoc</b></p> <p style="text-align: right;">Cat. No.: HY-138381</p>	<p><b>Br-PEG6-CH2COOtBu</b></p> <p style="text-align: right;">Cat. No.: HY-W190969</p>
<p>Br-PEG6-C2-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Br-PEG6-CH2COOtBu is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Br-PEG7-Br</b></p> <p style="text-align: right;">Cat. No.: HY-133299</p>	<p><b>Br-PEG7-NHBoc</b></p> <p style="text-align: right;">Cat. No.: HY-138515</p>
<p>Br-PEG7-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Br-PEG7-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Br-PEG9-C2-NHBoc</b></p> <p style="text-align: right;">Cat. No.: HY-138384</p>	<p><b>Bromo-C10-OBn</b></p> <p style="text-align: right;">Cat. No.: HY-132859</p>
<p>Br-PEG9-C2-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-C10-OBn is a PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>



<p><b>Bromo-C4-PEG4-t-butyl ester</b></p> <p>Cat. No.: HY-144079</p>	<p><b>Bromo-PEG1-acid</b></p> <p>Cat. No.: HY-130425</p>
<p>Bromo-C4-PEG4-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG1-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG1-C2-azide</b></p> <p>Cat. No.: HY-140819</p>	<p><b>Bromo-PEG1-C2-Boc</b></p> <p>Cat. No.: HY-141364</p>
<p>Bromo-PEG1-C2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG1-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.58%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Bromo-PEG1-CH2-Boc</b></p> <p>Cat. No.: HY-141370</p>	<p><b>Bromo-PEG1-CH2COOH</b></p> <p>Cat. No.: HY-130163</p>
<p>Bromo-PEG1-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Bromo-PEG1-CH2COOH is a PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG1-NH2 hydrobromide</b></p> <p>Cat. No.: HY-134736</p>	<p><b>Bromo-PEG10-t-butyl ester</b></p> <p>Cat. No.: HY-132118</p>
<p>Bromo-PEG1-NH2 hydrobromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG10-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG12-acid</b></p> <p>Cat. No.: HY-134754</p>	<p><b>Bromo-PEG12-t-butyl ester</b></p> <p>Cat. No.: HY-132117</p>
<p>Bromo-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG12-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Bromo-PEG2-acetic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W190843</p> <p>Bromo-PEG2-acetic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bromo-PEG2-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141213</p> <p>Bromo-PEG2-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Bromo-PEG2-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-130589</p> <p>Bromo-PEG2-bromide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 97.78%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p><b>Bromo-PEG2-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141362</p> <p>Bromo-PEG2-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG2-C2-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130485</p> <p>Bromo-PEG2-C2-azide is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Bromo-PEG2-C2-azide is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.10%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p><b>Bromo-PEG2-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141365</p> <p>Bromo-PEG2-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg</p>
<p><b>Bromo-PEG2-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141371</p> <p>Bromo-PEG2-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bromo-PEG2-MS</b></p> <p style="text-align: right;">Cat. No.: HY-132042</p> <p>Bromo-PEG2-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG2-NH2 hydrobromide</b></p> <p style="text-align: right;">Cat. No.: HY-134694</p> <p>Bromo-PEG2-NH2 hydrobromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bromo-PEG2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-132018</p> <p>Bromo-PEG2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>

<p><b>Bromo-PEG2-phosphonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-141297</p>	<p><b>Bromo-PEG2-phosphonic acid diethyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141299</p>
<p>Bromo-PEG2-phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG2-phosphonic acid diethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG2-THP</b></p> <p style="text-align: right;">Cat. No.: HY-W096071</p>	<p><b>Bromo-PEG24-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141369</p>
<p>Bromo-PEG2-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG24-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140820</p>	<p><b>Bromo-PEG3-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-141373</p>
<p>Bromo-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG3-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> 98.10%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg, 250 mg, 500 mg</p>
<p><b>Bromo-PEG3-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W039168</p>	<p><b>Bromo-PEG3-C2-phosphonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-130164</p>
<p>Bromo-PEG3-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Bromo-PEG3-C2-phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG3-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141372</p>	<p><b>Bromo-PEG3-CO-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-134708</p>
<p>Bromo-PEG3-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>Bromo-PEG3-CO-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>






<p><b>Bromo-PEG5-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141366</p>	<p><b>Bromo-PEG5-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-141374</p>
<p>Bromo-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG5-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG5-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130489</p>	<p><b>Bromo-PEG5-CH2COOtBu</b></p> <p style="text-align: right;">Cat. No.: HY-144080</p>
<p>Bromo-PEG5-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG5-CH2COOtBu is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG5-phosphonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-141298</p>	<p><b>Bromo-PEG5-phosphonic acid diethyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141301</p>
<p>Bromo-PEG5-phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG5-phosphonic acid diethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG6-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141215</p>	<p><b>Bromo-PEG6-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140823</p>
<p>Bromo-PEG6-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG6-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG6-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141367</p>	<p><b>Bromo-PEG6-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-135049</p>
<p>Bromo-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG6-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Bromo-PEG7-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-132087</p>	<p><b>Bromo-PEG7-amine</b></p> <p style="text-align: right;">Cat. No.: HY-143844</p>
<p>Bromo-PEG7-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG7-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG7-azide</b></p> <p style="text-align: right;">Cat. No.: HY-W096078</p>	<p><b>Bromo-PEG7-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-134677</p>
<p>Bromo-PEG7-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG7-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG7-CH2COOtBu</b></p> <p style="text-align: right;">Cat. No.: HY-144077</p>	<p><b>Bromo-PEG8-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141368</p>
<p>Bromo-PEG7-CH2COOtBu is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromo-PEG8-CH2COOtBu</b></p> <p style="text-align: right;">Cat. No.: HY-144078</p>	<p><b>Bromo-PEG9-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-134755</p>
<p>Bromo-PEG8-CH2COOtBu is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromo-PEG9-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromoacetamide-PEG3-C1-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133506</p>	<p><b>Bromoacetamide-PEG3-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-130942</p>
<p>Bromoacetamide-PEG3-C1-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Bromoacetamide-PEG3-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Bromoacetamido-C2-PEG2-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141199</p> <p>Bromoacetamido-C2-PEG2-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bromoacetamido-PEG2-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141380</p> <p>Bromoacetamido-PEG2-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.11% <b>Clinical Data:</b> <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Bromoacetamido-PEG2-C2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141386</p> <p>Bromoacetamido-PEG2-C2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bromoacetamido-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140825</p> <p>Bromoacetamido-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromoacetamido-PEG3-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141381</p> <p>Bromoacetamido-PEG3-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bromoacetamido-PEG3-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141388</p> <p>Bromoacetamido-PEG3-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromoacetamido-PEG3-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141200</p> <p>Bromoacetamido-PEG3-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bromoacetamido-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141382</p> <p>Bromoacetamido-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Bromoacetamido-PEG4-acid is also a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Bromoacetamido-PEG4-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130520</p> <p>Bromoacetamido-PEG4-C2-Boc is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Bromoacetamido-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141387</p> <p>Bromoacetamido-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

**Bromoacetamido-PEG5-azide** Cat. No.: HY-140826


Bromoacetamido-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**Bromoacetamido-PEG5-DOTA** Cat. No.: HY-140754


Bromoacetamido-PEG5-DOTA is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**Bromoacetamido-PEG8-acid** Cat. No.: HY-141383


Bromoacetamido-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**Bromoacetamido-PEG8-Boc** Cat. No.: HY-141389


Bromoacetamido-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Bromoacetamido-PEG8-t-butyl acetate** Cat. No.: HY-132084


Bromoacetamido-PEG8-t-butyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Bromoacetamido-PEG9-ethylcarbamoyl-C12-Boc** Cat. No.: HY-141392


Bromoacetamido-PEG9-ethylcarbamoyl-C12-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**Bromoacetamido-PEG9-ethylcarbamoyl-C4-Boc** Cat. No.: HY-141390


Bromoacetamido-PEG9-ethylcarbamoyl-C4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**Bromoacetamido-PEG9-ethylcarbamoyl-C8-Boc** Cat. No.: HY-141391


Bromoacetamido-PEG9-ethylcarbamoyl-C8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**Bromoacetic-PEG1-CH2-NHS ester** Cat. No.: HY-138414


Bromoacetic-PEG1-CH2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.













**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg











**Bromoacetic-PEG2-NHS ester** Cat. No.: HY-138412







Bromoacetic-PEG2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.













**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg





<p><b>Bromoacetyl-PEG3-DBCO</b></p> <p>Cat. No.: HY-133480</p>	<p><b>Butane-1,4-diyldiphosphonic acid</b></p> <p>Cat. No.: HY-W075674</p>
<p>Bromoacetyl-PEG3-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>Butane-1,4-diyldiphosphonic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p><b>Butoxycarbonyl-PEG5-sulfonic acid</b></p> <p>Cat. No.: HY-140175</p>	<p><b>C-NH-Boc-C-Bis-(C-PEG1-Boc)</b></p> <p>Cat. No.: HY-140336</p>
<p>Butoxycarbonyl-PEG5-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>C-NH-Boc-C-Bis-(C-PEG1-Boc) is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>
<p><b>C-NH-Boc-C-Bis-(C1-PEG1-PFP)</b></p> <p>Cat. No.: HY-141259</p>	<p><b>C18-PEG13-acid</b></p> <p>Cat. No.: HY-144082</p>
<p>C-NH-Boc-C-Bis-(C1-PEG1-PFP) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>	<p>C18-PEG13-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>
<p><b>C2-Bis-phosphoramidic acid diethyl ester</b></p> <p>Cat. No.: HY-141323</p>	<p><b>C6-Bis-phosphoramidic acid diethyl ester</b></p> <p>Cat. No.: HY-141325</p>
<p>C2-Bis-phosphoramidic acid diethyl ester is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>C6-Bis-phosphoramidic acid diethyl ester is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>
<p><b>Carboxy-PEG2-sulfonic acid</b></p> <p>Cat. No.: HY-140167</p>	<p><b>Carboxy-PEG4-phosphonic acid</b></p> <p>Cat. No.: HY-141302</p>
<p>Carboxy-PEG2-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: No Development Reported Size: 1 mg, 5 mg</p>	<p>Carboxy-PEG4-phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Purity: &gt;98% Clinical Data: Size: 1 mg, 5 mg</p>











<p><b>Carboxy-PEG4-phosphonic acid ethyl ester</b></p> <p>Cat. No.: HY-141303</p>	<p><b>Carboxy-PEG4-sulfonic acid</b></p> <p>Cat. No.: HY-140168</p>
<p>Carboxy-PEG4-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Carboxy-PEG4-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Carboxy-PEG5-sulfonic acid</b></p> <p>Cat. No.: HY-140169</p>	<p><b>Carboxyfluorescein-PEG12-NHS</b></p> <p>Cat. No.: HY-141085</p>
<p>Carboxy-PEG5-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Carboxyfluorescein-PEG12-NHS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 5 mg</p>
<p><b>Carboxymethyl-PEG-Carboxymethyl (MW 5000)</b></p> <p>Cat. No.: HY-140723</p>	<p><b>Carboxyrhodamine 110-PEG3-Azide</b></p> <p>Cat. No.: HY-141090</p>
<p>Carboxymethyl-PEG-Carboxymethyl (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Carboxyrhodamine 110-PEG3-Azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Carboxyrhodamine 110-PEG4-alkyne</b></p> <p>Cat. No.: HY-141089</p>	<p><b>Carboxyrhodamine 110-PEG4-DBCO</b></p> <p>Cat. No.: HY-140297</p>
<p>Carboxyrhodamine 110-PEG4-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Carboxyrhodamine 110-PEG4-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cbz-aminoxy-PEG8-acid</b></p> <p>Cat. No.: HY-140443</p>	<p><b>Cbz-aminoxy-PEG8-Boc</b></p> <p>Cat. No.: HY-140444</p>
<p>Cbz-aminoxy-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-aminoxy-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>

<p><b>Cbz-N-amido-PEG20-acid</b></p> <p>Cat. No.: HY-140491</p>	<p><b>Cbz-N-PEG10-acid</b></p> <p>Cat. No.: HY-140490</p>
<p>Cbz-N-amido-PEG20-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-N-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cbz-N-PEG15-amine</b></p> <p>Cat. No.: HY-140238</p>	<p><b>Cbz-NH-PEG1-CH2CH2COOH</b></p> <p>Cat. No.: HY-135926</p>
<p>Cbz-N-PEG15-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-NH-PEG1-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cbz-NH-PEG1-CH2COOH</b></p> <p>Cat. No.: HY-135925</p>	<p><b>Cbz-NH-PEG10-CH2COOH</b></p> <p>Cat. No.: HY-133357</p>
<p>Cbz-NH-PEG1-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.64% <b>Clinical Data:</b> <b>Size:</b> 250 mg, 500 mg</p>	<p>Cbz-NH-PEG10-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cbz-NH-PEG12-C2-acid</b></p> <p>Cat. No.: HY-133358</p>	<p><b>Cbz-NH-PEG2-C2-acid</b></p> <p>Cat. No.: HY-140487</p>
<p>Cbz-NH-PEG12-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-NH-PEG2-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cbz-NH-PEG2-CH2COOH</b></p> <p>Cat. No.: HY-135923</p>	<p><b>Cbz-NH-PEG24-C2-acid</b></p> <p>Cat. No.: HY-133359</p>
<p>Cbz-NH-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-NH-PEG24-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Cbz-NH-PEG3-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140488</p>	<p><b>Cbz-NH-PEG3-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-135913</p>
<p>Cbz-NH-PEG3-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-NH-PEG3-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cbz-NH-PEG36-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133360</p>	<p><b>Cbz-NH-PEG4-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W019796</p>
<p>Cbz-NH-PEG36-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-NH-PEG4-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cbz-NH-PEG5-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140489</p>	<p><b>Cbz-NH-PEG5-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-135919</p>
<p>Cbz-NH-PEG5-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-NH-PEG5-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cbz-NH-PEG6-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130443</p>	<p><b>Cbz-NH-PEG8-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130192</p>
<p>Cbz-NH-PEG6-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-NH-PEG8-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cbz-NH-PEG8-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-135942</p>	<p><b>Cbz-PEG2-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-W190968</p>
<p>Cbz-NH-PEG8-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Cbz-PEG2-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Cbz-PEG5-Br</b></p> <p>Cat. No.: HY-143832</p>	<p><b>CG-PEG5-azido</b></p> <p>Cat. No.: HY-138516</p>
<p>Cbz-PEG5-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>CG-PEG5-azido is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>CH2COOH-PEG12-CH2COOH</b></p> <p>Cat. No.: HY-132113</p>	<p><b>CH2COOH-PEG3-CH2COOH</b></p> <p>Cat. No.: HY-122696</p>
<p>CH2COOH-PEG12-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>CH2COOH-PEG3-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>CH2COOH-PEG6-CH2COOH</b></p> <p>Cat. No.: HY-W096106</p>	<p><b>CH2COOH-PEG9-CH2COOH</b></p> <p>Cat. No.: HY-138514</p>
<p>CH2COOH-PEG6-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>CH2COOH-PEG9-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Chloro-PEG2-Boc</b></p> <p>Cat. No.: HY-W096145</p>	<p><b>Chloro-PEG5-chloride</b></p> <p>Cat. No.: HY-W096131</p>
<p>Chloro-PEG2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Chloro-PEG5-chloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Chloroacetamido-C-PEG3-C3-NHBoc</b></p> <p>Cat. No.: HY-W096151</p>	<p><b>Chloroacetamido-C4-NHBoc</b></p> <p>Cat. No.: HY-W096152</p>
<p>Chloroacetamido-C-PEG3-C3-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Chloroacetamido-C4-NHBoc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>


<p><b>Chloroacetamido-PEG4-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130470</p>	<p><b>Chloroacetamido-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141394</p>
<p>Chloroacetamido-PEG4-C2-Boc is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Chloroacetamido-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>CHO-C-PEG2-C-CHO</b></p> <p style="text-align: right;">Cat. No.: HY-138383</p>	<p><b>CHO-CH2-PEG1-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138355</p>
<p>CHO-C-PEG2-C-CHO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>CHO-CH2-PEG1-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>CHO-PEG12-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-134707</p>	<p><b>cis-4-Hydroxy-D-proline hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-76104</p>
<p>CHO-PEG12-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>cis-4-Hydroxy-D-proline hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). cis-4-Hydroxy-D-proline hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 g, 5 g</p>
<p><b>cis-4-Hydroxy-L-proline hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-W019213</p>	<p><b>Cl-C6-PEG4-C3-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-135306</p>
<p>cis-4-Hydroxy-L-proline hydrochloride is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). cis-4-Hydroxy-L-proline hydrochloride is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>	<p>Cl-C6-PEG4-C3-COOH is a PROTAC linker can be used in the synthesis of chloroalkane-containing PROTACs (HaloPROTACs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cl-C6-PEG4-O-CH2COOH (PROTAC Linker 4)</b></p> <p style="text-align: right;">Cat. No.: HY-112496</p>	<p><b>Cl-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-43055</p>
<p>Cl-C6-PEG4-O-CH2COOH (PROTAC Linker 4) is a PEG-based PROTAC linker can be used in the synthesis of chloroalkane-containing PROTACs (HaloPROTACs).</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p>Cl-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>

<p><b>Cl-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W096129</p>	<p><b>Cl-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138411</p>
<p>Cl-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Cl-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cy5-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-141058</p>	<p><b>Cy5-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141033</p>
<p>Cy5-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 5 mg, 10 mg</p>	<p>Cy5-PEG4-acid (chloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cy5-PEG5-amine hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-141055</p>	<p><b>Cy5-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-141059</p>
<p>Cy5-PEG5-amine (hydrochloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Cy5-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cy5-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141034</p>	<p><b>Cy5-PEG6-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141044</p>
<p>Cy5-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Cy5-PEG6-NHS ester (chloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Cyclohexane-PEG1-Br</b></p> <p style="text-align: right;">Cat. No.: HY-W092895</p>	<p><b>Cystamine</b></p> <p style="text-align: right;">Cat. No.: HY-124476</p>
<p>Cyclohexane-PEG1-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Cystamine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 96.78%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>










<p><b>D-Proline, 4-hydroxy-, methyl ester hydrochloride</b></p> <p>Cat. No.: HY-76105</p> <p>D-Proline, 4-hydroxy-, methyl ester hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-(PEG2-Val-Cit-PAB)2</b></p> <p>Cat. No.: HY-126676</p> <p>DBCO-(PEG2-Val-Cit-PAB)2 is a dual cleavable ADC linker for antibody-drug conjugates (ADCs). DBCO-(PEG2-Val-Cit-PAB)2 is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-Biotin</b></p> <p>Cat. No.: HY-123916</p> <p>DBCO-Biotin is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>DBCO-C-PEG1</b></p> <p>Cat. No.: HY-140281</p> <p>DBCO-C-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-C2-PEG4-amine</b></p> <p>Cat. No.: HY-140287</p> <p>DBCO-C2-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-C2-PEG4-NH-Boc</b></p> <p>Cat. No.: HY-140294</p> <p>DBCO-C2-PEG4-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-C2-SulfoNHS ester</b></p> <p>Cat. No.: HY-133509</p> <p>DBCO-C2-SulfoNHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-C3-PEG4-amine</b></p> <p>Cat. No.: HY-140288</p> <p>DBCO-C3-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-C3-PEG4-NH-Boc</b></p> <p>Cat. No.: HY-140295</p> <p>DBCO-C3-PEG4-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-mPEG (MW 10kDa)</b></p> <p>Cat. No.: HY-140320</p> <p>DBCO-mPEG (MW 10kDa) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>DBCO-mPEG (MW 20kDa)</b></p> <p style="text-align: right;">Cat. No.: HY-140321</p> <p>DBCO-mPEG (MW 20kDa) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 20kDa</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-mPEG (MW 2kDa)</b></p> <p style="text-align: right;">Cat. No.: HY-140318</p> <p>DBCO-mPEG (MW 2kDa) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 2kDa</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-mPEG (MW 30kDa)</b></p> <p style="text-align: right;">Cat. No.: HY-140322</p> <p>DBCO-mPEG (MW 30kDa) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 30kDa</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg</p>	<p><b>DBCO-mPEG (MW 5kDa)</b></p> <p style="text-align: right;">Cat. No.: HY-140319</p> <p>DBCO-mPEG (MW 5kDa) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 5kDa</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-N-bis(PEG4-NHS ester)</b></p> <p style="text-align: right;">Cat. No.: HY-145090</p> <p>DBCO-N-bis(PEG4-NHS ester) is a PEG linker which contains two PEG4-NHS ester and a DBCO group. DBCO-N-bis(PEG4-NHS ester) is useful for protein modification or labeling.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-NH-(CH<sub>2</sub>)<sub>4</sub>COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130912</p> <p>DBCO-NH-(CH<sub>2</sub>)<sub>4</sub>COOH is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140293</p> <p>DBCO-NH-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-NH-PEG7-C2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-135959</p> <p>DBCO-NH-PEG7-C2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-NHCO-C4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-133371</p> <p>DBCO-NHCO-C4-NHS ester is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-NHCO-PEG12-amine</b></p> <p style="text-align: right;">Cat. No.: HY-133368</p> <p>DBCO-NHCO-PEG12-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>DBCO-NHCO-PEG12-biotin</b></p> <p style="text-align: right;">Cat. No.: HY-133376</p>	<p><b>DBCO-NHCO-PEG12-maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-133374</p>
<p>DBCO-NHCO-PEG12-biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-NHCO-PEG12-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 100 mg</p>
<p><b>DBCO-NHCO-PEG13-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140270</p>	<p><b>DBCO-NHCO-PEG13-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140276</p>
<p>DBCO-NHCO-PEG13-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-NHCO-PEG13-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-NHCO-PEG2-amine</b></p> <p style="text-align: right;">Cat. No.: HY-133366</p>	<p><b>DBCO-NHCO-PEG2-Biotin</b></p> <p style="text-align: right;">Cat. No.: HY-135939</p>
<p>DBCO-NHCO-PEG2-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-NHCO-PEG2-Biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-NHCO-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-135935</p>	<p><b>DBCO-NHCO-PEG2-maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-133372</p>
<p>DBCO-NHCO-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-NHCO-PEG2-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-NHCO-PEG2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-133375</p>	<p><b>DBCO-NHCO-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133369</p>
<p>DBCO-NHCO-PEG2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-NHCO-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

<p><b>DBCO-NHCO-PEG3-Fmoc</b></p> <p style="text-align: right;">Cat. No.: HY-133472</p> <p>DBCO-NHCO-PEG3-Fmoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-NHCO-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-125541</p> <p>DBCO-Amide-PEG5-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-Amide-PEG5-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-NHCO-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-124386</p> <p>DBCO-NHCO-PEG4-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-NHCO-PEG4-amine is a cleavable ADC linker used to conjugate MMAE (HY-15162) and antibody (e.g.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-NHCO-PEG4-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-126884</p> <p>DBCO-NHCO-PEG4-NH-Boc is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-NHCO-PEG4-NH-Boc is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-NHCO-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-111456</p> <p>DBCO-NHCO-PEG4-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-NHCO-PEG4-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-NHCO-PEG5-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140275</p> <p>DBCO-NHCO-PEG5-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-NHCO-PEG6-amine</b></p> <p style="text-align: right;">Cat. No.: HY-133367</p> <p>DBCO-NHCO-PEG6-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-NHCO-PEG6-Biotin</b></p> <p style="text-align: right;">Cat. No.: HY-135958</p> <p>DBCO-NHCO-PEG6-Biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-NHCO-PEG6-maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-133373</p> <p>DBCO-NHCO-PEG6-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-NHCO-PEG7-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133370</p> <p>DBCO-NHCO-PEG7-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

















<p><b>DBCO-PEG1</b></p> <p style="text-align: right;">Cat. No.: HY-140280</p>	<p><b>DBCO-PEG1-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140265</p>
<p>DBCO-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG1-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG1-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140282</p>	<p><b>DBCO-PEG1-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140289</p>
<p>DBCO-PEG1-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG1-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG1-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140271</p>	<p><b>DBCO-PEG10-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140304</p>
<p>DBCO-PEG1-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG10-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140269</p>	<p><b>DBCO-PEG12-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140274</p>
<p>DBCO-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG13-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140305</p>	<p><b>DBCO-PEG13-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-138744</p>
<p>DBCO-PEG13-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG13-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>




<p><b>DBCO-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140266</p>	<p><b>DBCO-PEG2-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140283</p>
<p>DBCO-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG2-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG2-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140290</p>	<p><b>DBCO-PEG3-amide-N-Fmoc</b></p> <p style="text-align: right;">Cat. No.: HY-138318</p>
<p>DBCO-PEG2-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG3-amide-N-Fmoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG3-amine</b></p> <p style="text-align: right;">Cat. No.: HY-134714</p>	<p><b>DBCO-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-120678</p>
<p>DBCO-PEG3-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG4-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140279</p>	<p><b>DBCO-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-130435</p>
<p>DBCO-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg, 250 mg, 500 mg</p>	<p>DBCO-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. DBCO-PEG4-amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>
<p><b>DBCO-PEG4-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130396</p>	<p><b>DBCO-PEG4-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-130346</p>
<p>DBCO-PEG4-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>DBCO-PEG4-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. DBCO-PEG4-DBCO is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>DBCO-PEG4-Desthiobiotin</b></p> <p style="text-align: right;">Cat. No.: HY-140301</p>	<p><b>DBCO-PEG4-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140291</p>
<p>DBCO-PEG4-Desthiobiotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG4-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140272</p>	<p><b>DBCO-PEG4-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140278</p>
<p>DBCO-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 97.97%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg</p>	<p>DBCO-PEG4-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>DBCO-PEG4-triethoxysilane</b></p> <p style="text-align: right;">Cat. No.: HY-140308</p>	<p><b>DBCO-PEG5-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140267</p>
<p>DBCO-PEG4-triethoxysilane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG5-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140302</p>	<p><b>DBCO-PEG5-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-126885</p>
<p>DBCO-PEG5-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>DBCO-PEG5-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. DBCO-PEG5-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>DBCO-PEG6-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140284</p>	<p><b>DBCO-PEG6-amine TFA</b></p> <p style="text-align: right;">Cat. No.: HY-140284A</p>
<p>DBCO-PEG6-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 50 mg, 100 mg</p>	<p>DBCO-PEG6-amine TFA is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>DBCO-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140268</p> <p>DBCO-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-PEG8-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140273</p> <p>DBCO-PEG8-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 50 mg, 100 mg</p>
<p><b>DBCO-PEG9-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140285</p> <p>DBCO-PEG9-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 95.90%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg</p>	<p><b>DBCO-PEG9-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140303</p> <p>DBCO-PEG9-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DBCO-PEG9-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140292</p> <p>DBCO-PEG9-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DBCO-S-S-PEG3-biotin</b></p> <p style="text-align: right;">Cat. No.: HY-140128</p> <p>DBCO-S-S-PEG3-biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 95.07%  <b>Clinical Data:</b>  <b>Size:</b> 5 mg, 10 mg, 50 mg, 100 mg</p>
<p><b>Dde Biotin-PEG4-alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-140924</p> <p>Dde Biotin-PEG4-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 5 mg, 10 mg</p>	<p><b>Dde Biotin-PEG4-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140916</p> <p>Dde Biotin-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Dde Biotin-PEG4-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140931</p> <p>Dde Biotin-PEG4-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Dde Biotin-PEG4-Picolyl azide</b></p> <p style="text-align: right;">Cat. No.: HY-140917</p> <p>Dde Biotin-PEG4-Picolyl azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Dde Biotin-PEG4-TAMRA-PEG4 Alkyne</b></p> <p>Cat. No.: HY-140877</p> <p>Dde Biotin-PEG4-TAMRA-PEG4 Alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Decaethylene glycol</b> (HO-PEG10-OH)</p> <p>Cat. No.: HY-141232</p> <p>Decaethylene glycol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.09%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg</p>
<p><b>Decaethylene glycol dodecyl ether</b> (PEG10-dodecyl)</p> <p>Cat. No.: HY-132065</p> <p>Decaethylene glycol dodecyl ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Desthiobiotin-PEG4-acid</b></p> <p>Cat. No.: HY-138507</p> <p>Desthiobiotin-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 95.91%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 250 mg</p>
<p><b>Desthiobiotin-PEG4-propargyl</b></p> <p>Cat. No.: HY-W096125</p> <p>Desthiobiotin-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Di(N-succinimidyl)adipate</b></p> <p>Cat. No.: HY-141099</p> <p>Di(N-succinimidyl)adipate is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Diazo Biotin-PEG3-alkyne</b></p> <p>Cat. No.: HY-140925</p> <p>Diazo Biotin-PEG3-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 10 mg, 25 mg, 50 mg, 100 mg</p>	<p><b>Diazo Biotin-PEG3-azide</b></p> <p>Cat. No.: HY-140918</p> <p>Diazo Biotin-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Diethoxy-phosphorylethyl-PEG5-ethylphosphonic acid</b></p> <p>Cat. No.: HY-141305</p> <p>Diethoxy-phosphorylethyl-PEG5-ethylphosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Diethyl 10-bromodecylphosphonate</b></p> <p>Cat. No.: HY-140333</p> <p>Diethyl 10-bromodecylphosphonate is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Diethyl 12-bromododecylphosphonate</b></p> <p>Cat. No.: HY-140334</p>	<p><b>Diethyl 7-bromoheptylphosphonate</b></p> <p>Cat. No.: HY-W035417</p>
<p>Diethyl 12-bromododecylphosphonate is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Diethyl 7-bromoheptylphosphonate is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Diethyl 8-bromooctylphosphonate</b></p> <p>Cat. No.: HY-140332</p>	<p><b>Diethylene glycol bis(p-toluenesulfonate)</b> (Diethylene glycol di(p-toluenesulfonate); Bis-Tos-PEG2) Cat. No.: HY-W010948</p>
<p>Diethyl 8-bromooctylphosphonate is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Diethylene glycol bis(p-toluenesulfonate) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.44% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 500 mg</p>
<p><b>Diethylene glycol diacetate</b></p> <p>Cat. No.: HY-W096085</p>	<p><b>Diethylene Glycol Monobenzyl Ether</b></p> <p>Cat. No.: HY-22393</p>
<p>Diethylene glycol diacetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Diethylene Glycol Monobenzyl Ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>Diketone-PEG11-Diketone</b></p> <p>Cat. No.: HY-140451</p>	<p><b>Diketone-PEG11-PFP ester</b></p> <p>Cat. No.: HY-140447</p>
<p>Diketone-PEG11-Diketone is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Diketone-PEG11-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Diketone-PEG12-Biotin</b></p> <p>Cat. No.: HY-140449</p>	<p><b>Diketone-PEG12-DBCO</b></p> <p>Cat. No.: HY-140450</p>
<p>Diketone-PEG12-Biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Diketone-PEG12-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Diketone-PEG4-Biotin</b></p> <p style="text-align: right;">Cat. No.: HY-140448</p> <p>Diketone-PEG4-Biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Diketone-PEG4-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140006</p> <p>Diketone-PEG4-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Dimethylamine-PEG19</b></p> <p style="text-align: right;">Cat. No.: HY-138401</p> <p>Dimethylamine-PEG19 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Dimethylamino-PEG11</b></p> <p style="text-align: right;">Cat. No.: HY-138409</p> <p>Dimethylamino-PEG11 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Dimethylamino-PEG2-C2-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-W096132</p> <p>Dimethylamino-PEG2-C2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Dimethylamino-PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-W096150</p> <p>Dimethylamino-PEG3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Dioxoisindolin-O-PEG-OH (MW 2000)</b>            ((1,3-dioxoisindolin-2-yl)-O-PEG-OH (MW 2000))</p> <p style="text-align: right;">Cat. No.: HY-140732</p> <p>Dioxoisindolin-O-PEG-OH (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Dioxoisindolin-O-PEG-OMe (MW 2000)</b>            ((1,3-dioxoisindolin-2-yl)-O-PEG-OMe (MW 2000))</p> <p style="text-align: right;">Cat. No.: HY-140733</p> <p>Dioxoisindolin-O-PEG-OMe (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DNP-NH-PEG2-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130467</p> <p>DNP-NH-PEG2-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DNP-NH-PEG4-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130491</p> <p>DNP-NH-PEG4-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>





<p><b>DNP-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140496</p> <p>DNP-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DNP-PEG12-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140615</p> <p>DNP-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DNP-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140617</p> <p>DNP-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.40%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p><b>DNP-PEG3-DNP</b></p> <p style="text-align: right;">Cat. No.: HY-130350</p> <p>DNP-PEG3-DNP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DNP-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130198</p> <p>DNP-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DNP-PEG4-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140618</p> <p>DNP-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DNP-PEG4-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140619</p> <p>DNP-PEG4-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DNP-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140614</p> <p>DNP-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DNP-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140495</p> <p>DNP-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DNP-PEG6-Boc</b> (DNP-PEG6-t-butyl ester)</p> <p style="text-align: right;">Cat. No.: HY-140616</p> <p>DNP-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>













<p><b>Docosanedioic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W034918</p>	<p><b>Dodecaethylene glycol</b> (HO-PEG12-OH)</p> <p style="text-align: right;">Cat. No.: HY-141233</p>
<p>Docosanedioic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Docosanedioic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 250 mg</p>	<p>Dodecaethylene glycol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 98.36% <b>Clinical Data:</b> <b>Size:</b> 100 mg</p>
<p><b>Dodecylheptaglycol</b></p> <p style="text-align: right;">Cat. No.: HY-W140896</p>	<p><b>DOTA-(t-butyl)3-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140753</p>
<p>Dodecylheptaglycol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>DOTA-(t-butyl)3-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DOTA-PEG5-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140008</p>	<p><b>DOTA-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140752</p>
<p>DOTA-PEG5-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> <b>Size:</b> 5 mg, 10 mg, 50 mg, 100 mg</p>	<p>DOTA-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> <b>Size:</b> 10 mg</p>
<p><b>DOTA-PEG5-C4-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140314</p>	<p><b>Dox-Ph-PEG1-Cl</b> (PROTAC Linker 34)</p> <p style="text-align: right;">Cat. No.: HY-125907</p>
<p>DOTA-PEG5-C4-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 50 mg</p>	<p>Dox-Ph-PEG1-Cl (PROTAC Linker 34) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DSPE-PEG-2-Aminoethyl-alpha-mannopyranoside (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140960</p>	<p><b>DSPE-PEG-Amine (MW 3400)</b></p> <p style="text-align: right;">Cat. No.: HY-140734</p>
<p>DSPE-PEG-2-Aminoethyl-alpha-mannopyranoside (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>DSPE-PEG-Amine (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>DSPE-PEG-Amine (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140735</p>	<p><b>DSPE-PEG-Biotin (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140736</p>
<p>DSPE-PEG-Amine (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>DSPE-PEG-Biotin (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DSPE-PEG-CH2COOH (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140737</p>	<p><b>DSPE-PEG-CH2COOH (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140738</p>
<p>DSPE-PEG-CH2COOH (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>DSPE-PEG-CH2COOH (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DSPE-PEG-Maleimide (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140740</p>	<p><b>DSPE-PEG-OH (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140741</p>
<p>DSPE-PEG-Maleimide (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>DSPE-PEG-OH (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DSPE-PEG13-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140959</p>	<p><b>DSPE-PEG14-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-134700</p>
<p>DSPE-PEG13-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>DSPE-PEG14-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>
<p><b>DSPE-PEG2-mal</b></p> <p style="text-align: right;">Cat. No.: HY-134745</p>	<p><b>DSPE-PEG36-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-133381</p>
<p>DSPE-PEG2-mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>DSPE-PEG36-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>DSPE-PEG36-mal</b></p> <p style="text-align: right;">Cat. No.: HY-133380</p> <p>DSPE-PEG36-mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DSPE-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140953</p> <p>DSPE-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DSPE-PEG4-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140955</p> <p>DSPE-PEG4-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DSPE-PEG46-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-133382</p> <p>DSPE-PEG46-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DSPE-PEG46-Folate</b></p> <p style="text-align: right;">Cat. No.: HY-133383</p> <p>DSPE-PEG46-Folate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DSPE-PEG46-N3</b></p> <p style="text-align: right;">Cat. No.: HY-130906</p> <p>DSPE-PEG46-N3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DSPE-PEG46-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-130905</p> <p>DSPE-PEG46-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DSPE-PEG47-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130904</p> <p>DSPE-PEG47-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>DSPE-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140954</p> <p>DSPE-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DSPE-PEG5-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140958</p> <p>DSPE-PEG5-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

<p><b>DSPE-PEG6-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-138395</p> <p>DSPE-PEG6-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>DSPE-PEG8-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-140956</p> <p>DSPE-PEG8-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>EGNHS</b> (EGS crosslinker)</p> <p style="text-align: right;">Cat. No.: HY-130458</p> <p>EGNHS is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>Eicosanedioic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W034595</p> <p>Eicosanedioic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Eicosanedioic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Eicosanedioic acid-d4</b></p> <p style="text-align: right;">Cat. No.: HY-W034595S</p> <p>Eicosanedioic acid-d4 is the deuterium labeled Eicosanedioic acid. Eicosanedioic acid is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>endo-BCN-O-PNB</b></p> <p style="text-align: right;">Cat. No.: HY-43581</p> <p>endo-BCN-O-PNB is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>endo-BCN-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140066</p> <p>endo-BCN-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>endo-BCN-PEG12-NH2 hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-138386</p> <p>endo-BCN-PEG12-NH2 hydrochloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>endo-BCN-PEG12-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140071</p> <p>endo-BCN-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>endo-BCN-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130557</p> <p>endo-BCN-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>endo-BCN-PEG2-alcohol</b></p> <p>Cat. No.: HY-140076</p>	<p><b>endo-BCN-PEG2-C2-NHS ester</b></p> <p>Cat. No.: HY-140067</p>
<p>endo-BCN-PEG2-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>endo-BCN-PEG2-C2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>endo-BCN-PEG2-NH2</b></p> <p>Cat. No.: HY-W072752</p>	<p><b>endo-BCN-PEG2-PFP ester</b></p> <p>Cat. No.: HY-140072</p>
<p>endo-BCN-PEG2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>Relative stereochemistry</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>endo-BCN-PEG2-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>endo-BCN-PEG3-acid</b></p> <p>Cat. No.: HY-133008</p>	<p><b>endo-BCN-PEG3-mal</b></p> <p>Cat. No.: HY-133400</p>
<p>endo-BCN-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>endo-BCN-PEG3-mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>endo-BCN-PEG3-NH-Boc</b></p> <p>Cat. No.: HY-140077</p>	<p><b>endo-BCN-PEG3-NH2</b></p> <p>Cat. No.: HY-133401</p>
<p>endo-BCN-PEG3-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>endo-BCN-PEG3-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>endo-BCN-PEG3-NHS ester</b></p> <p>Cat. No.: HY-140068</p>	<p><b>endo-BCN-PEG4-acid</b></p> <p>Cat. No.: HY-133009</p>
<p>endo-BCN-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg</p>	<p>endo-BCN-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>


<p><b>endo-BCN-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140074</p> <p>endo-BCN-PEG4-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>endo-BCN-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140069</p> <p>endo-BCN-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>endo-BCN-PEG4-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140073</p> <p>endo-BCN-PEG4-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>endo-BCN-PEG6-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140075</p> <p>endo-BCN-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>endo-BCN-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140065</p> <p>endo-BCN-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>endo-BCN-PEG8-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140070</p> <p>endo-BCN-PEG8-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ethyl acetate-PEG1</b></p> <p style="text-align: right;">Cat. No.: HY-W096086</p> <p>Ethyl acetate-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Ethyl azetidine-3-carboxylate hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-W052600</p> <p>Ethyl azetidine-3-carboxylate hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Ethyl azetidine-3-carboxylate hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>4/5</sup>.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>
<p><b>Ethyl-PEG4-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W096083</p> <p>Ethyl-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>exo BCN-O-PNB</b></p> <p style="text-align: right;">Cat. No.: HY-133406</p> <p>exo BCN-O-PNB is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>F-PEG2-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138498</p>	<p><b>F-PEG2-S-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138493</p>
<p>F-PEG2-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>F-PEG2-S-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>F-PEG2-S-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138497</p>	<p><b>F-PEG2-SO-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138496</p>
<p>F-PEG2-S-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>F-PEG2-SO-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>F-PEG2-SO2-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138495</p>	<p><b>FA-PEG5-Mal</b> (Folic acid-PEG5-Mal)</p> <p style="text-align: right;">Cat. No.: HY-132070</p>
<p>F-PEG2-SO2-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>FA-PEG5-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fluorescein-PEG3-amine</b></p> <p style="text-align: right;">Cat. No.: HY-141080</p>	<p><b>Fluorescein-PEG3-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141081</p>
<p>Fluorescein-PEG3-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fluorescein-PEG3-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fluorescein-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-120368</p>	<p><b>Fluorescein-PEG5-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133006</p>
<p>Fluorescein-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fluorescein-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

### Fluorescein-PEG5-NHS ester

Cat. No.: HY-141082

Fluorescein-PEG5-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### Fluorescein-PEG6-bis-NHS ester

Cat. No.: HY-141084

Fluorescein-PEG6-bis-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### Fluorescein-PEG6-NHS ester

Cat. No.: HY-141083

Fluorescein-PEG6-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### Fluorescein-thiourea-PEG2-azide

Cat. No.: HY-130160

Fluorescein-thiourea-PEG2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### Fluorescein-thiourea-PEG4-azide

Cat. No.: HY-120781

Fluorescein-thiourea-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### Fluorescein-thiourea-PEG6-acid

Cat. No.: HY-133061

Fluorescein-thiourea-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### Fmoc-8-amino-3,6-dioxaoctanoic acid

(Fmoc-NH-PEG2-CH2COOH)

Cat. No.: HY-W007713

Fmoc-8-amino-3,6-dioxaoctanoic acid (Fmoc-NH-PEG2-CH2COOH) is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-8-amino-3,6-dioxaoctanoic acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** 99.65%  
**Clinical Data:** No Development Reported  
**Size:** 100 mg, 500 mg

### Fmoc-amino-PEG3-CH2COOH

Cat. No.: HY-W008568

Fmoc-amino-PEG3-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** 98.14%  
**Clinical Data:** No Development Reported  
**Size:** 500 mg

### Fmoc-amino-PEG5-acid

Cat. No.: HY-W040245

Fmoc-amino-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** 99.51%  
**Clinical Data:** No Development Reported  
**Size:** 100 mg

### Fmoc-aminoxy-PEG12-acid

Cat. No.: HY-140441

Fmoc-aminoxy-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.























**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg













<p><b>Fmoc-aminoxy-PEG12-NHS ester</b></p> <p>Cat. No.: HY-140442</p>	<p><b>Fmoc-aminoxy-PEG4-acid</b></p> <p>Cat. No.: HY-140440</p>
<p>Fmoc-aminoxy-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-aminoxy-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-aminoxy-PFP ester</b></p> <p>Cat. No.: HY-133404</p>	<p><b>Fmoc-azetidine-3-carboxylic acid</b></p> <p>Cat. No.: HY-W011277</p>
<p>Fmoc-aminoxy-PFP ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-azetidine-3-carboxylic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-azetidine-3-carboxylic acid is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Fmoc-Hyp(Bom)-OH</b></p> <p>Cat. No.: HY-79125</p>	<p><b>Fmoc-Lys (biotin-PEG12)-OH</b></p> <p>Cat. No.: HY-140945</p>
<p>Fmoc-Hyp(Bom)-OH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-Hyp(Bom)-OH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-Lys (biotin-PEG12)-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-Lys (biotin-PEG4)-OH</b></p> <p>Cat. No.: HY-130477</p>	<p><b>Fmoc-Lys (Pal-Glu-OtBu)-OH</b></p> <p>Cat. No.: HY-W045822</p>
<p>Fmoc-Lys (biotin-PEG4)-OH is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-Lys (Pal-Glu-OtBu)-OH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-Lys (Pal-Glu-OtBu)-OH is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-Lys-OH hydrochloride</b></p> <p>Cat. No.: HY-W010975</p>	<p><b>Fmoc-N-amido-PEG2-alcohol</b></p> <p>Cat. No.: HY-W096094</p>
<p>Fmoc-Lys-OH hydrochloride is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-Lys-OH hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 g, 5 g</p>	<p>Fmoc-N-amido-PEG2-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Fmoc-N-amido-PEG2-azide</b></p> <p style="text-align: right;">Cat. No.: HY-132099</p>	<p><b>Fmoc-N-amido-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-132110</p>
<p>Fmoc-N-amido-PEG2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-N-amido-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-N-amido-PEG36-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138320</p>	<p><b>Fmoc-N-amido-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-134715</p>
<p>Fmoc-N-amido-PEG36-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-N-amido-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-N-amido-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-132111</p>	<p><b>Fmoc-N-amido-PEG6-amine</b></p> <p style="text-align: right;">Cat. No.: HY-138331</p>
<p>Fmoc-N-amido-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-N-amido-PEG6-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-N-methyl-PEG3-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W035378</p>	<p><b>Fmoc-N-PEG-CH2COOH (MW 3400)</b></p> <p style="text-align: right;">Cat. No.: HY-140668</p>
<p>Fmoc-N-methyl-PEG3-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-N-methyl-PEG3-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-N-amido-PEG-CH2COOH (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Fmoc-N-PEG-CH2COOH (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140669</p>	<p><b>Fmoc-N-PEG20-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140462</p>
<p>Fmoc-N-amido-PEG-CH2COOH (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-N-amido-PEG20-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Fmoc-N-PEG23-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140463</p>	<p><b>Fmoc-N-PEG24-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130907</p>
<p>Fmoc-N-amido-PEG23-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-N-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-N-PEG3-CH2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-135029</p>	<p><b>Fmoc-N-PEG36-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140464</p>
<p>Fmoc-N-PEG3-CH2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-N-amido-PEG36-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-N-PEG7-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140460</p>	<p><b>Fmoc-NH-PEG1-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140013</p>
<p>Fmoc-N-amido-PEG7-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p>Fmoc-NH-PEG1-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-NH-PEG1-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W055861</p>	<p><b>Fmoc-NH-PEG10-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140461</p>
<p>Fmoc-NH-PEG1-CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG1-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.91%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>	<p>Fmoc-NH-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-NH-PEG11-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130870</p>	<p><b>Fmoc-NH-PEG11-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140466</p>
<p>Fmoc-NH-PEG11-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p>Fmoc-NH-PEG11-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>













<p><b>Fmoc-NH-PEG3-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W040231</p>	<p><b>Fmoc-NH-PEG30-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133365</p>
<p>Fmoc-NH-PEG3-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG3-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.80%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 50 mg, 100 mg</p>	<p>Fmoc-NH-PEG30-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-NH-PEG4-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-133470</p>	<p><b>Fmoc-NH-PEG4-CH2CH2COOH</b> (Fmoc-15-amino-4,7,10,13-tetraoxapentadecanoic acid) Cat. No.: HY-W000434</p>
<p>Fmoc-NH-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-NH-PEG4-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG4-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.92%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Fmoc-NH-PEG4-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130175</p>	<p><b>Fmoc-NH-PEG5-C2-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-134682</p>
<p>Fmoc-NH-PEG4-CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG4-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Fmoc-NH-PEG5-C2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-NH-PEG5-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133062</p>	<p><b>Fmoc-NH-PEG5-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-134681</p>
<p>Fmoc-NH-PEG5-CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG5-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-NH-PEG5-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-NH-PEG6-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W040246</p>	<p><b>Fmoc-NH-PEG6-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130364</p>
<p>Fmoc-NH-PEG6-CH2CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> 98.86%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 500 mg, 1 g</p>	<p>Fmoc-NH-PEG6-CH2COOH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Fmoc-NH-PEG6-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>























<p><b>Fmoc-PEG2-C2-NHS ester</b></p> <p>Cat. No.: HY-126881</p>	<p><b>Fmoc-PEG24-NHS ester</b></p> <p>Cat. No.: HY-141102</p>
<p>Fmoc-PEG2-CH<sub>2</sub>CH<sub>2</sub>-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-PEG24-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-PEG3-alcohol</b></p> <p>Cat. No.: HY-W143484</p>	<p><b>Fmoc-PEG3-C2-NHS ester</b></p> <p>Cat. No.: HY-120773</p>
<p>Fmoc-PEG3-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-PEG3-CH<sub>2</sub>CH<sub>2</sub>-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-PEG4-NHS ester</b></p> <p>Cat. No.: HY-122456</p>	<p><b>Fmoc-PEG5-NHBoc</b></p> <p>Cat. No.: HY-138330</p>
<p>Fmoc-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Fmoc-PEG5-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-PEG5-NHS ester</b></p> <p>Cat. No.: HY-122459</p>	<p><b>Fmoc-PEG6-NHS ester</b></p> <p>Cat. No.: HY-126882</p>
<p>Fmoc-PEG5-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Fmoc-PEG6-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Fmoc-PEG9-NHS ester</b></p> <p>Cat. No.: HY-138720</p>	<p><b>FmocNH-PEG3-CH2CH2NH2 hydrochloride</b></p> <p>Cat. No.: HY-W190961</p>
<p>Fmoc-PEG9-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>FmocNH-PEG3-CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub> (hydrochloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>FmocNH-PEG4-t-butyl acetate</b></p> <p>Cat. No.: HY-132079</p>	<p><b>FmocNH-PEG4-t-butyl ester</b></p> <p>Cat. No.: HY-132064</p>
<p>FmocNH-PEG4-t-butyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>FmocNH-PEG4-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Folate-PEG1-mal</b></p> <p>Cat. No.: HY-133494</p>	<p><b>Folate-PEG2-amine</b></p> <p>Cat. No.: HY-133495</p>
<p>Folate-PEG1-mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Folate-PEG2-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Folate-PEG3-alkyne</b></p> <p>Cat. No.: HY-133496</p>	<p><b>Folate-PEG3-azide</b></p> <p>Cat. No.: HY-133483</p>
<p>Folate-PEG3-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Folate-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mg, 25 mg, 50 mg</p>
<p><b>Folate-PEG3-NHS ester</b></p> <p>Cat. No.: HY-133493</p>	<p><b>Gly-PEG3-endo-BCN</b></p> <p>Cat. No.: HY-140080</p>
<p>Folate-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Gly-PEG3-endo-BCN is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Glycocholic acid-PEG10-iodoacetamide</b></p> <p>Cat. No.: HY-134718</p>	<p><b>H-cis-Hyp-OMe hydrochloride</b></p> <p>Cat. No.: HY-W016429</p>
<p>Glycocholic acid-PEG10-iodoacetamide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>H-cis-Hyp-OMe hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). H-cis-Hyp-OMe hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PR.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 g, 5 g</p> <p>HCl</p>













<p><b>H-Glu-OtBu</b></p> <p style="text-align: right;">Cat. No.: HY-W018154</p> <p>H-Glu-OtBu is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). H-Glu-OtBu is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>2</sup>.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 g</p>	<p><b>H-Hyp-OMe hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-76043</p> <p>H-Hyp-OMe hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). H-Hyp-OMe hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>2</sup>.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 g, 5 g</p>
<p><b>H2N-PEG12-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133343</p> <p>H2N-PEG12-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>H2N-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W006524</p> <p>H2N-PEG2-CH2COOH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>H2N-PEG2-CH2COOtBu</b></p> <p style="text-align: right;">Cat. No.: HY-135927</p> <p>H2N-PEG2-CH2COOtBu is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p><b>H2N-PEG4-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133339</p> <p>H2N-PEG4-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>H2N-PEG5-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133340</p> <p>H2N-PEG5-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>H2N-PEG6-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133341</p> <p>H2N-PEG6-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>H2N-PEG8-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133342</p> <p>H2N-PEG8-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Heptaethylene glycol</b> (HO-PEG7-OH)</p> <p style="text-align: right;">Cat. No.: HY-141231</p> <p>Heptaethylene glycol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>

<p><b>Hexaethylene glycol</b></p> <p>Cat. No.: HY-141230</p>	<p><b>Hexaethylene glycol dimethyl ether</b></p> <p>Cat. No.: HY-134746</p>
<p>Hexaethylene glycol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.62% <b>Clinical Data:</b> <b>Size:</b> 100 mg</p>	<p>Hexaethylene glycol dimethyl ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hexaethylene glycol monomethyl ether</b></p> <p>Cat. No.: HY-W042625</p>	<p><b>Hexane-1,6-diylidiphosphonic acid</b></p> <p>Cat. No.: HY-140326</p>
<p>Hexaethylene glycol monomethyl ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Hexane-1,6-diylidiphosphonic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG-amine (MW 1000)</b></p> <p>Cat. No.: HY-140713</p>	<p><b>HO-PEG-amine (MW 10000)</b></p> <p>Cat. No.: HY-140641</p>
<p>HO-PEG-amine (MW 1000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 1000</p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG-amine (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 10000</p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG-amine (MW 2000)</b></p> <p>Cat. No.: HY-140714</p>	<p><b>HO-PEG-amine (MW 3400)</b></p> <p>Cat. No.: HY-140715</p>
<p>HO-PEG-amine (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 2000</p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG-amine (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 3400</p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG-amine (MW 5000)</b></p> <p>Cat. No.: HY-140716</p>	<p><b>HO-PEG-CH2COOH (MW 3400)</b></p> <p>Cat. No.: HY-140711</p>
<p>HO-PEG-amine (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG-CH2COOH (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 3400</p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>









<p><b>HO-PEG-CH<sub>2</sub>COOH (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140712</p>	<p><b>HO-PEG-mal (MW 3400)</b></p> <p style="text-align: right;">Cat. No.: HY-140717</p>
<p>HO-PEG-CH<sub>2</sub>COOH (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 5000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG-mal (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 3400</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG-mal (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140718</p>	<p><b>HO-PEG1-benzyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-133315</p>
<p>HO-PEG-mal (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 5000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG1-benzyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG10-CH<sub>2</sub>COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133304</p>	<p><b>HO-PEG11-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130342</p>
<p>HO-PEG10-CH<sub>2</sub>COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG11-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>
<p><b>HO-PEG12-CH<sub>2</sub>COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133303</p>	<p><b>HO-PEG13-OH</b></p> <p style="text-align: right;">Cat. No.: HY-141234</p>
<p>HO-PEG12-CH<sub>2</sub>COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG13-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG14-OH</b></p> <p style="text-align: right;">Cat. No.: HY-120430</p>	<p><b>HO-PEG15-OH</b></p> <p style="text-align: right;">Cat. No.: HY-120724</p>
<p>HO-PEG14-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG15-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>HO-PEG16-OH</b></p> <p style="text-align: right;">Cat. No.: HY-141235</p> <p>HO-PEG16-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>HO-PEG17-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130600</p> <p>HO-PEG17-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG18-OH</b></p> <p style="text-align: right;">Cat. No.: HY-144081</p> <p>HO-PEG18-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>HO-PEG20-OH</b></p> <p style="text-align: right;">Cat. No.: HY-141236</p> <p>HO-PEG20-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG21-OH</b></p> <p style="text-align: right;">Cat. No.: HY-141237</p> <p>HO-PEG21-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>HO-PEG22-OH</b></p> <p style="text-align: right;">Cat. No.: HY-133300</p> <p>HO-PEG22-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG24-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133309</p> <p>HO-PEG24-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mg, 25 mg, 50 mg</p>	<p><b>HO-PEG24-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133305</p> <p>HO-PEG24-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG24-OH</b></p> <p style="text-align: right;">Cat. No.: HY-141238</p> <p>HO-PEG24-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p><b>HO-PEG4-benzyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-133316</p> <p>HO-PEG4-benzyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>HO-PEG5-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-134750</p>	<p><b>HO-PEG6-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133302</p>
<p>HO-PEG5-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG6-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG7-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-134748</p>	<p><b>HO-PEG8-CH2CH2COOH</b> (Hydroxy-PEG8-acid)</p> <p style="text-align: right;">Cat. No.: HY-133053</p>
<p>HO-PEG7-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>HO-PEG8-CH2CH2COOH (Hydroxy-PEG8-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HO-PEG8-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133301</p>	<p><b>HOOCH2O-PEG4-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-124780</p>
<p>HO-PEG8-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>HOOCH2O-PEG4-CH2COOH, compound 5, is a symmetric PEG linker, used for the synthesis of the first class of Homo-PROTAC.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b>  <b>Size:</b> 10 mg, 50 mg, 100 mg</p>
<p><b>HS-C6-PEG9-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133387</p>	<p><b>HS-PEG10-CH2CH2COOH</b> (Thiol-PEG10-propionic acid)</p> <p style="text-align: right;">Cat. No.: HY-130876</p>
<p>HS-C6-PEG9-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>HS-PEG10-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HS-PEG11-CH2CH2N3</b></p> <p style="text-align: right;">Cat. No.: HY-130880</p>	<p><b>HS-PEG24-CH2CH2COOH</b> (Thiol-PEG24-acid)</p> <p style="text-align: right;">Cat. No.: HY-130878</p>
<p>HS-PEG11-CH2CH2N3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>HS-PEG24-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Hydroxy-Amino-bis(PEG1-C2-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-141250</p>	<p><b>Hydroxy-Amino-bis(PEG2-propargyl)</b></p> <p style="text-align: right;">Cat. No.: HY-140082</p>
<p>Hydroxy-Amino-bis(PEG1-C2-Boc) is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-Amino-bis(PEG2-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG1-(CH2)2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-W055870</p>	<p><b>Hydroxy-PEG1-C2-methyl ester</b> (Methyl 3-(2-hydroxyethoxy)propanoate)</p> <p style="text-align: right;">Cat. No.: HY-141239</p>
<p>Hydroxy-PEG1-(CH2)2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p>Hydroxy-PEG1-C2-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG1-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141208</p>	<p><b>Hydroxy-PEG10-acid</b> (HO-PEG10-CH2CH2COOH)</p> <p style="text-align: right;">Cat. No.: HY-133307</p>
<p>Hydroxy-PEG1-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Hydroxy-PEG10-CH2-Boc</b> (HO-PEG10-CH2COOtBu)</p> <p style="text-align: right;">Cat. No.: HY-133312</p>	<p><b>Hydroxy-PEG11-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141206</p>
<p>Hydroxy-PEG10-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG11-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG12-acid</b> (HO-PEG12-CH2CH2COOH)</p> <p style="text-align: right;">Cat. No.: HY-133306</p>	<p><b>Hydroxy-PEG12-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130831</p>
<p>Hydroxy-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG12-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Hydroxy-PEG12-CH2-Boc</b> (HO-PEG12-CH2COOtBu)</p> <p>Cat. No.: HY-133313</p> <p>Hydroxy-PEG12-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Hydroxy-PEG13-Boc</b></p> <p>Cat. No.: HY-130832</p> <p>Hydroxy-PEG13-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG14-t-butyl ester</b></p> <p>Cat. No.: HY-132056</p> <p>Hydroxy-PEG14-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Hydroxy-PEG16-acid</b> (HO-PEG16-CH2CH2COOH)</p> <p>Cat. No.: HY-133308</p> <p>Hydroxy-PEG16-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG16-Boc</b></p> <p>Cat. No.: HY-130833</p> <p>Hydroxy-PEG16-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Hydroxy-PEG2-(CH2)2-Boc</b></p> <p>Cat. No.: HY-W067061</p> <p>Hydroxy-PEG2-(CH2)2-Boc is a uncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Hydroxy-PEG2-(CH2)2-Boc is extracted from patent WO2004008101A2 (compound 196).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 50 mg, 100 mg</p>
<p><b>Hydroxy-PEG2-acid</b></p> <p>Cat. No.: HY-124380</p> <p>Hydroxy-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 250 mg, 500 mg</p>	<p><b>Hydroxy-PEG2-C2-methyl ester</b> (Methyl 3-[2-(2-hydroxyethoxy)ethoxy]propanoate)</p> <p>Cat. No.: HY-141240</p> <p>Hydroxy-PEG2-C2-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 25 mg, 100 mg</p>
<p><b>Hydroxy-PEG2-C2-PFP ester</b></p> <p>Cat. No.: HY-130488</p> <p>Hydroxy-PEG2-C2-PFP ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Hydroxy-PEG2-C2-sulfonic acid</b></p> <p>Cat. No.: HY-121507</p> <p>Hydroxy-PEG2-C2-sulfonic acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>













<p><b>Hydroxy-PEG2-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130599</p>	<p><b>Hydroxy-PEG2-CH2CH2COO-PEG2-propionic acid</b></p> <p style="text-align: right;">Cat. No.: HY-132083</p>
<p>Hydroxy-PEG2-CH2-Boc is a PEG/Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG2-CH2CH2COO-PEG2-propionic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140497</p>	<p><b>Hydroxy-PEG2-CH2COONa</b> (HO-PEG2-CH2COONa)</p> <p style="text-align: right;">Cat. No.: HY-140497A</p>
<p>Hydroxy-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG2-CH2COONa is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG20-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141207</p>	<p><b>Hydroxy-PEG24-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130834</p>
<p>Hydroxy-PEG20-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG24-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG24-CH2-Boc</b> (HO-PEG24-CH2COOtBu)</p> <p style="text-align: right;">Cat. No.: HY-133314</p>	<p><b>Hydroxy-PEG3-(CH2)2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-42488</p>
<p>Hydroxy-PEG24-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG2-(CH2)2-Boc is a uncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Hydroxy-PEG2-(CH2)2-Boc is extracted from patent WO2004008101A2 (compound 196).</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>Hydroxy-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133052</p>	<p><b>Hydroxy-PEG3-acrylate</b></p> <p style="text-align: right;">Cat. No.: HY-141244</p>
<p>Hydroxy-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG3-acrylate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Hydroxy-PEG3-C2-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141241</p> <p>Hydroxy-PEG3-C2-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Hydroxy-PEG3-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141209</p> <p>Hydroxy-PEG3-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG3-ethyl acetate</b></p> <p style="text-align: right;">Cat. No.: HY-W096154</p> <p>Hydroxy-PEG3-ethyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Hydroxy-PEG3-MS</b></p> <p style="text-align: right;">Cat. No.: HY-W096069</p> <p>Hydroxy-PEG3-MS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG3-NHS</b></p> <p style="text-align: right;">Cat. No.: HY-130672</p> <p>Hydroxy-PEG3-NHS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Hydroxy-PEG3-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-141212</p> <p>Hydroxy-PEG3-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG4-(CH2)2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-W039178</p> <p>Hydroxy-PEG4-(CH2)2-Boc is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Hydroxy-PEG4-(CH2)2-Boc is extracted from patent WO2004008101A2 (compound 191).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 500 mg</p>	<p><b>Hydroxy-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-117104</p> <p>Hydroxy-PEG4-acid is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Hydroxy-PEG4-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG4-C2-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141242</p> <p>Hydroxy-PEG4-C2-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Hydroxy-PEG4-C2-nitrile</b></p> <p style="text-align: right;">Cat. No.: HY-141245</p> <p>Hydroxy-PEG4-C2-nitrile is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Hydroxy-PEG4-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-42617</p>	<p><b>Hydroxy-PEG4-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140498</p>
<p>Hydroxy-PEG4-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Hydroxy-PEG4-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG4-methyl acetate</b></p> <p style="text-align: right;">Cat. No.: HY-138421</p>	<p><b>Hydroxy-PEG4-methylamine</b></p> <p style="text-align: right;">Cat. No.: HY-140162</p>
<p>Hydroxy-PEG4-methyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG4-methylamine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG4-O-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141210</p>	<p><b>Hydroxy-PEG5-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133051</p>
<p>Hydroxy-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG5-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141202</p>	<p><b>Hydroxy-PEG5-C2-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141243</p>
<p>Hydroxy-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG5-C2-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133050</p>	<p><b>Hydroxy-PEG6-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141203</p>
<p>Hydroxy-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG7-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Hydroxy-PEG6-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141211</p>	<p><b>Hydroxy-PEG7-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141204</p>
<p>Hydroxy-PEG6-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG7-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Hydroxy-PEG7-CH2-Boc</b> (HO-PEG7-CH2COOtBu)</p> <p style="text-align: right;">Cat. No.: HY-133310</p>	<p><b>Hydroxy-PEG8-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130662</p>
<p>Hydroxy-PEG7-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 500 mg, 1 g</p>
<p><b>Hydroxy-PEG8-CH2-Boc</b> (HO-PEG8-CH2COOtBu)</p> <p style="text-align: right;">Cat. No.: HY-133311</p>	<p><b>Hydroxy-PEG9-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141205</p>
<p>Hydroxy-PEG8-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Hydroxy-PEG9-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>HyNic-PEG2-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-133498</p>	<p><b>HyNic-PEG2-TCO</b></p> <p style="text-align: right;">Cat. No.: HY-133497</p>
<p>HyNic-PEG2-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>HyNic-PEG2-TCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>I-PEG5-OH</b></p> <p style="text-align: right;">Cat. No.: HY-W096157</p>	<p><b>I-PEG6-OH</b></p> <p style="text-align: right;">Cat. No.: HY-W096156</p>
<p>I-PEG5-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>I-PEG6-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>







<p><b>ICG-Sulfo-OSu sodium</b></p> <p>Cat. No.: HY-134719</p>	<p><b>Iodo-PEG12-acid</b></p> <p>Cat. No.: HY-133481</p>
<p>ICG-Sulfo-OSu sodium is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Iodo-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Iodo-PEG12-NHS ester</b></p> <p>Cat. No.: HY-133482</p>	<p><b>Iodo-PEG4-N3</b></p> <p>Cat. No.: HY-130525</p>
<p>Iodo-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Iodo-PEG4-N3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Iodo-PEG7-alcohol</b></p> <p>Cat. No.: HY-143833</p>	<p><b>Iodoacetamide-PEG3-azide</b></p> <p>Cat. No.: HY-140857</p>
<p>Iodo-PEG7-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Iodoacetamide-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.93%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Iodoacetamide-PEG5-azide</b></p> <p>Cat. No.: HY-134687</p>	<p><b>Iodoacetamide-PEG5-NH-Boc</b></p> <p>Cat. No.: HY-134738</p>
<p>Iodoacetamide-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Iodoacetamide-PEG5-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Iodoacetamide-PEG5-NH2</b></p> <p>Cat. No.: HY-134688</p>	<p><b>Iodoacetamido-PEG6-acid</b></p> <p>Cat. No.: HY-138505</p>
<p>Iodoacetamide-PEG5-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Iodoacetamido-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Iodoacetyl-PEG4-biotin</b></p> <p style="text-align: right;">Cat. No.: HY-138491</p>	<p><b>Iodoacetyl-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-138422</p>
<p>Iodoacetyl-PEG4-biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 50 mg, 100 mg</p>	<p>Iodoacetyl-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Iodoacetyl-PEG8-biotin</b></p> <p style="text-align: right;">Cat. No.: HY-138490</p>	<p><b>L-Azidohomoalanine</b></p> <p style="text-align: right;">Cat. No.: HY-140346</p>
<p>Iodoacetyl-PEG8-biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>L-Azidohomoalanine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 50 mg, 100 mg</p>
<p><b>L-Azidohomoalanine hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-140346A</p>	<p><b>L-Azidohomoalanine-1,2,3,4-13C4 hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-140346AS</p>
<p>L-Azidohomoalanine hydrochloride is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.76%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 10 mM × 1 mL, 50 mg</p>	<p>L-Azidohomoalanine-1,2,3,4-13C4 hydrochloride is the 13C- and 15N-labeled L-Azidohomoalanine hydrochloride. L-Azidohomoalanine hydrochloride is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>L-Homopropargylglycine</b></p> <p style="text-align: right;">Cat. No.: HY-140345</p>	<p><b>L-Homopropargylglycine hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-140345A</p>
<p>L-Homopropargylglycine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs. L-homopropargylglycine is an amino acid analog of methionine containing an alkyne moiety that can undergo a classic click chemical reaction with azide containing Alexa Fluor.</p>  <p><b>Purity:</b> ≥98.0%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 5 mg, 10 mg, 25 mg</p>	<p>L-Homopropargylglycine hydrochloride is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 5 mg, 10 mg, 25 mg</p>
<p><b>Leukotriene B4</b> (LTB4; 5(S),12(R)-DIHETE)</p> <p style="text-align: right;">Cat. No.: HY-107608</p>	<p><b>Leukotriene B4-d4</b> (LTB4-d4; 5(S),12(R)-DIHETE-d4)</p> <p style="text-align: right;">Cat. No.: HY-107608S</p>
<p>Leukotriene B4 (LTB4) is known as one of the most potent chemoattractants and activators of leukocytes and is involved in inflammatory diseases. Leukotriene B4 is also an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%</p> <p><b>Clinical Data:</b> Phase 2</p> <p><b>Size:</b> 25 µg (297.2 µM * 250 µL in Ethanol)</p>	<p>Leukotriene B4-d4 (LTB4-d4) is the deuterium labeled Leukotriene B4. Leukotriene B4 (LTB4) is known as one of the most potent chemoattractants and activators of leukocytes and is involved in inflammatory diseases.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 25 µg</p>









<p><b>LG-PEG10-click-DBCO-Oleic</b></p> <p style="text-align: right;">Cat. No.: HY-141130</p>	<p><b>Lipoamide-PEG11-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-141339</p>
<p>LG-PEG10-click-DBCO-Oleic is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Lipoamide-PEG11-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Lipoamide-PEG3-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-141338</p>	<p><b>Lipoamido-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141334</p>
<p>Lipoamide-PEG3-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Lipoamido-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Lipoamido-PEG2-OH</b></p> <p style="text-align: right;">Cat. No.: HY-141337</p>	<p><b>Lipoamido-PEG24-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141335</p>
<p>Lipoamido-PEG2-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg</p>	<p>Lipoamido-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Lipoamido-PEG3-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130407</p>	<p><b>Lipoamido-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141332</p>
<p>Lipoamido-PEG3-OH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Lipoamido-PEG3-OH (compound TA-TEG-G2CN) can be used in the formation of a highly stable, dendronized gold nanoparticle (AuNP)-based drug delivery platform.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Lipoamido-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Lipoamido-PEG4-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130197</p>	<p><b>Lipoamido-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141333</p>
<p>Lipoamido-PEG4-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Lipoamido-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130292</p>	<p><b>m-PEG-acrylate (MW 10000)</b></p> <p style="text-align: right;">Cat. No.: HY-140671</p>
<p>m-PEG-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-acrylate (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p style="text-align: center;">MW 10000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-acrylate (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140670</p>	<p><b>m-PEG-acrylate (MW 20000)</b></p> <p style="text-align: right;">Cat. No.: HY-140672</p>
<p>m-PEG-acrylate (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p style="text-align: center;">MW 2000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-acrylate (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p style="text-align: center;">MW 20000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-acrylate (MW 30000)</b></p> <p style="text-align: right;">Cat. No.: HY-140673</p>	<p><b>m-PEG-Aminoxy (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140695</p>
<p>m-PEG-acrylate (MW 30000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p style="text-align: center;">MW 30000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-Aminoxy (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p style="text-align: center;">MW 2000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-azide (MW 10000)</b></p> <p style="text-align: right;">Cat. No.: HY-140684</p>	<p><b>m-PEG-azide (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140682</p>
<p>m-PEG-azide (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p style="text-align: center;">MW 10000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-azide (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p style="text-align: center;">MW 2000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-azide (MW 20000)</b></p> <p style="text-align: right;">Cat. No.: HY-140685</p>	<p><b>m-PEG-azide (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140683</p>
<p>m-PEG-azide (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p style="text-align: center;">MW 20000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-azide (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p style="text-align: center;">MW 5000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>























<p><b>m-PEG-Butyraldehyde (MW 5000)</b></p> <p>Cat. No.: HY-140674</p>	<p><b>m-PEG-CH<sub>2</sub>COOH (MW 2000)</b></p> <p>Cat. No.: HY-140686</p>
<p>m-PEG-Butyraldehyde (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-CH<sub>2</sub>COOH (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 2000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-CH<sub>2</sub>COOH (MW 20000)</b></p> <p>Cat. No.: HY-140688</p>	<p><b>m-PEG-CH<sub>2</sub>COOH (MW 5000)</b></p> <p>Cat. No.: HY-140687</p>
<p>m-PEG-CH<sub>2</sub>COOH (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 20000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-CH<sub>2</sub>COOH (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-Lys-NHS ester (MW 20000)</b></p> <p>Cat. No.: HY-140680</p>	<p><b>m-PEG-Lys-NHS ester (MW 40000)</b></p> <p>Cat. No.: HY-140681</p>
<p>m-PEG-Lys-NHS ester (MW 20000) is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 20000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-Lys-NHS ester (MW 40000) is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 40000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-mal (MW 10000)</b></p> <p>Cat. No.: HY-140691</p>	<p><b>m-PEG-mal (MW 2000)</b></p> <p>Cat. No.: HY-140689</p>
<p>m-PEG-mal (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 10000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-mal (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 2000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 500 mg</p>
<p><b>m-PEG-mal (MW 20000)</b></p> <p>Cat. No.: HY-140692</p>	<p><b>m-PEG-mal (MW 30000)</b></p> <p>Cat. No.: HY-140693</p>
<p>m-PEG-mal (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 20000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-mal (MW 30000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACS.</p>  <p>MW 30000</p> <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>


<p><b>m-PEG-mal (MW 5000)</b></p> <p>Cat. No.: HY-140690</p>	<p><b>m-PEG-NH2 (MW 1000)</b></p> <p>Cat. No.: HY-140675</p>
<p>m-PEG-mal (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-NH2 (MW 1000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 1000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-NH2 (MW 10000)</b></p> <p>Cat. No.: HY-140678</p>	<p><b>m-PEG-NH2 (MW 2000)</b></p> <p>Cat. No.: HY-140676</p>
<p>m-PEG-NH2 (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 10000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-NH2 (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 2000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>m-PEG-NH2 (MW 20000)</b></p> <p>Cat. No.: HY-140679</p>	<p><b>m-PEG-NH2 (MW 5000)</b></p> <p>Cat. No.: HY-140677</p>
<p>m-PEG-NH2 (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 20000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-NH2 (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>m-PEG-NHS ester (MW 10000)</b></p> <p>Cat. No.: HY-140699</p>	<p><b>m-PEG-NHS ester (MW 20000)</b></p> <p>Cat. No.: HY-140700</p>
<p>m-PEG-NHS ester (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 10000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-NHS ester (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 20000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-NHS ester (MW 5000)</b></p> <p>Cat. No.: HY-140698</p>	<p><b>m-PEG-NPC (MW 20000)</b></p> <p>Cat. No.: HY-140694</p>
<p>m-PEG-NHS ester (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> 99.58%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>m-PEG-NPC (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 20000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG-OH (MW5000)</b></p> <p>Cat. No.: HY-140697</p>	<p><b>m-PEG-OH (MW 2000)</b></p> <p>Cat. No.: HY-140696</p>
<p>m-PEG-OH (MW5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-OH (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 2000</p> <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>
<p><b>m-PEG-Succinimidyl Succinate (MW 5000)</b></p> <p>Cat. No.: HY-140701</p>	<p><b>m-PEG-thiol (MW 10000)</b></p> <p>Cat. No.: HY-140708</p>
<p>m-PEG-Succinimidyl Succinate (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg</p>	<p>m-PEG-thiol (MW 10000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 10000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-thiol (MW 2000)</b></p> <p>Cat. No.: HY-140706</p>	<p><b>m-PEG-thiol (MW 20000)</b></p> <p>Cat. No.: HY-140709</p>
<p>m-PEG-thiol (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 2000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-thiol (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 20000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-thiol (MW 5000)</b></p> <p>Cat. No.: HY-140707</p>	<p><b>m-PEG-Tos (MW 2000)</b></p> <p>Cat. No.: HY-140705</p>
<p>m-PEG-thiol (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-Tos (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 2000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG-Tresyl (MW 5000)</b></p> <p>Cat. No.: HY-140710</p>	<p><b>m-PEG-triethoxysilane (MW 1000)</b></p> <p>Cat. No.: HY-140702</p>
<p>m-PEG-Tresyl (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-triethoxysilane (MW 1000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 1000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>m-PEG-triethoxysilane (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140703</p>	<p><b>m-PEG-triethoxysilane (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140704</p>
<p>m-PEG-triethoxysilane (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 2000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG-triethoxysilane (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 5000</p> </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG1-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-114512</p>	<p><b>m-PEG10-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140500</p>
<p>m-PEG1-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>m-PEG10-acid is a non-cleavable 10 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG10-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG10-alcohol</b> (Decaethylene glycol monomethyl ether)</p> <p style="text-align: right;">Cat. No.: HY-141218</p>	<p><b>m-PEG10-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140226</p>
<p>m-PEG10-alcohol (Decaethylene glycol monomethyl ether) is a non-cleavable 10 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG10-alcohol is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG10-amine is a non-cleavable 10 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG10-amine is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG10-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130424</p>	<p><b>m-PEG10-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141399</p>
<p>m-PEG10-azide a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG10-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG10-Br</b></p> <p style="text-align: right;">Cat. No.: HY-133278</p>	<p><b>m-PEG10-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133285</p>
<p>m-PEG10-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG10-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg</p>











<p><b>m-PEG10-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141104</p>	<p><b>m-PEG10-SH</b></p> <p style="text-align: right;">Cat. No.: HY-133270</p>
<p>m-PEG10-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG10-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG10-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140360</p>	<p><b>m-PEG1000-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138312</p>
<p>m-PEG10-Tos is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG1000-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG11-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140501</p>	<p><b>m-PEG11-Amine</b></p> <p style="text-align: right;">Cat. No.: HY-W040222</p>
<p>m-PEG11-acid is a non-cleavable 11 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG11-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG11-Amino is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG11-Amine is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG11-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140829</p>	<p><b>m-PEG11-Br</b></p> <p style="text-align: right;">Cat. No.: HY-133279</p>
<p>m-PEG11-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG11-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG11-C2-NHS Ester</b></p> <p style="text-align: right;">Cat. No.: HY-135931</p>	<p><b>m-PEG11-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133347</p>
<p>m-PEG11-C2-NHS Ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG11-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG11-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141105</p>	<p><b>m-PEG11-OH</b></p> <p style="text-align: right;">Cat. No.: HY-141219</p>
<p>m-PEG11-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG11-OH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG11-SH</b></p> <p style="text-align: right;">Cat. No.: HY-133269</p>	<p><b>m-PEG11-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140361</p>
<p>m-PEG11-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG11-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG12-2-methylacrylate</b></p> <p style="text-align: right;">Cat. No.: HY-135824</p>	<p><b>m-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-135820</p>
<p>m-PEG12-2-methylacrylate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG12-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140227</p>	<p><b>m-PEG12-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140830</p>
<p>m-PEG12-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.  m-PEG12-amine is also a non-cleavable 12 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG12-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG12-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141400</p>	<p><b>m-PEG12-Br</b></p> <p style="text-align: right;">Cat. No.: HY-141378</p>
<p>m-PEG12-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG12-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG12-COO-propanoic acid</b> (3-(m-PEG12-ethoxycarbonyl)propanoic acid) <span style="float: right;">Cat. No.: HY-140504</span></p> <p>m-PEG12-COO-propanoic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG12-DBCO</b> <span style="float: right;">Cat. No.: HY-140317</span></p> <p>m-PEG12-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.78% <b>Clinical Data:</b> <b>Size:</b> 50 mg, 100 mg</p>
<p><b>m-PEG12-DSPE</b> <span style="float: right;">Cat. No.: HY-140951</span></p> <p>m-PEG12-DSPE is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG12-Hydrazide</b> <span style="float: right;">Cat. No.: HY-133348</span></p> <p>m-PEG12-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG12-Mal</b> <span style="float: right;">Cat. No.: HY-140983</span></p> <p>m-PEG12-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG12-NH-C2-acid</b> <span style="float: right;">Cat. No.: HY-140508</span></p> <p>m-PEG12-NH-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG12-NHS ester</b> <span style="float: right;">Cat. No.: HY-141106</span></p> <p>m-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> <b>Size:</b> 100 mg, 500 mg</p>	<p><b>m-PEG12-OH</b> <span style="float: right;">Cat. No.: HY-141220</span></p> <p>m-PEG12-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. m-PEG12-OH is also a non-cleavable 12 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG12-Thiol</b> <span style="float: right;">Cat. No.: HY-141331</span></p> <p>m-PEG12-Thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg, 250 mg</p>	<p><b>m-PEG13-azide</b> <span style="float: right;">Cat. No.: HY-138717</span></p> <p>m-PEG13-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>










<p><b>m-PEG13-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141401</p> <p>m-PEG13-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG13-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133349</p> <p>m-PEG13-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG13-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141107</p> <p>m-PEG13-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG14-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-138718</p> <p>m-PEG14-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG15-acetic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W190959</p> <p>m-PEG15-acetic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG15-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141221</p> <p>m-PEG15-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG15-amine</b></p> <p style="text-align: right;">Cat. No.: HY-138424</p> <p>m-PEG15-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p><b>m-PEG15-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138314</p> <p>m-PEG15-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG15-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130829</p> <p>m-PEG15-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG16-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141222</p> <p>m-PEG16-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>






















<p><b>m-PEG16-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140831</p>	<p><b>m-PEG16-Br</b></p> <p style="text-align: right;">Cat. No.: HY-133281</p>
<p>m-PEG16-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG16-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG16-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-133326</p>	<p><b>m-PEG16-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-133288</p>
<p>m-PEG16-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG16-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG16-SH</b></p> <p style="text-align: right;">Cat. No.: HY-133274</p>	<p><b>m-PEG17-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140502</p>
<p>m-PEG16-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG17-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG17-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133350</p>	<p><b>m-PEG17-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141108</p>
<p>m-PEG17-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG17-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG18-acid</b></p> <p style="text-align: right;">Cat. No.: HY-143815</p>	<p><b>m-PEG18-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-138427</p>
<p>m-PEG18-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG18-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>






<p><b>m-PEG19-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141223</p>	<p><b>m-PEG2-4-nitrophenyl carbonate</b></p> <p style="text-align: right;">Cat. No.: HY-117023</p>
<p>m-PEG19-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG2-4-nitrophenyl carbonate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130231</p>	<p><b>m-PEG2-amido-Ph-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-134705</p>
<p>m-PEG2-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG2-amido-Ph-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG2-Amine</b></p> <p style="text-align: right;">Cat. No.: HY-W008429</p>	<p><b>m-PEG2-Amino</b></p> <p style="text-align: right;">Cat. No.: HY-140397</p>
<p>m-PEG2-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG2-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> 99.58%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 50 mg, 100 mg</p>	<p>m-PEG2-Amino is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG2-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130578</p>	<p><b>m-PEG2-Br</b></p> <p style="text-align: right;">Cat. No.: HY-133277</p>
<p>m-PEG2-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG2-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>
<p><b>m-PEG2-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W043725</p>	<p><b>m-PEG2-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-140362</p>
<p>m-PEG2-CH2CH2COOH is PEG-based based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG2-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG2-O-Ph-3-NH2</b></p> <p>Cat. No.: HY-W096118</p>	<p><b>m-PEG2-O-Ph-NH2</b></p> <p>Cat. No.: HY-W096113</p>
<p>m-PEG2-O-Ph-3-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG2-O-Ph-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg, 50 mg, 100 mg</p>
<p><b>m-PEG2-phosphonic acid</b></p> <p>Cat. No.: HY-141306</p>	<p><b>m-PEG2-Tos</b></p> <p>Cat. No.: HY-42745</p>
<p>m-PEG2-phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG2-Tos is a uncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG2-Tos is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>m-PEG20-alcohol</b></p> <p>Cat. No.: HY-W096138</p>	<p><b>m-PEG21-acid</b></p> <p>Cat. No.: HY-133322</p>
<p>m-PEG20-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG21-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG21-OH</b></p> <p>Cat. No.: HY-134743</p>	<p><b>m-PEG23-alcohol</b></p> <p>Cat. No.: HY-141224</p>
<p>m-PEG21-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG23-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG24-acid</b></p> <p>Cat. No.: HY-133323</p>	<p><b>m-PEG24-alcohol</b></p> <p>Cat. No.: HY-141225</p>
<p>m-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p>m-PEG24-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>m-PEG24-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140832</p>	<p><b>m-PEG24-Br</b></p> <p style="text-align: right;">Cat. No.: HY-133282</p>
<p>m-PEG24-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG24-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG24-DSPE</b></p> <p style="text-align: right;">Cat. No.: HY-140952</p>	<p><b>m-PEG24-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-140984</p>
<p>m-PEG24-DSPE is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG24-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG24-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140228</p>	<p><b>m-PEG24-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130830</p>
<p>m-PEG24-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG24-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>m-PEG24-SH</b></p> <p style="text-align: right;">Cat. No.: HY-133275</p>	<p><b>m-PEG25-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130938</p>
<p>m-PEG24-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG25-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG25-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133351</p>	<p><b>m-PEG25-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141109</p>
<p>m-PEG25-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG25-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>



<p><b>m-PEG25-Propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-130901</p>	<p><b>m-PEG3-0-benzaldehyde</b></p> <p style="text-align: right;">Cat. No.: HY-140621</p>
<p>m-PEG25-Propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-0-benzaldehyde is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-aldehyde</b></p> <p style="text-align: right;">Cat. No.: HY-124011</p>	<p><b>m-PEG3-amido-C3-triethoxysilane</b></p> <p style="text-align: right;">Cat. No.: HY-141407</p>
<p>m-PEG3-aldehyde is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-amido-C3-triethoxysilane is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-Amine</b></p> <p style="text-align: right;">Cat. No.: HY-W018174</p>	<p><b>m-PEG3-Aminoxy</b></p> <p style="text-align: right;">Cat. No.: HY-140398</p>
<p>m-PEG3-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG3-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> 97.63%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 50 mg, 100 mg</p>	<p>m-PEG3-Aminoxy is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-aminoxy-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141201</p>	<p><b>m-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140827</p>
<p>m-PEG3-aminoxy-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130404</p>	<p><b>m-PEG3-CH2-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-117191</p>
<p>m-PEG3-Boc is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-CH2-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>m-PEG3-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W067509</p>	<p><b>m-PEG3-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130445</p>
<p>m-PEG3-CH2CH2COOH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG3-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-CH2COOH is a PEG-based based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>m-PEG3-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133344</p>	<p><b>m-PEG3-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-135940</p>
<p>m-PEG3-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-133064</p>	<p><b>m-PEG3-OH</b></p> <p style="text-align: right;">Cat. No.: HY-135796</p>
<p>m-PEG3-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-OMs</b></p> <p style="text-align: right;">Cat. No.: HY-140363</p>	<p><b>m-PEG3-phosphonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W096153</p>
<p>m-PEG3-OMs is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-Propanoyl chloride</b></p> <p style="text-align: right;">Cat. No.: HY-140505</p>	<p><b>m-PEG3-S-Acetyl</b></p> <p style="text-align: right;">Cat. No.: HY-141351</p>
<p>m-PEG3-Propanoyl chloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-S-Acetyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG3-S-PEG1-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140595</p>	<p><b>m-PEG3-S-PEG2-OH</b></p> <p style="text-align: right;">Cat. No.: HY-140593</p>
<p>m-PEG3-S-PEG1-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-S-PEG2-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-S-PEG3-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140596</p>	<p><b>m-PEG3-S-PEG4-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140594</p>
<p>m-PEG3-S-PEG3-Boc is a PEG- and Alkyl/ether -based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-S-PEG4-propargyl is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-SH</b></p> <p style="text-align: right;">Cat. No.: HY-133273</p>	<p><b>m-PEG3-succinimidyl carbonate</b></p> <p style="text-align: right;">Cat. No.: HY-141112</p>
<p>m-PEG3-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-succinimidyl carbonate is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-Sulfone-PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-140607</p>	<p><b>m-PEG3-Sulfone-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140605</p>
<p>m-PEG3-Sulfone-PEG3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-Sulfone-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG3-Sulfone-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140606</p>	<p><b>m-PEG3-Sulfone-PEG3-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140609</p>
<p>m-PEG3-Sulfone-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-Sulfone-PEG3-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG3-Sulfone-PEG4-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140608</p>	<p><b>m-PEG3-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-W043277</p>
<p>m-PEG3-Sulfone-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG3-Tos is a PEG-based PROTAC linker can be used in the synthesis of Silymarin (HY-W043277).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG36-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141226</p>	<p><b>m-PEG36-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140229</p>
<p>m-PEG36-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG36-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG36-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140833</p>	<p><b>m-PEG36-Br</b></p> <p style="text-align: right;">Cat. No.: HY-133283</p>
<p>m-PEG36-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG36-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG36-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-140985</p>	<p><b>m-PEG36-OH</b></p> <p style="text-align: right;">Cat. No.: HY-133287</p>
<p>m-PEG36-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG36-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG37-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133324</p>	<p><b>m-PEG37-hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-141403</p>
<p>m-PEG37-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG37-hydrazide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>














<p><b>m-PEG37-NHS ester</b></p> <p>Cat. No.: HY-141110</p>	<p><b>m-PEG37-Propargyl</b></p> <p>Cat. No.: HY-130902</p>
<p>m-PEG37-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG37-Propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-(CH2)6-Phosphonic acid</b></p> <p>Cat. No.: HY-141309</p>	<p><b>m-PEG4-aldehyde</b></p> <p>Cat. No.: HY-140620</p>
<p>m-PEG4-(CH2)6-Phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-aldehyde is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-Amine</b></p> <p>Cat. No.: HY-W040214</p>	<p><b>m-PEG4-amino-Mal</b></p> <p>Cat. No.: HY-130181</p>
<p>m-PEG4-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG4-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-amino-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-azide</b> (13-Azido-2,5,8,11-tetraoxatridecane)</p> <p>Cat. No.: HY-140828</p>	<p><b>m-PEG4-Boc</b></p> <p>Cat. No.: HY-141395</p>
<p>m-PEG4-azide (13-Azido-2,5,8,11-tetraoxatridecane) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-Boc is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG4-Boc is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-Br</b></p> <p>Cat. No.: HY-130161</p>	<p><b>m-PEG4-C6-phosphonic acid ethyl ester</b></p> <p>Cat. No.: HY-141314</p>
<p>m-PEG4-Br is a cleavable ADC linker used in the synthesis of antibody-drug conjugate (ADC) for Trastuzumab (HY-P9907).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-C6-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG4-CH2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140507</p>	<p><b>m-PEG4-CH2-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-133057</p>
<p>m-PEG4-CH2-acid is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-CH2-alcohol is PEG-based based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-CH2-aldehyde</b></p> <p style="text-align: right;">Cat. No.: HY-117041</p>	<p><b>m-PEG4-CH2-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141404</p>
<p>m-PEG4-CH2-aldehyde is a PEG-based based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-CH2-methyl ester is a PEG- and Alkyl/ester-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130478</p>	<p><b>m-PEG4-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-141402</p>
<p>m-PEG4-CH2COOH is a PEG-based based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>m-PEG4-Hydrazide is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-130457</p>	<p><b>m-PEG4-NH-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140315</p>
<p>m-PEG4-Ms is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs. m-PEG4-Ms is a cleavable <b>ADC linker</b> used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-NH-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-124323</p>	<p><b>m-PEG4-O-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141113</p>
<p>m-PEG4-NHS ester is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-O-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG4-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-138382</p>	<p><b>m-PEG4-phosphonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-141307</p>
<p>m-PEG4-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-phosphonic acid ethyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141311</p>	<p><b>m-PEG4-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-113921</p>
<p>m-PEG4-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-SH</b></p> <p style="text-align: right;">Cat. No.: HY-133271</p>	<p><b>m-PEG4-sulfonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-140170</p>
<p>m-PEG4-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG4-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG4-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-114661</p>	<p><b>m-PEG48-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140230</p>
<p>m-PEG4-Tos is a derivative of silybin ethers, extracted from patent CN105037337A (compound III-b). m-PEG4-Tos is a PEG-based PROTAC linker that can be used in the synthesis of Silymarin (HY-W043277).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG48-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG48-Br</b></p> <p style="text-align: right;">Cat. No.: HY-133284</p>	<p><b>m-PEG48-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-140986</p>
<p>m-PEG48-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG48-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>m-PEG48-OH</b></p> <p style="text-align: right;">Cat. No.: HY-133289</p> <p>m-PEG48-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG49-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133325</p> <p>m-PEG49-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG49-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130903</p> <p>m-PEG49-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG5-2-methylacrylate</b></p> <p style="text-align: right;">Cat. No.: HY-141406</p> <p>m-PEG5-2-methylacrylate is a PEG- and Alkyl/ester-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG5-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W040195</p> <p>m-PEG5-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG5-amino-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-138732</p> <p>m-PEG5-amino-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130168</p> <p>m-PEG5-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG5-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141396</p> <p>m-PEG5-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG5-Br</b></p> <p style="text-align: right;">Cat. No.: HY-141375</p> <p>m-PEG5-Br is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>m-PEG5-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W043840</p> <p>m-PEG5-CH2CH2COOH is a PEG-based based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG5-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-120537</p>	<p><b>m-PEG5-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-135921</p>
<p>m-PEG5-CH2COOH is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG5-CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG5-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG5-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-116186</p>	<p><b>m-PEG5-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140225</p>
<p>m-PEG5-Ms is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG5-Ms is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG5-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG5-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-133065</p>	<p><b>m-PEG5-nitrile</b></p> <p style="text-align: right;">Cat. No.: HY-130657</p>
<p>m-PEG5-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>	<p>m-PEG5-nitrile is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG5-phosphonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-141308</p>	<p><b>m-PEG5-phosphonic acid ethyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141312</p>
<p>m-PEG5-phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG5-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG5-Propyne</b></p> <p style="text-align: right;">Cat. No.: HY-138719</p>	<p><b>m-PEG5-SH</b></p> <p style="text-align: right;">Cat. No.: HY-133272</p>
<p>m-PEG5-Propyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG5-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 500 mg</p>











<p><b>m-PEG5-succinimidyl carbonate</b></p> <p style="text-align: right;">Cat. No.: HY-130150</p>	<p><b>m-PEG5-sulfonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-140171</p>
<p>m-PEG5-succinimidyl carbonate is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG5-succinimidyl carbonate is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG5-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG5-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-W042501</p>	<p><b>m-PEG5-triethoxysilane</b></p> <p style="text-align: right;">Cat. No.: HY-141408</p>
<p>m-PEG5-Tos is a derivative of silybin ethers, extracted from patent CN105037337A (compound III-c). m-PEG5-Tos is a PEG-based PROTAC linker can be used in the synthesis of Silymarin (HY-W043277).</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG5-triethoxysilane is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG6-(CH2)6-Phosphonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-141310</p>	<p><b>m-PEG6-2-methylacrylate</b></p> <p style="text-align: right;">Cat. No.: HY-130548</p>
<p>m-PEG6-(CH2)6-Phosphonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG6-2-methylacrylate is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG6-Amine</b></p> <p style="text-align: right;">Cat. No.: HY-130408</p>	<p><b>m-PEG6-amino-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-130141</p>
<p>m-PEG6-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG6-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG6-amino-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG6-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138449</p>	<p><b>m-PEG6-Br</b></p> <p style="text-align: right;">Cat. No.: HY-141376</p>
<p>m-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG6-Br is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>m-PEG6-C6-phosphonic acid ethyl ester</b></p> <p>Cat. No.: HY-141315</p>	<p><b>m-PEG6-CH2CH2CHO</b></p> <p>Cat. No.: HY-W035376</p>
<p>m-PEG6-C6-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG6-CH2CH2CHO is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG6-CH2CH2CHO is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG6-CH2CH2COOH</b></p> <p>Cat. No.: HY-W040239</p>	<p><b>m-PEG6-Hydrazide</b></p> <p>Cat. No.: HY-135922</p>
<p>m-PEG6-CH2CH2COOH is a PEG-based based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG6-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG6-Ms</b></p> <p>Cat. No.: HY-140364</p>	<p><b>m-PEG6-NHS ester</b></p> <p>Cat. No.: HY-133066</p>
<p>m-PEG6-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG6-NHS ester is a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). m-PEG6-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG6-O-CH2COOH</b></p> <p>Cat. No.: HY-140506</p>	<p><b>m-PEG6-thiol</b></p> <p>Cat. No.: HY-141329</p>
<p>m-PEG6-O-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG6-thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG6-Tos</b></p> <p>Cat. No.: HY-130336</p>	<p><b>m-PEG7-4-nitrophenyl carbonate</b></p> <p>Cat. No.: HY-141405</p>
<p>m-PEG6-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG7-4-nitrophenyl carbonate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>




















<p><b>m-PEG7-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-122413</p>	<p><b>m-PEG7-Silane</b></p> <p style="text-align: right;">Cat. No.: HY-138420</p>
<p>m-PEG7-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG7-Silane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG7-thiol</b></p> <p style="text-align: right;">Cat. No.: HY-W052006</p>	<p><b>m-PEG7-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140359</p>
<p>m-PEG7-thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 500 mg</p>	<p>m-PEG7-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG750-Br</b></p> <p style="text-align: right;">Cat. No.: HY-138448</p>	<p><b>m-PEG8-(CH<sub>2</sub>)<sub>12</sub>-phosphonic acid ethyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141316</p>
<p>m-PEG750-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG8-(CH<sub>2</sub>)<sub>12</sub>-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG8-amide-C10-Thiol</b></p> <p style="text-align: right;">Cat. No.: HY-138532</p>	<p><b>m-PEG8-Amine</b></p> <p style="text-align: right;">Cat. No.: HY-W040236</p>
<p>m-PEG8-amide-C10-Thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG8-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg</p>
<p><b>m-PEG8-azide</b></p> <p style="text-align: right;">Cat. No.: HY-130204</p>	<p><b>m-PEG8-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141397</p>
<p>m-PEG8-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>




















<p><b>m-PEG8-succinimidyl carbonate</b></p> <p style="text-align: right;">Cat. No.: HY-141114</p>	<p><b>m-PEG8-thiol</b></p> <p style="text-align: right;">Cat. No.: HY-141330</p>
<p>m-PEG8-succinimidyl carbonate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG8-thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG8-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-135090</p>	<p><b>m-PEG9-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140499</p>
<p>Tos-PEG8-m is a derivative of silybin ethers, extracted from patent CN105037337A (compound III-d). Tos-PEG8-m is a PEG-based PROTAC linker can be used in the synthesis of Silymarin (HY-W043277).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG9-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG9-Amine</b></p> <p style="text-align: right;">Cat. No.: HY-130571</p>	<p><b>m-PEG9-azide</b></p> <p style="text-align: right;">Cat. No.: HY-138716</p>
<p>m-PEG9-Amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. m-PEG9-Amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg</p>	<p>m-PEG9-azide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG9-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141398</p>	<p><b>m-PEG9-Br</b></p> <p style="text-align: right;">Cat. No.: HY-141377</p>
<p>m-PEG9-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG9-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG9-C4-SH</b></p> <p style="text-align: right;">Cat. No.: HY-135936</p>	<p><b>m-PEG9-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-133286</p>
<p>m-PEG9-C4-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG9-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

<p><b>m-PEG9-Hydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-133346</p>	<p><b>m-PEG9-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-138428</p>
<p>m-PEG9-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG9-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG9-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130446</p>	<p><b>m-PEG9-phosphonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-130554</p>
<p>m-PEG9-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG9-phosphonic acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEG9-phosphonic acid ethyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-141313</p>	<p><b>m-PEG9-SH</b></p> <p style="text-align: right;">Cat. No.: HY-133280</p>
<p>m-PEG9-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>m-PEG9-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>m-PEGn-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-W019795</p>	<p><b>Mal-amide-PEG2-oxyamine</b></p> <p style="text-align: right;">Cat. No.: HY-133507</p>
<p>m-PEGn-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amide-PEG2-oxyamine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amide-PEG2-oxyamine-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133503</p>	<p><b>Mal-amido-PEG10-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140973</p>
<p>Mal-amide-PEG2-oxyamineBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amido-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Mal-amido-PEG12-acid</b></p> <p>Cat. No.: HY-140974</p>	<p><b>Mal-amido-PEG12-NHS ester</b></p> <p>Cat. No.: HY-140977</p>
<p>Mal-amido-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amido-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amido-PEG12-TFP ester</b></p> <p>Cat. No.: HY-140980</p>	<p><b>Mal-amido-PEG15-acid</b></p> <p>Cat. No.: HY-143841</p>
<p>Mal-amido-PEG12-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amido-PEG15-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amido-PEG16-acid</b> (Maleimide-NH-PEG16-CH2CH2COOH)</p> <p>Cat. No.: HY-130858</p>	<p><b>Mal-amido-PEG2-C2-acid</b></p> <p>Cat. No.: HY-42151</p>
<p>Mal-amido-PEG16-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amido-PEG2-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.54% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 10 mg, 50 mg</p>
<p><b>Mal-amido-PEG2-C2-amido-Ph-C2-CO-AZD</b></p> <p>Cat. No.: HY-113697</p>	<p><b>Mal-amido-PEG2-TFP ester</b></p> <p>Cat. No.: HY-130232</p>
<p>Mal-amido-PEG2-C2-amido-Ph-C2-CO-AZD is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.97% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 5 mg, 10 mg, 25 mg, 50 mg, 100 mg</p>	<p>Mal-amido-PEG2-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 95.09% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>Mal-amido-PEG24-acid</b></p> <p>Cat. No.: HY-140975</p>	<p><b>Mal-amido-PEG24-TFP ester</b></p> <p>Cat. No.: HY-140981</p>
<p>Mal-amido-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amido-PEG24-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>









<p><b>Mal-amido-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138727</p>	<p><b>Mal-amido-PEG3-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-143842</p>
<p>Mal-amido-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amido-PEG3-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amido-PEG3-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-138729</p>	<p><b>Mal-amido-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130414</p>
<p>Mal-amido-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amido-PEG4-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 97.01%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Mal-Amido-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140967</p>	<p><b>Mal-amido-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-W040134</p>
<p>Mal-Amido-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amido-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amido-PEG4-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-140978</p>	<p><b>Mal-amido-PEG5-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138728</p>
<p>Mal-amido-PEG4-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 99.32%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Mal-amido-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amido-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130399</p>	<p><b>Mal-amido-PEG6-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-W008879</p>
<p>Mal-amido-PEG6-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-amido-PEG6-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Mal-amido-PEG7-acid</b> (Mal-NH-PEG7-COOH; Maleimide-NH-PEG7-CH<sub>2</sub>CH<sub>2</sub>COOH) Cat. No.: HY-130855</p> <p>Mal-amido-PEG7-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Mal-amido-PEG7-NHS ester</b> Cat. No.: HY-138730</p> <p>Mal-amido-PEG7-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amido-PEG8-NHS ester</b> Cat. No.: HY-W040133</p> <p>Mal-amido-PEG8-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.23% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 25 mg, 50 mg</p>	<p><b>Mal-amido-PEG8-TFP ester</b> Cat. No.: HY-140979</p> <p>Mal-amido-PEG8-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amido-PEG9-acid</b> (Maleimide-NH-PEG9-CH<sub>2</sub>CH<sub>2</sub>COOH) Cat. No.: HY-130856</p> <p>Mal-amido-PEG9-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Mal-amido-PEG9-amine</b> Cat. No.: HY-140976</p> <p>Mal-amido-PEG9-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amido-PEG9-NH-Boc</b> Cat. No.: HY-140972</p> <p>Mal-amido-PEG9-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Mal-amido-PEG9-NHS ester</b> Cat. No.: HY-138731</p> <p>Mal-amido-PEG9-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-amino-sulfo</b> Cat. No.: HY-136168</p> <p>Mal-amino-sulfo is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Mal-C2-cyclohexylcarboxyl-hydrazide hydrochloride</b> Cat. No.: HY-133410A</p> <p>Mal-C2-cyclohexylcarboxyl-hydrazide hydrochloride is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.23% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg</p>

<p><b>Mal-C2-cyclohexylcarboxyl-hydrazide TFA</b></p> <p>Cat. No.: HY-133410</p> <p>Mal-C2-cyclohexylcarboxyl-hydrazide (TFA) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Mal-C4-NH-Boc</b></p> <p>Cat. No.: HY-140992</p> <p>Mal-C4-NH-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-C6-amine TFA</b></p> <p>Cat. No.: HY-W018291</p> <p>Mal-C6-amine (TFA) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>	<p><b>Mal-N-bis(PEG4-C2-acid)</b></p> <p>Cat. No.: HY-140530</p> <p>Mal-N-bis(PEG4-C2-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-NH-Boc</b></p> <p>Cat. No.: HY-140991</p> <p>Mal-NH-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Mal-NH-PEG10-CH2CH2COOPFP ester</b></p> <p>Cat. No.: HY-130865</p> <p>Mal-NH-PEG10-CH2CH2COOPFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-NH-PEG12-CH2CH2COOPFP ester</b></p> <p>Cat. No.: HY-130864</p> <p>Mal-NH-PEG12-CH2CH2COOPFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Mal-NH-PEG14-CH2CH2COOPFP ester</b></p> <p>Cat. No.: HY-130866</p> <p>Mal-NH-PEG14-CH2CH2COOPFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-NH-PEG16-CH2CH2COOPFP ester</b></p> <p>Cat. No.: HY-130867</p> <p>Mal-NH-PEG16-CH2CH2COOPFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Mal-NH-PEG2-CH2CH2COOPFP ester</b></p> <p>Cat. No.: HY-130860</p> <p>Mal-NH-PEG2-CH2CH2COOPFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>













<p><b>Mal-NH-PEG24-CH2CH2COOPFP ester</b></p> <p>Cat. No.: HY-130868</p>	<p><b>Mal-NH-PEG4-CH2CH2COOPFP ester</b></p> <p>Cat. No.: HY-130861</p>
<p>Mal-NH-PEG24-CH2CH2COOPFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-NH-PEG4-CH2CH2COOPFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-NH-PEG6-CH2CH2COOPFP ester</b></p> <p>Cat. No.: HY-130862</p>	<p><b>Mal-NH-PEG8-Boc</b></p> <p>Cat. No.: HY-138447</p>
<p>Mal-NH-PEG6-CH2CH2COOPFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-NH-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-NH-PEG8-CH2CH2COOPFP ester</b></p> <p>Cat. No.: HY-130863</p>	<p><b>Mal-NH2 TFA</b></p> <p>Cat. No.: HY-140988</p>
<p>Mal-NH-PEG8-CH2CH2COOPFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-NH2 (TFA) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-NO2-Ph-PEG12-NHS</b></p> <p>Cat. No.: HY-141115</p>	<p><b>Mal-PEG-mal (MW 2000)</b></p> <p>Cat. No.: HY-140719</p>
<p>Mal-NO2-Ph-PEG12-NHS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG-mal (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG-mal (MW 3400)</b></p> <p>Cat. No.: HY-140720</p>	<p><b>Mal-PEG-mal (MW 5000)</b></p> <p>Cat. No.: HY-140721</p>
<p>Mal-PEG-mal (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG-mal (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Mal-PEG-Succinimidyl Valerate (MW 20000)</b></p> <p>Cat. No.: HY-140722</p>	<p><b>Mal-PEG1-acid</b></p> <p>Cat. No.: HY-126960</p>
<p>Mal-PEG-Succinimidyl Valerate (MW 20000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p>MW 20000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG1-acid is a non-cleavable 1 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Mal-PEG1-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.40%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Mal-PEG1-Boc</b></p> <p>Cat. No.: HY-133162</p>	<p><b>Mal-PEG1-bromide</b></p> <p>Cat. No.: HY-140989</p>
<p>Mal-PEG1-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG1-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG1-NHS ester</b></p> <p>Cat. No.: HY-126886</p>	<p><b>Mal-PEG1-PFP ester</b></p> <p>Cat. No.: HY-130568</p>
<p>Mal-PEG1-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Mal-PEG1-NHS ester is PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.41%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 250 mg</p>	<p>Mal-PEG1-PFP ester is an Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG10-acid</b></p> <p>Cat. No.: HY-138723</p>	<p><b>Mal-PEG10-Boc</b></p> <p>Cat. No.: HY-138722</p>
<p>Mal-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG10-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG10-NHS ester</b></p> <p>Cat. No.: HY-138726</p>	<p><b>Mal-PEG11-mal</b></p> <p>Cat. No.: HY-133354</p>
<p>Mal-PEG10-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.01%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Mal-PEG11-mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>













<p><b>Mal-PEG2-C2-Boc</b></p> <p>Cat. No.: HY-140964</p>	<p><b>Mal-PEG2-NH-Boc</b></p> <p>Cat. No.: HY-130193</p>
<p>Mal-PEG2-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG2-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG2-NH2</b></p> <p>Cat. No.: HY-35261</p>	<p><b>Mal-PEG2-NH2 TFA</b></p> <p>Cat. No.: HY-35261A</p>
<p>Mal-PEG2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 100 mg, 250 mg, 500 mg</p>	<p>Mal-PEG2-NH2 (TFA) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.09%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 100 mg</p>
<p><b>Mal-PEG2-NHS</b></p> <p>Cat. No.: HY-138430</p>	<p><b>Mal-PEG2-oxyamine</b></p> <p>Cat. No.: HY-133454</p>
<p>Mal-PEG2-NHS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG2-oxyamine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 100 mg</p>
<p><b>Mal-PEG2-PFP ester</b></p> <p>Cat. No.: HY-130416</p>	<p><b>Mal-PEG2-Val-Cit-amido-PAB-OH</b></p> <p>Cat. No.: HY-130222</p>
<p>Mal-PEG2-PFP ester is a Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG2-Val-Cit-amido-PAB-OH is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Mal-PEG2-Val-Cit-amido-PAB-OH also can be used as a PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG24-acid</b></p> <p>Cat. No.: HY-130857</p>	<p><b>Mal-PEG24-NHS ester</b></p> <p>Cat. No.: HY-135819</p>
<p>Mal-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG24-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>

<p><b>Mal-PEG25-NH2 hydrochloride</b></p> <p>Cat. No.: HY-138392</p>	<p><b>Mal-PEG3-alcohol</b></p> <p>Cat. No.: HY-140969</p>
<p>Mal-PEG25-NH2 hydrochloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>Mal-PEG3-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG3-Boc</b></p> <p>Cat. No.: HY-135169</p>	<p><b>Mal-PEG3-O-Ac</b></p> <p>Cat. No.: HY-134684</p>
<p>Mal-PEG3-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG3-O-Ac is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG3-PFP ester</b></p> <p>Cat. No.: HY-130586</p>	<p><b>Mal-PEG36-NHS ester</b></p> <p>Cat. No.: HY-133389</p>
<p>Mal-PEG3-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p>Mal-PEG36-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG4-acid</b></p> <p>Cat. No.: HY-126961</p>	<p><b>Mal-PEG4-Boc</b></p> <p>Cat. No.: HY-130469</p>
<p>Mal-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Mal-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.93%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Mal-PEG4-C2-NH2 TFA</b></p> <p>Cat. No.: HY-138380</p>	<p><b>Mal-PEG4-Glu(OH)-NH-m-PEG24</b></p> <p>Cat. No.: HY-140997</p>
<p>Mal-PEG4-C2-NH2 TFA is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG4-Glu(OH)-NH-m-PEG24 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>



<p><b>Mal-PEG4-Glu(TFP ester)-NH-m-PEG24</b></p> <p>Cat. No.: HY-140998</p>	<p><b>Mal-PEG4-Lys(t-Boc)-NH-m-PEG24</b></p> <p>Cat. No.: HY-140999</p>
<p>Mal-PEG4-Glu(TFP ester)-NH-m-PEG24 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG4-Lys(t-Boc)-NH-m-PEG24 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG4-Lys(TFA)-NH-m-PEG24</b></p> <p>Cat. No.: HY-141000</p>	<p><b>Mal-PEG4-OH</b></p> <p>Cat. No.: HY-135048</p>
<p>Mal-PEG4-Lys(TFA)-NH-m-PEG24 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG4-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG5-acid</b></p> <p>Cat. No.: HY-126962</p>	<p><b>Mal-PEG5-Boc</b></p> <p>Cat. No.: HY-140965</p>
<p>Mal-PEG5-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.12%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Mal-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG5-C2-NH2 hydrochloride</b></p> <p>Cat. No.: HY-138375</p>	<p><b>Mal-PEG5-mal</b></p> <p>Cat. No.: HY-133352</p>
<p>Mal-PEG5-C2-NH2 hydrochloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG5-mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG5-NHS ester</b></p> <p>Cat. No.: HY-126888</p>	<p><b>Mal-PEG5-PFP ester</b></p> <p>Cat. No.: HY-133210</p>
<p>Mal-PEG5-NHS ester is an Alkyl/ether and PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG5-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>



<p><b>Mal-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-126963</p>	<p><b>Mal-PEG6-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133211</p>
<p>Mal-PEG6-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 98.96%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Mal-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG6-mal</b></p> <p style="text-align: right;">Cat. No.: HY-133353</p>	<p><b>Mal-PEG6-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-130454</p>
<p>Mal-PEG6-mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Mal-PEG6-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 96.89%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>
<p><b>Mal-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130334</p>	<p><b>Mal-PEG8-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140970</p>
<p>Mal-PEG8-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Mal-PEG8-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PEG8-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133212</p>	<p><b>Mal-PEG8-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-126887</p>
<p>Mal-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 95.08%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>Mal-PEG8-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Mal-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-138467</p>	<p><b>Mal-PEG4-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140037</p>
<p>Mal-PFP ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>Mal-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Maleimide-C10-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-W009037</p>	<p><b>Maleimido-tri(ethylene glycol)-propionic acid</b> (Mal-PEG3-acid)</p> <p style="text-align: right;">Cat. No.: HY-130426</p>
<p>Maleimide-C10-NHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Maleimido-tri(ethylene glycol)-propionic acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> 99.14% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Me-PEG18-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-138400</p>	<p><b>Me-PEG4-Me</b></p> <p style="text-align: right;">Cat. No.: HY-W013522</p>
<p>Me-PEG18-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Me-PEG4-Me is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.57% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>Methoxy-Tr-NH-PEG7</b></p> <p style="text-align: right;">Cat. No.: HY-141229</p>	<p><b>Methyl 1-Boc-azetidine-3-carboxylate</b></p> <p style="text-align: right;">Cat. No.: HY-40151</p>
<p>Methoxy-Tr-NH-PEG7 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyl 1-Boc-azetidine-3-carboxylate is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Methyl 1-Boc-azetidine-3-carboxylate is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>1</sup>.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 500 mg, 1 g</p>
<p><b>Methyl 1-Cbz-azetidine-3-carboxylate</b></p> <p style="text-align: right;">Cat. No.: HY-W019226</p>	<p><b>Methyl 3-hydroxypropanoate</b></p> <p style="text-align: right;">Cat. No.: HY-42569</p>
<p>Methyl 1-Cbz-azetidine-3-carboxylate is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Methyl 1-Cbz-azetidine-3-carboxylate is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>1</sup>.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyl 3-hydroxypropanoate is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 5 g</p>
<p><b>Methyl acetate-PEG1</b></p> <p style="text-align: right;">Cat. No.: HY-W096155</p>	<p><b>Methyl acetate-PEG1-methyl acetate</b></p> <p style="text-align: right;">Cat. No.: HY-134716</p>
<p>Methyl acetate-PEG1 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyl acetate-PEG1-methyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>













<p><b>Methyl acetate-PEG2-propanol</b></p> <p>Cat. No.: HY-138462</p>	<p><b>Methyl azetidine-3-carboxylate hydrochloride</b></p> <p>Cat. No.: HY-33615</p>
<p>Methyl acetate-PEG2-propanol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyl azetidine-3-carboxylate hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Methyl azetidine-3-carboxylate hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>
<p><b>Methyl propionate-PEG12</b></p> <p>Cat. No.: HY-130186</p>	<p><b>Methyl-PEG2-alcohol</b></p> <p>Cat. No.: HY-W012862</p>
<p>Methyl propionate-PEG12 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyl-PEG2-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyl-PEG3-Ald</b></p> <p>Cat. No.: HY-132092</p>	<p><b>Methyl-PEG3-bromide</b></p> <p>Cat. No.: HY-W094307</p>
<p>Methyl-PEG3-Ald is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyl-PEG3-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyl-PEG4-acyl chloride</b></p> <p>Cat. No.: HY-132095</p>	<p><b>Methylacetamide-PEG3-NH2</b></p> <p>Cat. No.: HY-134712</p>
<p>Methyl-PEG4-acyl chloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methylacetamide-PEG3-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methylamino-PEG1-acid</b></p> <p>Cat. No.: HY-140000</p>	<p><b>Methylamino-PEG1-Boc</b> (Methylamino-PEG1-t-butyl ester)</p> <p>Cat. No.: HY-140154</p>
<p>Methylamino-PEG1-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methylamino-PEG1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Methylamino-PEG2-acid</b></p> <p>Cat. No.: HY-140153</p>	<p><b>Methylamino-PEG2-Boc</b> (Methylamino-PEG2-t-butyl ester)</p> <p>Cat. No.: HY-140155</p>
<p>Methylamino-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Methylamino-PEG2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methylamino-PEG3-acid</b></p> <p>Cat. No.: HY-132075</p>	<p><b>Methylamino-PEG3-azide</b></p> <p>Cat. No.: HY-140158</p>
<p>Methylamino-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Methylamino-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methylamino-PEG3-t-butyl ester</b></p> <p>Cat. No.: HY-132071</p>	<p><b>Methylamino-PEG4-acid</b></p> <p>Cat. No.: HY-135153</p>
<p>Methylamino-PEG3-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Methylamino-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methylamino-PEG4-Boc</b> (Methylamino-PEG4-t-butyl ester)</p> <p>Cat. No.: HY-140156</p>	<p><b>Methylamino-PEG5-azide</b></p> <p>Cat. No.: HY-140159</p>
<p>Methylamino-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Methylamino-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methylamino-PEG7-benzyl</b></p> <p>Cat. No.: HY-132019</p>	<p><b>Methyltetrazine-acid</b></p> <p>Cat. No.: HY-141263</p>
<p>Methylamino-PEG7-benzyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.46% <b>Clinical Data:</b> <b>Size:</b> 25 mg, 100 mg</p>

<p><b>Methyltetrazine-amido-PEG5-alkyne</b></p> <p>Cat. No.: HY-141276</p>	<p><b>Methyltetrazine-amido-PEG7-azide</b></p> <p>Cat. No.: HY-141277</p>
<p>Methyltetrazine-amido-PEG5-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-amido-PEG7-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG12-DBCO</b></p> <p>Cat. No.: HY-140312</p>	<p><b>Methyltetrazine-PEG13-acid</b></p> <p>Cat. No.: HY-141266</p>
<p>Methyltetrazine-PEG12-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg</p>	<p>Methyltetrazine-PEG13-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG13-Boc</b></p> <p>Cat. No.: HY-141281</p>	<p><b>Methyltetrazine-PEG13-NHS ester</b></p> <p>Cat. No.: HY-141271</p>
<p>Methyltetrazine-PEG13-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG13-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG24-amine</b></p> <p>Cat. No.: HY-141262</p>	<p><b>Methyltetrazine-PEG24-Boc</b></p> <p>Cat. No.: HY-141282</p>
<p>Methyltetrazine-PEG24-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG24-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG24-NH-Boc</b></p> <p>Cat. No.: HY-141280</p>	<p><b>Methyltetrazine-PEG24-NHS ester</b></p> <p>Cat. No.: HY-141272</p>
<p>Methyltetrazine-PEG24-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG24-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Methyltetrazine-PEG25-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141267</p>	<p><b>Methyltetrazine-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141264</p>
<p>Methyltetrazine-PEG25-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-141261</p>	<p><b>Methyltetrazine-PEG4-amine hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-133505</p>
<p>Methyltetrazine-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG4-amine (hydrochloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG4-maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-141278</p>	<p><b>Methyltetrazine-PEG4-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141279</p>
<p>Methyltetrazine-PEG4-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 10 mg, 25 mg, 100 mg</p>	<p>Methyltetrazine-PEG4-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG4-SSPy</b></p> <p style="text-align: right;">Cat. No.: HY-133508</p>	<p><b>Methyltetrazine-PEG5-alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-141275</p>
<p>Methyltetrazine-PEG4-SSPy is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG5-alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG5-methyltetrazine</b></p> <p style="text-align: right;">Cat. No.: HY-133467</p>	<p><b>Methyltetrazine-PEG5-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141269</p>
<p>Methyltetrazine-PEG5-methyltetrazine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG5-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Methyltetrazine-PEG5-triethoxysilane</b></p> <p>Cat. No.: HY-141283</p>	<p><b>Methyltetrazine-PEG6-maleimide</b></p> <p>Cat. No.: HY-133468</p>
<p>Methyltetrazine-PEG5-triethoxysilane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG6-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG8-acid</b></p> <p>Cat. No.: HY-141265</p>	<p><b>Methyltetrazine-PEG8-N3</b></p> <p>Cat. No.: HY-133469</p>
<p>Methyltetrazine-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG8-N3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-PEG8-NHS ester</b></p> <p>Cat. No.: HY-141270</p>	<p><b>Methyltetrazine-PEG8-PFP ester</b></p> <p>Cat. No.: HY-141273</p>
<p>Methyltetrazine-PEG8-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-PEG8-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-Ph-NHS ester</b></p> <p>Cat. No.: HY-130283</p>	<p><b>Methyltetrazine-Ph-PEG4-azide</b></p> <p>Cat. No.: HY-130508</p>
<p>Methyltetrazine-Ph-NHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-Ph-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Methyltetrazine-propylamine</b></p> <p>Cat. No.: HY-141260</p>	<p><b>Methyltetrazine-Sulfo-NHS ester sodium</b></p> <p>Cat. No.: HY-141268</p>
<p>Methyltetrazine-propylamine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Methyltetrazine-Sulfo-NHS ester (sodium) is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

<p><b>ML 2-14</b></p> <p style="text-align: right;">Cat. No.: HY-132991</p>	<p><b>Monoethyl pimelate</b></p> <p style="text-align: right;">Cat. No.: HY-W016087</p>
<p>ML 2-14 is a PROTAC linker, which belongs to a polyethylene glycol (PEG) linker. ML 2-14 can be used in the synthesis of the PROTAC.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Monoethyl pimelate is a &lt;b&gt;PROTAC linker, which refers to the alkyl/ether composition. Monoethyl pimelate can be used in the synthesis of (S,R,S)-AHPC-Me-C7 ester, a specific BCL-X<sub>1</sub> PROTAC degrader.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>MS-PEG1-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138482</p>	<p><b>Ms-PEG10-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140386</p>
<p>MS-PEG1-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ms-PEG10-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>MS-PEG10-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138344</p>	<p><b>Ms-PEG12-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140387</p>
<p>MS-PEG10-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ms-PEG12-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ms-PEG12-m</b></p> <p style="text-align: right;">Cat. No.: HY-115392</p>	<p><b>MS-PEG12-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138371</p>
<p>Ms-PEG12-m is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>MS-PEG12-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ms-PEG2-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140383</p>	<p><b>Ms-PEG2-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-140382</p>
<p>Ms-PEG2-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ms-PEG2-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>



<p><b>Ms-PEG3-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-134680</p>	<p><b>MS-PEG3-dodecyl</b></p> <p style="text-align: right;">Cat. No.: HY-132063</p>
<p>Ms-PEG3-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>MS-PEG3-dodecyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ms-PEG3-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-134683</p>	<p><b>MS-PEG3-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132107</p>
<p>Ms-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>MS-PEG3-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ms-PEG4-Ms</b> (1,11-Bis(methanesulfonyloxy)-3,6,9-trioxandecane)</p> <p style="text-align: right;">Cat. No.: HY-42774</p>	<p><b>MS-PEG4-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132109</p>
<p>Ms-PEG4-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p>MS-PEG4-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>MS-PEG4-THP</b></p> <p style="text-align: right;">Cat. No.: HY-W096074</p>	<p><b>Ms-PEG5-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140384</p>
<p>MS-PEG4-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ms-PEG5-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>MS-PEG5-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138385</p>	<p><b>Ms-PEG6-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-138324</p>
<p>MS-PEG5-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ms-PEG6-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>






<p><b>Ms-PEG6-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138315</p>	<p><b>Ms-PEG7-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-130152</p>
<p>Ms-PEG6-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ms-PEG7-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ms-PEG8-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140385</p>	<p><b>MS-PEG8-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138337</p>
<p>Ms-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>MS-PEG8-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N'-Boc-N-(Gly-Oleoyl)-Lys</b></p> <p style="text-align: right;">Cat. No.: HY-141291</p>	<p><b>N,N'-bis-(Acid-PEG3)-benzothiazole Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141043</p>
<p>N'-Boc-N-(Gly-Oleoyl)-Lys is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N,N'-bis-(Acid-PEG3)-benzothiazole Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N,N'-bis-(azide-PEG3)-chlorocyclohexenyl Cy7</b></p> <p style="text-align: right;">Cat. No.: HY-141079</p>	<p><b>N,N'-bis-(azide-PEG3)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141060</p>
<p>N,N'-bis-(azide-PEG3)-chlorocyclohexenyl Cy7 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>NN'-bis-(azide-PEG3)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N,N'-bis-(propargyl-PEG4)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141054</p>	<p><b>N,N'-DME-N,N'-Bis-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141028</p>
<p>N,N'-bis-(propargyl-PEG4)-Cy5 (chloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N,N'-DME-N,N'-Bis-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>









<p><b>N,N'-DME-N-PEG2-Boc</b></p> <p>Cat. No.: HY-140163</p>	<p><b>N,N-Diethanol amine-PEG4-Boc</b></p> <p>Cat. No.: HY-141252</p>
<p>N,N'-DME-N-PEG2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N,N-Diethanol amine-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(4-Carboxycyclohexylmethyl)maleimide</b></p> <p>Cat. No.: HY-42359</p>	<p><b>N-(Ac-PEG3)-N'-(azide-PEG3)-Cy7 chloride</b></p> <p>Cat. No.: HY-141077</p>
<p>N-(4-Carboxycyclohexylmethyl)maleimide is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Ac-PEG3)-N'-(azide-PEG3)-Cy7 (chloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Acid-PEG2)-N-bis(PEG3-azide)</b></p> <p>Cat. No.: HY-140521</p>	<p><b>N-(acid-PEG3)-N-bis(PEG3-azide)</b></p> <p>Cat. No.: HY-140520</p>
<p>N-(Acid-PEG2)-N-bis(PEG3-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(acid-PEG3)-N-bis(PEG3-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Amino-PEG1)-N-bis(PEG2-propargyl)</b></p> <p>Cat. No.: HY-140086</p>	<p><b>N-(Amino-PEG2)-N-bis(PEG3-azide)</b></p> <p>Cat. No.: HY-140087</p>
<p>N-(Amino-PEG1)-N-bis(PEG2-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Amino-PEG2)-N-bis(PEG3-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>N-(Amino-PEG3)-N-bis(PEG3-acid)</b></p> <p>Cat. No.: HY-140245</p>	<p><b>N-(Amino-PEG3)-N-bis(PEG3-Boc)</b></p> <p>Cat. No.: HY-140253</p>
<p>N-(Amino-PEG3)-N-bis(PEG3-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Amino-PEG3)-N-bis(PEG3-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-(Amino-PEG3)-N-bis(PEG4-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140254</p> <p>N-(Amino-PEG3)-N-bis(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(Amino-PEG4)-N-Biotin-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140532</p> <p>N-(Amino-PEG4)-N-Biotin-PEG4-acid is a biotin-labeled, PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Amino-PEG4)-N-bis(PEG4-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140255</p> <p>N-(Amino-PEG4)-N-bis(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(Amino-PEG5)-N-bis(PEG4-acid)</b></p> <p style="text-align: right;">Cat. No.: HY-130695</p> <p>N-(Amino-PEG5)-N-bis(PEG4-acid) is a PEG-based PROTAC linker used in the synthesis of PROTACs. N-(Amino-PEG5)-N-bis(PEG4-acid) contains an amino group with two terminal carboxylic acids.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Aminoxy-PEG2)-N-bis(PEG3-propargyl)</b></p> <p style="text-align: right;">Cat. No.: HY-140091</p> <p>N-(Aminoxy-PEG2)-N-bis(PEG3-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(Aminoxy-PEG3)-N-bis(PEG4-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140446</p> <p>N-(Aminoxy-PEG3)-N-bis(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(azide-PEG3)-N'-(Amine-C3-Amide-PEG4)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141065</p> <p>N-(azide-PEG3)-N'-(Amine-C3-Amide-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(azide-PEG3)-N'-(m-PEG4)-Benzothiazole Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141064</p> <p>N-(azide-PEG3)-N'-(m-PEG4)-Benzothiazole Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(azide-PEG3)-N'-(Mal-PEG4)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141066</p> <p>N-(azide-PEG3)-N'-(Mal-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(Azide-PEG3)-N'-(PEG4-acid)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141041</p> <p>N-(Azide-PEG3)-N'-(PEG4-acid)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-(Azide-PEG3)-N'-(PEG4-NHS ester)-Cy5</b></p> <p>Cat. No.: HY-141048</p>	<p><b>N-(Azido-PEG2)-N-bis(PEG4-Boc)</b></p> <p>Cat. No.: HY-140867</p>
<p>N-(Azide-PEG3)-N'-(PEG4-NHS ester)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Azido-PEG2)-N-bis(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG2)-N-Boc-PEG3-acid</b></p> <p>Cat. No.: HY-140524</p>	<p><b>N-(Azido-PEG2)-N-Boc-PEG3-Boc</b></p> <p>Cat. No.: HY-140562</p>
<p>N-(Azido-PEG2)-N-Boc-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Azido-PEG2)-N-Boc-PEG3-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG2)-N-Boc-PEG3-NHS ester</b></p> <p>Cat. No.: HY-140559</p>	<p><b>N-(Azido-PEG2)-N-Boc-PEG4-acid</b></p> <p>Cat. No.: HY-140525</p>
<p>N-(Azido-PEG2)-N-Boc-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Azido-PEG2)-N-Boc-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG2)-N-Boc-PEG4-Boc</b></p> <p>Cat. No.: HY-140563</p>	<p><b>N-(Azido-PEG2)-N-Boc-PEG4-NHS ester</b></p> <p>Cat. No.: HY-140560</p>
<p>N-(Azido-PEG2)-N-Boc-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Azido-PEG2)-N-Boc-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG2)-N-Fluorescein-PEG3-acid</b></p> <p>Cat. No.: HY-140544</p>	<p><b>N-(Azido-PEG3)-N-(PEG2-amine)-PEG3-acid</b></p> <p>Cat. No.: HY-140247</p>
<p>N-(Azido-PEG2)-N-Fluorescein-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Azido-PEG3)-N-(PEG2-amine)-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.58%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-(Azido-PEG3)-N-(PEG2-NH-Boc)-PEG3-acid</b></p> <p>Cat. No.: HY-141287</p> <p>N-(Azido-PEG3)-N-(PEG2-NH-Boc)-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(Azido-PEG3)-N-Biotin-PEG4-methyl ester</b></p> <p>Cat. No.: HY-140581</p> <p>N-(Azido-PEG3)-N-Biotin-PEG4-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG3)-N-bis(PEG1-t-butyl ester)</b></p> <p>Cat. No.: HY-140873</p> <p>N-(Azido-PEG3)-N-bis(PEG1-t-butyl ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(Azido-PEG3)-N-bis(PEG3-acid)</b></p> <p>Cat. No.: HY-140517</p> <p>N-(Azido-PEG3)-N-bis(PEG3-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG3)-N-bis(PEG3-Boc)</b></p> <p>Cat. No.: HY-140868</p> <p>N-(Azido-PEG3)-N-bis(PEG3-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(Azido-PEG3)-N-bis(PEG4-acid)</b></p> <p>Cat. No.: HY-140518</p> <p>N-(Azido-PEG3)-N-bis(PEG4-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG3)-N-bis(PEG4-Boc)</b></p> <p>Cat. No.: HY-140869</p> <p>N-(Azido-PEG3)-N-bis(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(Azido-PEG3)-N-Boc-PEG3-acid</b></p> <p>Cat. No.: HY-140526</p> <p>N-(Azido-PEG3)-N-Boc-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG3)-N-Boc-PEG3-NHS ester</b></p> <p>Cat. No.: HY-140561</p> <p>N-(Azido-PEG3)-N-Boc-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>N-(Azido-PEG3)-N-Boc-PEG3-t-butyl ester</b></p> <p>Cat. No.: HY-140564</p> <p>N-(Azido-PEG3)-N-Boc-PEG3-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-(Azido-PEG3)-N-Boc-PEG4-acid</b></p> <p>Cat. No.: HY-130598</p>	<p><b>N-(Azido-PEG3)-N-Boc-PEG4-Boc</b></p> <p>Cat. No.: HY-140565</p>
<p>N-(Azido-PEG3)-N-Boc-PEG4-acid is a PEG-based PROTAC linker with a terminal azide group and is used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Azido-PEG3)-N-Boc-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG3)-N-Fluorescein-PEG3-acid</b></p> <p>Cat. No.: HY-130768</p>	<p><b>N-(Azido-PEG3)-N-Fluorescein-PEG4-acid</b></p> <p>Cat. No.: HY-140546</p>
<p>N-(Azido-PEG3)-N-Fluorescein-PEG3-acid is a PEG-based PROTAC linker which contains azide, fluorescein and carboxylic acid moieties.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg</p>	<p>N-(Azido-PEG3)-N-Fluorescein-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG3)-NH-PEG3-acid</b></p> <p>Cat. No.: HY-140552</p>	<p><b>N-(Azido-PEG4)-biotin</b></p> <p>Cat. No.: HY-140919</p>
<p>N-(Azido-PEG3)-NH-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Azido-PEG4)-biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG4)-N-bis(PEG4-acid)</b></p> <p>Cat. No.: HY-140519</p>	<p><b>N-(Azido-PEG4)-N-bis(PEG4-NHS ester)</b></p> <p>Cat. No.: HY-140866</p>
<p>N-(Azido-PEG4)-N-bis(PEG4-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Azido-PEG4)-N-bis(PEG4-NHS ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Azido-PEG4)-N-bis(PEG4-t-butyl ester)</b></p> <p>Cat. No.: HY-140870</p>	<p><b>N-(Azido-PEG4)-N-Boc-PEG4-Boc</b></p> <p>Cat. No.: HY-140567</p>
<p>N-(Azido-PEG4)-N-bis(PEG4-t-butyl ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Azido-PEG4)-N-Boc-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-(Azido-PEG4)-N-Boc-PEG4-NHS ester</b></p> <p>Cat. No.: HY-130312</p>	<p><b>N-(Biotin)-N-bis(PEG1-alcohol)</b></p> <p>Cat. No.: HY-140949</p>
<p>N-(Azido-PEG4)-N-Boc-PEG4-NHS ester is a PEG-based PROTAC linker with a terminal azide group. N-(Azido-PEG4)-N-Boc-PEG4-NHS ester is used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Biotin)-N-bis(PEG1-alcohol) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Biotin-PEG4)-N-bis(PEG4-acid)</b></p> <p>Cat. No.: HY-140531</p>	<p><b>N-(Biotin-PEG4)-N-bis(PEG4-Boc)</b></p> <p>Cat. No.: HY-141288</p>
<p>N-(Biotin-PEG4)-N-bis(PEG4-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Biotin-PEG4)-N-bis(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Boc-PEG1)-N-bis(PEG2-propargyl)</b></p> <p>Cat. No.: HY-140090</p>	<p><b>N-(Boc-PEG2)-N-bis(PEG3-azide)</b></p> <p>Cat. No.: HY-140872</p>
<p>N-(Boc-PEG1)-N-bis(PEG2-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Boc-PEG2)-N-bis(PEG3-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Boc-PEG3)-N-bis(PEG2-alcohol)</b></p> <p>Cat. No.: HY-141254</p>	<p><b>N-(Boc-PEG3)-N-bis(PEG3-acid)</b></p> <p>Cat. No.: HY-140536</p>
<p>N-(Boc-PEG3)-N-bis(PEG2-alcohol) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Boc-PEG3)-N-bis(PEG3-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b>  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>N-(Boc-PEG3)-N-bis(PEG3-azide)</b></p> <p>Cat. No.: HY-140871</p>	<p><b>N-(Boc-PEG4)-NH-PEG4-NH-Boc</b></p> <p>Cat. No.: HY-140259</p>
<p>N-(Boc-PEG3)-N-bis(PEG3-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Boc-PEG4)-NH-PEG4-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-(Boc-PEG5)-N-bis(PEG4-acid)</b></p> <p>Cat. No.: HY-140537</p>	<p><b>N-(DBCO-PEG4)-N-Biotin-PEG4-NHS</b></p> <p>Cat. No.: HY-140582</p>
<p>N-(Boc-PEG5)-N-bis(PEG4-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(DBCO-PEG4)-N-Biotin-PEG4-NHS is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Hydroxy-PEG3)-N-bis(PEG4-Boc)</b></p> <p>Cat. No.: HY-141251</p>	<p><b>N-(Hydroxy-PEG3)-N-Boc-PEG4-Boc</b> (N-(Hydroxy-PEG3)-N-Boc-PEG4-t-butyl ester)</p> <p>Cat. No.: HY-140568</p>
<p>N-(Hydroxy-PEG3)-N-bis(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Hydroxy-PEG3)-N-Boc-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(m-PEG4)-3,3-Dimethyl-3H-indole-N'-(acid-PEG3)-benzothiazole</b></p> <p>Cat. No.: HY-141039</p>	<p><b>N-(m-PEG4)-N'-(4-Hydroxycyclohexyl-1-amido-PEG4)-Cy5</b></p> <p>Cat. No.: HY-141075</p>
<p>N-(m-PEG4)-3,3-Dimethyl-3H-indole-N'-(acid-PEG3)-benzothiazole is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(m-PEG4)-N'-(4-Hydroxycyclohexyl-1-amido-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(m-PEG4)-N'-(Acid-PEG3)-benzothiazole Cy5</b></p> <p>Cat. No.: HY-141038</p>	<p><b>N-(m-PEG4)-N'-(amino-PEG3)-Cy5</b></p> <p>Cat. No.: HY-141056</p>
<p>N-(m-PEG4)-N'-(Acid-PEG3)-benzothiazole Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(m-PEG4)-N'-(amino-PEG3)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(m-PEG4)-N'-(azide-PEG3)-Cy5</b></p> <p>Cat. No.: HY-141063</p>	<p><b>N-(m-PEG4)-N'-(azide-PEG4)-Cy3</b></p> <p>Cat. No.: HY-141030</p>
<p>N-(m-PEG4)-N'-(azide-PEG3)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg</p>	<p>N-(m-PEG4)-N'-(azide-PEG4)-Cy3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-(m-PEG4)-N'-(azide-PEG4)-Cy7</b></p> <p>Cat. No.: HY-141078</p>	<p><b>N-(m-PEG4)-N'-(Biotin-PEG2-amido-PEG4)-Cy5</b></p> <p>Cat. No.: HY-141069</p>
<p>N-(m-PEG4)-N'-(azide-PEG4)-Cy7 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(m-PEG4)-N'-(Biotin-PEG2-amido-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(m-PEG4)-N'-(biotin-PEG3)-Cy5</b></p> <p>Cat. No.: HY-141070</p>	<p><b>N-(m-PEG4)-N'-(DBCO-PEG4)-Cy5</b></p> <p>Cat. No.: HY-141071</p>
<p>N-(m-PEG4)-N'-(biotin-PEG3)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(m-PEG4)-N'-(DBCO-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(m-PEG4)-N'-(hydroxy-PEG2)-Cy5</b></p> <p>Cat. No.: HY-141073</p>	<p><b>N-(m-PEG4)-N'-(m-PEG4)-O-(m-PEG4)-O'-(azide-PEG4)-Cy5</b></p> <p>Cat. No.: HY-141068</p>
<p>N-(m-PEG4)-N'-(hydroxy-PEG2)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(m-PEG4)-N'-(m-PEG4)-O-(m-PEG4)-O'-(azide-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(m-PEG4)-N'-(m-PEG4)-O-(m-PEG4)-O'-(propargyl-PEG4)-Cy5</b></p> <p>Cat. No.: HY-141053</p>	<p><b>N-(m-PEG4)-N'-(PEG2-acid)-Cy5</b></p> <p>Cat. No.: HY-141035</p>
<p>N-(m-PEG4)-N'-(m-PEG4)-O-(m-PEG4)-O'-(propargyl-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(m-PEG4)-N'-(PEG2-acid)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(m-PEG4)-N'-(PEG2-NHS ester)-Cy5</b></p> <p>Cat. No.: HY-141045</p>	<p><b>N-(m-PEG4)-N'-(PEG3-Mal)-Cy5</b></p> <p>Cat. No.: HY-141076</p>
<p>N-(m-PEG4)-N'-(PEG2-NHS ester)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(m-PEG4)-N'-(PEG3-Mal)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>




<p><b>N-(m-PEG4)-N'-(PEG4-acid)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141036</p>	<p><b>N-(m-PEG4)-N'-(PEG4-NHS ester)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141046</p>
<p>N-(m-PEG4)-N'-(PEG4-acid)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(m-PEG4)-N'-(PEG4-NHS ester)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(m-PEG4)-N'-hydroxypropyl-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141074</p>	<p><b>N-(m-PEG9)-N'-(PEG5-acid)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141037</p>
<p>N-(m-PEG4)-N'-hydroxypropyl-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(m-PEG9)-N'-(PEG5-acid)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(m-PEG9)-N'-(propargyl-PEG8)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141051</p>	<p><b>N-(Mal-PEG6)-N-bis(PEG3-amine)</b></p> <p style="text-align: right;">Cat. No.: HY-140248</p>
<p>N-(m-PEG9)-N'-(propargyl-PEG8)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Mal-PEG6)-N-bis(PEG3-amine) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Mal-PEG6)-N-bis(PEG7-TCO)</b></p> <p style="text-align: right;">Cat. No.: HY-141009</p>	<p><b>N-(NHS-PEG3)-N-bis(PEG3-azide)</b></p> <p style="text-align: right;">Cat. No.: HY-140864</p>
<p>N-(Mal-PEG6)-N-bis(PEG7-TCO) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(NHS-PEG3)-N-bis(PEG3-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(PEG1-OH)-N-Boc-PEG2-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140569</p>	<p><b>N-(PEG2-Boc)-N-bis(PEG2-propargyl)</b></p> <p style="text-align: right;">Cat. No.: HY-140093</p>
<p>N-(PEG1-OH)-N-Boc-PEG2-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(PEG2-Boc)-N-bis(PEG2-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-(PEG2-C2-acid)-N-bis(PEG2-propargyl)</b></p> <p>Cat. No.: HY-140085</p>	<p><b>N-(PEG3-acid)-N-bis(PEG3-amine)</b></p> <p>Cat. No.: HY-140246</p>
<p>N-(PEG2-C2-acid)-N-bis(PEG2-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(PEG3-acid)-N-bis(PEG3-amine) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Propanoic acid)-N-bis(m-PEG12)</b></p> <p>Cat. No.: HY-141289</p>	<p><b>N-(Propargyl-PEG2)-N-Boc-PEG3-t-butyl ester</b></p> <p>Cat. No.: HY-140094</p>
<p>N-(Propanoic acid)-N-bis(m-PEG12) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Propargyl-PEG2)-N-Boc-PEG3-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Propargyl-PEG4)-biocytin</b></p> <p>Cat. No.: HY-140926</p>	<p><b>N-(Propargyl-PEG4)-N-bis(PEG4-acid)</b></p> <p>Cat. No.: HY-140084</p>
<p>N-(Propargyl-PEG4)-biocytin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(Propargyl-PEG4)-N-bis(PEG4-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Propargyl-PEG4-carbonyl)-N-bis(PEG1-methyl ester)</b></p> <p>Cat. No.: HY-140095</p>	<p><b>N-(t-Boc-Aminoxy-PEG2)-N-bis(PEG3-propargyl)</b></p> <p>Cat. No.: HY-140092</p>
<p>N-(Propargyl-PEG4-carbonyl)-N-bis(PEG1-methyl ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-(t-Boc-Aminoxy-PEG2)-N-bis(PEG3-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-(Tos-PEG4)-N-bis(PEG4-Boc)</b></p> <p>Cat. No.: HY-140393</p>	<p><b>N-Acetylpyrrolidine-PEG2-Br</b></p> <p>Cat. No.: HY-134730</p>
<p>N-(Tos-PEG4)-N-bis(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Acetylpyrrolidine-PEG2-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

**N-Azido-PEG4-N-Boc-N-PEG3-Boc** Cat. No.: HY-140566


N-Azido-PEG4-N-Boc-N-PEG3-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**N-Benzyl-N-bis(PEG3-acid)** Cat. No.: HY-140542


N-Benzyl-N-bis(PEG3-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**N-Benzyl-N-bis-PEG2** Cat. No.: HY-140586


N-Benzyl-N-bis-PEG2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**N-Benzyl-N-bis-PEG4** Cat. No.: HY-140587


N-Benzyl-N-bis-PEG4 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**N-Biotin-N-bis(PEG4-acid)** Cat. No.: HY-140528


N-Biotin-N-bis(PEG4-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**N-bis(t-boc-N-amido-PEG3)-N-(PEG3-acid) (hydrochloride)** Cat. No.: HY-143849


N-bis(t-boc-N-amido-PEG3)-N-(PEG3-acid) hydrochloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**N-Boc-4-pentyne-1-amine** Cat. No.: HY-43721


N-Boc-4-pentyne-1-amine is a PROTAC linker, which refers to the alkyl chain composition. N-Boc-4-pentyne-1-amine can be used in the synthesis of the PROTAC MG-277 (HY-130122).



**Purity:** ≥98.0%  
**Clinical Data:** No Development Reported  
**Size:** 10 mM × 1 mL, 10 mg, 25 mg, 50 mg, 100 mg

**N-Boc-C1-PEG3-C3-NH2** Cat. No.: HY-W004640


N-Boc-C1-PEG3-C3-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** ≥98.0%  
**Clinical Data:** No Development Reported  
**Size:** 100 mg

**N-Boc-C1-PEG5-C3-NH2** Cat. No.: HY-134679


N-Boc-C1-PEG5-C3-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**N-Boc-cis-4-Hydroxy-D-proline** Cat. No.: HY-W002887

N-Boc-cis-4-Hydroxy-D-proline is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-Boc-cis-4-Hydroxy-D-proline is also an alkyl chain-based PROTAC linker that can be used in the synth.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 g, 5 g

### N-Boc-cis-4-hydroxy-D-proline methyl ester

Cat. No.: HY-W002680

N-Boc-cis-4-hydroxy-D-proline methyl ester is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-Boc-cis-4-hydroxy-D-proline methyl ester is also an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>1</sup>.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 g

### N-Boc-cis-4-hydroxy-L-proline

Cat. No.: HY-W002886

N-Boc-cis-4-hydroxy-L-proline is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-Boc-cis-4-hydroxy-L-proline is also an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>1</sup>.




**Purity:** ≥97.0%  
**Clinical Data:** No Development Reported  
**Size:** 500 mg

### N-Boc-cis-4-hydroxy-L-proline methyl ester

Cat. No.: HY-Y0755

N-Boc-cis-4-hydroxy-L-proline methyl ester is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-Boc-cis-4-hydroxy-L-proline methyl ester is also an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs<sup>1</sup>.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 100 mg, 250 mg

### N-Boc-diethanolamine

Cat. No.: HY-W044078

N-Boc-diethanolamine is an Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-diethanolamine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 500 mg

### N-Boc-N-bis(C2-PEG1-azide)

Cat. No.: HY-140554

N-Boc-N-bis(C2-PEG1-azide) is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### N-Boc-N-bis(PEG2-acid)

Cat. No.: HY-140538

N-Boc-N-bis(PEG2-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### N-Boc-N-bis(PEG2-OH)

Cat. No.: HY-117079

N-Boc-N-bis(PEG2-OH) is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-N-bis(PEG2-OH) is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### N-Boc-N-bis(PEG2-propargyl)

Cat. No.: HY-140089

N-Boc-N-bis(PEG2-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

### N-Boc-N-bis(PEG3-acid)

Cat. No.: HY-140539

N-Boc-N-bis(PEG3-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg











### N-Boc-N-bis(PEG3-azide)











Cat. No.: HY-140555











N-Boc-N-bis(PEG3-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.






**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg


<p><b>N-Boc-N-bis(PEG3-NHS ester)</b></p> <p style="text-align: right;">Cat. No.: HY-140557</p>	<p><b>N-Boc-N-bis(PEG4-acid)</b></p> <p style="text-align: right;">Cat. No.: HY-140540</p>
<p>N-Boc-N-bis(PEG3-NHS ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-N-bis(PEG4-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Boc-N-bis(PEG4-azide)</b></p> <p style="text-align: right;">Cat. No.: HY-140556</p>	<p><b>N-Boc-N-bis(PEG4-NHS ester)</b></p> <p style="text-align: right;">Cat. No.: HY-140558</p>
<p>N-Boc-N-bis(PEG4-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-N-bis(PEG4-NHS ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Boc-N-bis(PEG4-OH)</b></p> <p style="text-align: right;">Cat. No.: HY-130449</p>	<p><b>N-Boc-N-bis-PEG5</b></p> <p style="text-align: right;">Cat. No.: HY-140553</p>
<p>N-Boc-N-bis(PEG4-OH) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-N-bis(PEG4-OH) is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-N-bis-PEG5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Boc-PEG-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132088</p>	<p><b>N-Boc-PEG10-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W190857</p>
<p>N-Boc-PEG-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-PEG10-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Boc-PEG12-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141193</p>	<p><b>N-Boc-PEG16-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141194</p>
<p>N-Boc-PEG12-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-PEG16-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-Boc-PEG2-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-130503</p>	<p><b>N-Boc-PEG23-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-141197</p>
<p>N-Boc-PEG2-bromide is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG2-bromide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>N-Boc-PEG23-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Boc-PEG24-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141195</p>	<p><b>N-Boc-PEG3-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-W006445</p>
<p>N-Boc-PEG24-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-PEG3-bromide is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG3-bromide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Boc-PEG3-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132089</p>	<p><b>N-Boc-PEG36-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141196</p>
<p>N-Boc-PEG3-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-PEG36-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Boc-PEG4-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-W046471</p>	<p><b>N-Boc-PEG5-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141191</p>
<p>N-Boc-PEG4-bromide is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG4-bromide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg</p>	<p>N-Boc-PEG5-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg</p>
<p><b>N-Boc-PEG5-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-120702</p>	<p><b>N-Boc-PEG6-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W071584</p>
<p>N-Boc-PEG5-bromide is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG5-bromide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-PEG6-alcohol is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG6-alcohol is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>











<p><b>N-Boc-PEG7-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-130505</p>	<p><b>N-Boc-PEG8-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141192</p>
<p>N-Boc-PEG7-alcohol is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG7-alcohol is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-PEG8-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Boc-PEG9-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W071583</p>	<p><b>N-Boc-piperazine</b></p> <p style="text-align: right;">Cat. No.: HY-30105</p>
<p>N-Boc-PEG9-alcohol is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG9-alcohol is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-piperazine is a Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTAC PD-1/PD-L1 degrader-1 (HY-131183).</p>  <p><b>Purity:</b> 99.96%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 500 mg</p>
<p><b>N-Boc-piperazine-C3-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-131184</p>	<p><b>N-Boc-serinol</b></p> <p style="text-align: right;">Cat. No.: HY-W018464</p>
<p>N-Boc-piperazine-C3-COOH is a PROTAC linker, which refers to the alkyl/ether composition. Boc-N-piperazine-C3-COOH can be used in the synthesis of PROTAC PD-1/PD-L1 degrader-1 (HY-131183).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Boc-serinol is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>
<p><b>N-Bromoacetyl-β-alanine</b></p> <p style="text-align: right;">Cat. No.: HY-141379</p>	<p><b>N-DBCO-N-bis(PEG2-C2-acid)</b></p> <p style="text-align: right;">Cat. No.: HY-140543</p>
<p>N-Bromoacetyl-β-alanine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs. N-Bromoacetyl-β-alanine is also a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p>N-DBCO-N-bis(PEG2-C2-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-DBCO-N-bis(PEG2-C2-NHS ester)</b></p> <p style="text-align: right;">Cat. No.: HY-140585</p>	<p><b>N-Desthiobiotin-N-bis(PEG4-NHS ester)</b></p> <p style="text-align: right;">Cat. No.: HY-140583</p>
<p>N-DBCO-N-bis(PEG2-C2-NHS ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Desthiobiotin-N-bis(PEG4-NHS ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>





<p><b>N-Desthiobiotin-N-bis(PEG4-t-butyl ester)</b></p> <p>Cat. No.: HY-140584</p>	<p><b>N-Ethyl-3,3,3-trifluoro-N-methylpropanamide-PEG2-Br</b></p> <p>Cat. No.: HY-134739</p>
<p>N-Desthiobiotin-N-bis(PEG4-t-butyl ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Ethyl-333-trifluoro-N-methylpropanamide-PEG2-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Ethyl-N-methylpropionamide-PEG1-Br</b></p> <p>Cat. No.: HY-134733</p>	<p><b>N-Ethylacetamide-PEG1-Br</b></p> <p>Cat. No.: HY-134735</p>
<p>N-Ethyl-N-methylpropionamide-PEG1-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Ethylacetamide-PEG1-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Ethylacetamide-PEG2-Br</b></p> <p>Cat. No.: HY-134731</p>	<p><b>N-Ethylpropionamide-PEG1-Br</b></p> <p>Cat. No.: HY-134732</p>
<p>N-Ethylacetamide-PEG2-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Ethylpropionamide-PEG1-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Fmoc-N'-(azido-PEG4)-L-Lysine</b></p> <p>Cat. No.: HY-140846</p>	<p><b>N-Fmoc-N'-(azido-PEG4)-L-Lysine-PFP ester</b></p> <p>Cat. No.: HY-140847</p>
<p>N-Fmoc-N'-(azido-PEG4)-L-Lysine is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Fmoc-N'-(azido-PEG4)-L-Lysine-PFP ester is an alkyl/ether and PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Hydroxypropyl-N'-(azide-PEG3)-Cy3</b></p> <p>Cat. No.: HY-141029</p>	<p><b>N-Mal-N-bis(PEG2-acid)</b></p> <p>Cat. No.: HY-140529</p>
<p>N-Hydroxypropyl-N'-(azide-PEG3)-Cy3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Mal-N-bis(PEG2-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>






















<p><b>N-Mal-N-bis(PEG2-amine)</b></p> <p style="text-align: right;">Cat. No.: HY-140570</p>	<p><b>N-Mal-N-bis(PEG2-C2-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140576</p>
<p>N-Mal-N-bis(PEG2-amine) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Mal-N-bis(PEG2-C2-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Mal-N-bis(PEG2-NH-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140574</p>	<p><b>N-Mal-N-bis(PEG2-NHS ester)</b></p> <p style="text-align: right;">Cat. No.: HY-140571</p>
<p>N-Mal-N-bis(PEG2-NH-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Mal-N-bis(PEG2-NHS ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.36% <b>Clinical Data:</b> <b>Size:</b> 25 mg</p>
<p><b>N-Mal-N-bis(PEG4-amine)</b></p> <p style="text-align: right;">Cat. No.: HY-140573</p>	<p><b>N-Mal-N-bis(PEG4-NH-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140575</p>
<p>N-Mal-N-bis(PEG4-amine) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Mal-N-bis(PEG4-NH-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Mal-N-bis(PEG4-NHS ester)</b></p> <p style="text-align: right;">Cat. No.: HY-140572</p>	<p><b>N-Me-N-bis(PEG2-propargyl)</b></p> <p style="text-align: right;">Cat. No.: HY-140043</p>
<p>N-Mal-N-bis(PEG4-NHS ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Me-N-bis(PEG2-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Me-N-bis(PEG4-acid)</b></p> <p style="text-align: right;">Cat. No.: HY-140579</p>	<p><b>N-Me-N-bis(PEG4-C2-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140157</p>
<p>N-Me-N-bis(PEG4-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Me-N-bis(PEG4-C2-Boc) is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>



<p><b>N-Me-N-bis-PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-140577</p>	<p><b>N-Me-N-bis-PEG4</b></p> <p style="text-align: right;">Cat. No.: HY-140578</p>
<p>N-Me-N-bis-PEG3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Me-N-bis-PEG4 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-methyl-N'-(azide-PEG3)-Cy3</b></p> <p style="text-align: right;">Cat. No.: HY-140010</p>	<p><b>N-Methyl-N'-(azido-PEG2-C5)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141062</p>
<p>N-methyl-N'-(azide-PEG3)-Cy3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Methyl-N'-(azido-PEG2-C5)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Methyl-N'-(hydroxy-PEG2)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141072</p>	<p><b>N-methyl-N'-(propargyl-PEG4)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141049</p>
<p>N-Methyl-N'-(hydroxy-PEG2)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-methyl-N'-(propargyl-PEG4)-Cy5 (chloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-methyl-N'-methyl-O-(m-PEG4)-O'-(acid-PEG5)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141040</p>	<p><b>N-Methyl-N'-methyl-O-(m-PEG4)-O'-(azide-PEG4)-Cy3</b></p> <p style="text-align: right;">Cat. No.: HY-141031</p>
<p>N-methyl-N'-methyl-O-(m-PEG4)-O'-(acid-PEG5)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Methyl-N'-methyl-O-(m-PEG4)-O'-(azide-PEG4)-Cy3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-methyl-N'-methyl-O-(m-PEG4)-O'-(azide-PEG4)-Cy5</b></p> <p style="text-align: right;">Cat. No.: HY-141067</p>	<p><b>N-methyl-N'-methyl-O-(m-PEG4)-O'-(propargyl-PEG4)-Cy3</b></p> <p style="text-align: right;">Cat. No.: HY-141032</p>
<p>N-methyl-N'-methyl-O-(m-PEG4)-O'-(azide-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>N-methyl-N'-methyl-O-(m-PEG4)-O'-(propargyl-PEG4)-Cy3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N-methyl-N'-methyl-O-(m-PEG4)-O'-(propargyl-PEG4)-Cy5</b> Cat. No.: HY-141052</p>	<p><b>N-Methyl-N-(t-Boc)-PEG4-acid</b> Cat. No.: HY-140510</p>
<p>N-methyl-N'-methyl-O-(m-PEG4)-O'-(propargyl-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N-Methyl-N-(t-Boc)-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-PEG3-N'-(azide-PEG3)-Cy5</b> Cat. No.: HY-141061</p>	<p><b>N-PEG3-N'-(propargyl-PEG4)-Cy5</b> Cat. No.: HY-141050</p>
<p>N-PEG3-N'-(azide-PEG3)-Cy5 (chloride) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-PEG3-N'-(propargyl-PEG4)-Cy5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N-Succinimidyl 3-(Bromoacetamido)propionate</b> (3-(2-Bromoacetamido)propanoic acid NHS ester) Cat. No.: HY-141385</p>	<p><b>N-tert-Butoxycarbonyl-trans-4-hydroxy-D-proline</b> Cat. No.: HY-77593</p>
<p>N-Succinimidyl 3-(Bromoacetamido)propionate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. N-Succinimidyl 3-(Bromoacetamido)propionate is also a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N-tert-Butoxycarbonyl-trans-4-hydroxy-D-proline is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N-tert-Butoxycarbonyl-trans-4-hydroxy-D-proline is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 g, 5 mg</p>
<p><b>N3-PEG12-Hydrazide</b> Cat. No.: HY-130886</p>	<p><b>N3-PEG16-Hydrazide</b> Cat. No.: HY-133320</p>
<p>N3-PEG12-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N3-PEG16-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N3-PEG24-Hydrazide</b> Cat. No.: HY-133321</p>	<p><b>N3-PEG3-CH2CH2-Boc</b> Cat. No.: HY-42489</p>
<p>N3-PEG24-Hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>N3-PEG3-CH2CH2-Boc is a cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). N3-PEG3-CH2CH2-Boc is also a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>N3-PEG3-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-42490</p>	<p><b>N3-PEG3-CH2COOH</b> (PROTAC Linker 14)</p> <p style="text-align: right;">Cat. No.: HY-42637</p>
<p>N3-PEG3-CH2CH2COOH a PEG-based PROTAC linker can be used in the synthesis of BI-3663 (HY-111546), BI-4216 and BI-0319. Azido-PEG3-acid is also a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg</p>	<p>N3-PEG3-CH2COOH (PROTAC Linker 14) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N3-PEG3-Propanehydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-130883</p>	<p><b>N3-PEG4-C2-NH2</b> (PROTAC Linker 20)</p> <p style="text-align: right;">Cat. No.: HY-128834</p>
<p>N3-PEG3-Propanehydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N3-PEG4-C2-NH2 (PROTAC Linker 20) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg</p>
<p><b>N3-PEG6-Propanehydrazide</b></p> <p style="text-align: right;">Cat. No.: HY-130885</p>	<p><b>N3-PEG8-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130228</p>
<p>N3-PEG6-Propanehydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>N3-PEG8-CH2COOH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>N33-TEG-COOH</b> (N3-TEG-COOH; 14-Azido-3,6,9,12-tetraoxatetradecanoic acid)</p> <p style="text-align: right;">Cat. No.: HY-108370</p>	<p><b>NH-bis(C1-PEG1-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140252</p>
<p>N33-TEG-COOH (N3-TEG-COOH) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 98.08% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 100 mg, 250 mg</p>	<p>NH-bis(C1-PEG1-Boc) is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH-bis(C2-PEG1-azide)</b></p> <p style="text-align: right;">Cat. No.: HY-140550</p>	<p><b>NH-bis(C2-PEG2-NH-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-130531</p>
<p>NH-bis(C2-PEG1-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH-bis(C2-PEG2-NH-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>











<p><b>NH-bis(m-PEG4)</b></p> <p style="text-align: right;">Cat. No.: HY-140263</p> <p>NH-bis(m-PEG4) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>NH-bis(m-PEG8)</b></p> <p style="text-align: right;">Cat. No.: HY-140264</p> <p>NH-bis(m-PEG8) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH-bis(PEG2-C2-acid)</b></p> <p style="text-align: right;">Cat. No.: HY-140514</p> <p>NH-bis(PEG2-C2-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>NH-bis(PEG2-C2-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140256</p> <p>NH-bis(PEG2-C2-Boc) is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH-bis(PEG2-propargyl)</b></p> <p style="text-align: right;">Cat. No.: HY-140088</p> <p>NH-bis(PEG2-propargyl) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>NH-bis(PEG3-acid)</b></p> <p style="text-align: right;">Cat. No.: HY-140515</p> <p>NH-bis(PEG3-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH-bis(PEG3-azide)</b></p> <p style="text-align: right;">Cat. No.: HY-140551</p> <p>NH-bis(PEG3-azide) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>NH-bis(PEG3-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140257</p> <p>NH-bis(PEG3-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 250 mg, 500 mg</p>
<p><b>NH-bis(PEG3-C2-NH-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140260</p> <p>NH-bis(PEG3-C2-NH-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>NH-bis(PEG4-acid)</b></p> <p style="text-align: right;">Cat. No.: HY-140516</p> <p>NH-bis(PEG4-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

<p><b>NH-bis(PEG4-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140258</p> <p>NH-bis(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>NH-bis(PEG4-C2-NH-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140261</p> <p>NH-bis(PEG4-C2-NH-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH-bis-PEG2</b></p> <p style="text-align: right;">Cat. No.: HY-130328</p> <p>NH-bis-PEG2 is a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). NH-bis-PEG2 is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>NH-bis-PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-140547</p> <p>NH-bis-PEG3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH-bis-PEG4</b></p> <p style="text-align: right;">Cat. No.: HY-140548</p> <p>NH-bis-PEG4 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>NH-bis-PEG5</b></p> <p style="text-align: right;">Cat. No.: HY-140549</p> <p>NH-bis-PEG5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH2-C2-amido-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130712</p> <p>NH2-C2-amido-C2-Boc is a PROTAC linker, which refers to the alkyl/ether composition. NH2-C5-NH-Boc can be used in the synthesis of a series of PROTACs, such as the PROTAC CDK2/9 Degrader-1 (HY-130709).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>NH2-C2-NH-Boc (PROTAC Linker 22)</b></p> <p style="text-align: right;">Cat. No.: HY-40171</p> <p>NH2-C2-NH-Boc (PROTAC Linker 22) is a alkyl chain-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.52%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>NH2-C4-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-40178</p> <p>NH2-C4-NH-Boc (compound 15) is a PROTAC linker, which refers to the Alkyl/ether composition. NH2-C4-NH-Boc can be used in the synthesis of a series of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>NH2-C5-NH-Boc (PROTAC Linker 23)</b></p> <p style="text-align: right;">Cat. No.: HY-W004710</p> <p>NH2-C5-NH-Boc (PROTAC Linker 23) is an alkyl chain-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.27%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>










<p><b>NH2-C6-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-W008296</p>	<p><b>NH2-O-C5-COOH hydrobromide</b> (6-Aminoxy-hexanoic acid hydrobromide)</p> <p style="text-align: right;">Cat. No.: HY-133411A</p>
<p>NH2-C6-NH-Boc is a <b>PROTAC linker</b> which refers to the alkyl/ether composition. NH2-C6-NH-Boc can be used in the synthesis the Mcl-1 inhibitor based on PROTAC.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 99.60% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>	<p>NH2-O-C5-COOH (hydrobromide) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH2-PEG-Strt (MW 10000)</b></p> <p style="text-align: right;">Cat. No.: HY-138307</p>	<p><b>NH2-PEG-Strt (MW 3000)</b></p> <p style="text-align: right;">Cat. No.: HY-138308</p>
<p>NH2-PEG-Strt (MW 10000) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG-Strt (MW 3000) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH2-PEG1-C1-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-W073709</p>	<p><b>NH2-PEG1-CH2CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-W067489</p>
<p>NH2-PEG1-C1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG1-CH2CH2-Boc is a PEG- and Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>NH2-PEG10-C2-dimethylamino</b></p> <p style="text-align: right;">Cat. No.: HY-138390</p>	<p><b>NH2-PEG2-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-42149</p>
<p>NH2-PEG10-C2-dimethylamino is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG2-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. NH2-PEG2-C2-Boc is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>NH2-PEG2-C6-Cl</b></p> <p style="text-align: right;">Cat. No.: HY-W096119</p>	<p><b>NH2-PEG2-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-42427</p>
<p>NH2-PEG2-C6-Cl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 50 mg, 100 mg</p>	<p>NH2-PEG2-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 250 mg</p>







<p><b>NH2-PEG2-methyl acetate</b></p> <p style="text-align: right;">Cat. No.: HY-W096162</p>	<p><b>NH2-PEG3</b> (PROTAC Linker 35)</p> <p style="text-align: right;">Cat. No.: HY-W007545</p>
<p>NH2-PEG2-methyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG3 (PROTAC Linker 35) is a PROTAC linker, which belongs to a polyethylene glycol (PEG) linker. NH2-PEG3 (PROTAC Linker 35) can be used in the synthesis of the PROTAC (<math>\beta</math>-NF-JQ1).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 97.58% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>NH2-PEG3 hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-W096143</p>	<p><b>NH2-PEG3-C1-Boc</b> (PROTAC Linker 5)</p> <p style="text-align: right;">Cat. No.: HY-128801</p>
<p>NH2-PEG3 hydrochloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG3-C1-Boc (PROTAC Linker 5) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> <math>\geq</math>98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>NH2-PEG3-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-135804</p>	<p><b>NH2-PEG3-C2-NH-Boc</b> (PROTAC Linker 15)</p> <p style="text-align: right;">Cat. No.: HY-42776</p>
<p>NH2-PEG3-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 250 mg, 500 mg</p>	<p>NH2-PEG3-C2-NH-Boc (PROTAC Linker 15) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> <math>\geq</math>97.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH2-PEG3-C6-Cl</b></p> <p style="text-align: right;">Cat. No.: HY-138469</p>	<p><b>NH2-PEG3500-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-138523</p>
<p>NH2-PEG3-C6-Cl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG3500-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH2-PEG4-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W021787</p>	<p><b>NH2-PEG4-Glu(OH)-NH-m-PEG24</b></p> <p style="text-align: right;">Cat. No.: HY-140241</p>
<p>NH2-PEG4-CH2CH2COOH is a cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). NH2-PEG4-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> <math>\geq</math>98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 500 mg</p>	<p>NH2-PEG4-Glu(OH)-NH-m-PEG24 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>













<p><b>NH2-PEG5-C1-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-135938</p>	<p><b>NH2-PEG5-C2-NH-Boc</b> (PROTAC Linker 17)</p> <p style="text-align: right;">Cat. No.: HY-W022240</p>
<p>NH2-PEG5-C1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG5-C2-NH-Boc (PROTAC Linker 17) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>NH2-PEG5-C6-Cl</b></p> <p style="text-align: right;">Cat. No.: HY-129622</p>	<p><b>NH2-PEG5-C6-Cl hydrochloride</b></p> <p style="text-align: right;">Cat. No.: HY-129622A</p>
<p>NH2-PEG5-C6-Cl (K-7) is a linker which refers to the PEG composition. NH2-PEG5-C6-Cl can be used in the synthesis of a series of compounds that induce degradation of intracellular molecules by autophagy.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG5-C6-Cl hydrochloride is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NH2-PEG5-OH</b></p> <p style="text-align: right;">Cat. No.: HY-129637</p>	<p><b>NH2-PEG6-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130486</p>
<p>NH2-PEG5-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. NH2-PEG5-OH is also a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg, 500 mg</p>	<p>NH2-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. NH2-PEG6-Boc is also a non-cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mg, 50 mg, 100 mg</p>
<p><b>NH2-PEG6-C1-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-135918</p>	<p><b>NH2-PEG6-CH2CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W040257</p>
<p>NH2-PEG6-C1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG6-CH2CH2COOH is a cleavable 6 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). NH2-PEG6-CH2CH2COOH is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 100 mg</p>
<p><b>NH2-PEG7</b></p> <p style="text-align: right;">Cat. No.: HY-120918</p>	<p><b>NH2-PEG8-C1-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-135943</p>
<p>NH2-PEG7 is a PROTAC linker, which refers to the PEG composition. NH2-PEG7 can be used in the synthesis of the PROTAC PARP1 degrader iRucaparib-AP6.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 250 mg, 500 mg</p>	<p>NH2-PEG8-C1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

<p><b>NH2-PEG8-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130200</p>	<p><b>NH2-PEG9-acid</b></p> <p style="text-align: right;">Cat. No.: HY-W019798</p>
<p>NH2-PEG8-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>NH2-PEG9-acid is a non-cleavable 9 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). NH2-PEG9-acid also is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 500 mg</p>
<p><b>NH2-Ph-C4-acid-NH2-Me</b> (PROTAC Linker 31)</p> <p style="text-align: right;">Cat. No.: HY-128931</p>	<p><b>NHS ester-PEG10-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138510</p>
<p>NH2-Ph-C4-acid-NH2-Me (PROTAC Linker 31) is an alkyl chain-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>NHS ester-PEG10-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NHS ester-PEG13-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138509</p>	<p><b>NHS ester-PEG3-S-methyl ethanethioate</b></p> <p style="text-align: right;">Cat. No.: HY-132103</p>
<p>NHS ester-PEG13-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>NHS ester-PEG3-S-methyl ethanethioate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NHS ester-PEG7-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-138511</p>	<p><b>NHS-bis-PEG2-amide-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-138522</p>
<p>NHS ester-PEG7-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>NHS-bis-PEG2-amide-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>NHS-PEG4-(m-PEG12)3-ester</b></p> <p style="text-align: right;">Cat. No.: HY-141125</p>	<p><b>NHS-PEG4-(m-PEG4)3-ester</b></p> <p style="text-align: right;">Cat. No.: HY-141124</p>
<p>NHS-PEG4-(m-PEG12)3-ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>NHS-PEG4-(m-PEG4)3-ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>









<p><b>Nonaethylene glycol</b></p> <p style="text-align: right;">Cat. No.: HY-116899</p>	<p><b>Nonaethylene glycol monomethyl ether</b></p> <p style="text-align: right;">Cat. No.: HY-120175</p>
<p>Nonaethylene glycol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Nonaethylene glycol monomethyl ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Nonylbenzene-PEG5-OH</b></p> <p style="text-align: right;">Cat. No.: HY-W096105</p>	<p><b>Nonylbenzene-PEG8-OH</b></p> <p style="text-align: right;">Cat. No.: HY-W096099</p>
<p>Nonylbenzene-PEG5-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Nonylbenzene-PEG8-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Octaethylene glycol</b></p> <p style="text-align: right;">Cat. No.: HY-W050087</p>	<p><b>Octaethylene glycol monomethyl ether</b></p> <p style="text-align: right;">Cat. No.: HY-W042657</p>
<p>Octaethylene glycol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p>Octaethylene glycol monomethyl ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Oleic-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-141293</p>	<p><b>OPSS-PEG8-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130348</p>
<p>Oleic-DBCO is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>OPSS-PEG8-COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>OTs-C6-OBn</b></p> <p style="text-align: right;">Cat. No.: HY-130621</p>	<p><b>PC Azido-PEG11-NHS carbonate ester</b></p> <p style="text-align: right;">Cat. No.: HY-140142</p>
<p>OTs-C6-OBn is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTAC SGK3 degrader-1 (HY-125878).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> 98.07%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 500 mg</p>	<p>PC Azido-PEG11-NHS carbonate ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>PC DBCO-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-138756</p> <p>PC DBCO-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>PC Methyltetrazine-PEG4-NHS carbonate ester</b></p> <p style="text-align: right;">Cat. No.: HY-140141</p> <p>PC Methyltetrazine-PEG4-NHS carbonate ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>PC-Biotin-PEG4-NHS carbonate</b></p> <p style="text-align: right;">Cat. No.: HY-140135</p> <p>PC-Biotin-PEG4-NHS carbonate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>PC-PEG11-Azide</b></p> <p style="text-align: right;">Cat. No.: HY-140863</p> <p>PC-PEG11-Azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>PEG12-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-117050</p> <p>Tos-PEG12 is a noncleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). PEG12-Tos is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>PEG2-bis(phosphonic acid diethyl ester)</b></p> <p style="text-align: right;">Cat. No.: HY-141320</p> <p>PEG2-bis(phosphonic acid diethyl ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>PEG2-Cl</b></p> <p style="text-align: right;">Cat. No.: HY-W096116</p> <p>PEG2-Cl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>PEG2-ethyl acetate</b></p> <p style="text-align: right;">Cat. No.: HY-138436</p> <p>PEG2-ethyl acetate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>PEG20-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140358</p> <p>PEG20-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-W096121</p> <p>PEG3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>PEG3-bis(phosphonic acid diethyl ester)</b></p> <p>Cat. No.: HY-141321</p>	<p><b>PEG3-bis(phosphonic acid)</b></p> <p>Cat. No.: HY-141294</p>
<p>PEG3-bis(phosphonic acid diethyl ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>PEG3-bis(phosphonic acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>PEG3-bis-(ethyl phosphonate)</b></p> <p>Cat. No.: HY-141295</p>	<p><b>PEG3-C4-OBn</b></p> <p>Cat. No.: HY-130620</p>
<p>PEG3-bis-(ethyl phosphonate) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>PEG3-C4-OBn is a polyethylene glycol (PEG)-based PROTAC linker. PEG3-C4-OBn can be used in the synthesis of the PROTAC SGK3 degrader-1 (HY-125878). PROTAC SGK3 degrader-1 is a potent SGK3 degrader based on PROTAC.</p>  <p><b>Purity:</b> 99.29% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 500 mg</p>
<p><b>PEG3-methylamine</b></p> <p>Cat. No.: HY-140161</p>	<p><b>PEG3-O-CH2COOH</b> (PROTAC Linker 8)</p> <p>Cat. No.: HY-128804</p>
<p>PEG3-methylamine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>PEG3-O-CH2COOH (PROTAC Linker 8) is a PEG-based PROTAC linker can be used in the synthesis of SNIPERS.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>PEG4-bis(phosphonic acid diethyl ester)</b></p> <p>Cat. No.: HY-141322</p>	<p><b>PEG4-Ms</b></p> <p>Cat. No.: HY-140388</p>
<p>PEG4-bis(phosphonic acid diethyl ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>PEG4-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>PEG4-sulfonic acid</b></p> <p>Cat. No.: HY-133054</p>	<p><b>PEG5-bis-(Ethyl phosphonate)</b></p> <p>Cat. No.: HY-141296</p>
<p>PEG4-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>PEG5-bis-(Ethyl phosphonate) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>








<p><b>PEG5-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-23417</p>	<p><b>PEG6-(CH<sub>2</sub>CO<sub>2</sub>H)<sub>2</sub></b></p> <p style="text-align: right;">Cat. No.: HY-122702</p>
<p>PEG5-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.05%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 10 mg, 50 mg, 100 mg</p>	<p>PEG6-(CH<sub>2</sub>CO<sub>2</sub>H)<sub>2</sub> is a symmetric PEG PROTAC linker, for the synthesis of Homo-PROTACs which is bivalent small-molecule dimerizers of the VHL E3 ubiquitin ligase to induce self-degradation.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>
<p><b>PEG7-O-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-140389</p>	<p><b>PEG8-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-130555</p>
<p>PEG7-O-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>PEG8-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Pentaethylene glycol</b></p> <p style="text-align: right;">Cat. No.: HY-W007341</p>	<p><b>Pentaethylene glycol di(p-toluenesulfonate) (Penta(ethylene glycol) bis(p-toluenesulfonate); Bis-Tos-PEG5)</b></p> <p style="text-align: right;">Cat. No.: HY-W004447</p>
<p>Pentaethylene glycol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>	<p>Pentaethylene glycol di(p-toluenesulfonate) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.34%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Pentaethylene glycol monomethyl ether</b></p> <p style="text-align: right;">Cat. No.: HY-W042713</p>	<p><b>Ph-Bis(C1-N-(C2-NH-Boc)2)</b></p> <p style="text-align: right;">Cat. No.: HY-140337</p>
<p>Pentaethylene glycol monomethyl ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Ph-Bis(C1-N-(C2-NH-Boc)2) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Ph-PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-W096158</p>	<p><b>Phenol-amido-C1-PEG3-N3</b> (PROTAC Linker 21)</p> <p style="text-align: right;">Cat. No.: HY-128835</p>
<p>Ph-PEG3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Phenol-amido-C1-PEG3-N3 (PROTAC Linker 21) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Phthalamide-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140862</p>	<p><b>Phthalamide-PEG3-C2-OTs</b></p> <p style="text-align: right;">Cat. No.: HY-129772</p>
<p>Phthalamide-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Phthalamide-PEG3-C2-OTs (Compound 5) is a PROTAC linker, which refers to the PEGs composition. Phthalamide-PEG3-C2-OTs can be used in the synthesis of a series of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Phthalamide-PEG4-MPDM-OH</b></p> <p style="text-align: right;">Cat. No.: HY-129774</p>	<p><b>Phthalamide-PEG4-PDM-OTBS</b></p> <p style="text-align: right;">Cat. No.: HY-129773</p>
<p>Phthalamide-PEG4-MPDM-OH is a PROTAC linker, which refers to the PEGs composition. Phthalamide-PEG4-MPDM-OH can be used in the synthesis of a series of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Phthalamide-PEG4-PDM-OTBS is a PROTAC linker, which refers to the PEGs composition. Phthalamide-PEG4-PDM-OTBS can be used in the synthesis of a series of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Pip-alkyne-Ph-COOCH3</b></p> <p style="text-align: right;">Cat. No.: HY-130846</p>	<p><b>Propanol-PEG3-CH2OH</b></p> <p style="text-align: right;">Cat. No.: HY-134689</p>
<p>Pip-alkyne-Ph-COOCH3 is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTAC ARD-266.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propanol-PEG3-CH2OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propanol-PEG4-CH2OH</b></p> <p style="text-align: right;">Cat. No.: HY-134690</p>	<p><b>Propanol-PEG5-CH2OH</b></p> <p style="text-align: right;">Cat. No.: HY-134678</p>
<p>Propanol-PEG4-CH2OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propanol-PEG5-CH2OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propanol-PEG6-CH2OH</b></p> <p style="text-align: right;">Cat. No.: HY-134691</p>	<p><b>Propargyl-PEG-acid</b></p> <p style="text-align: right;">Cat. No.: HY-143820</p>
<p>Propanol-PEG6-CH2OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Propargyl-PEG-amine</b></p> <p>Cat. No.: HY-143819</p>	<p><b>Propargyl-PEG1-acid</b></p> <p>Cat. No.: HY-130504</p>
<p>Propargyl-PEG-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG1-acid is a PEG-based PROTAC linker can be used in the synthesis of BTK-CRBN PROTACs Ibrutinib(HY-10997)-based PROTAC 4 and PROTAC 5. PROTAC 5 causes the degradation of BTK and induces the degradation of CSK, LYN, and LAT2 at 10 μM.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG1-acrylate</b></p> <p>Cat. No.: HY-140064</p>	<p><b>Propargyl-PEG1-Boc</b></p> <p>Cat. No.: HY-140026</p>
<p>Propargyl-PEG1-acrylate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG1-Boc is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG1-NH2</b></p> <p>Cat. No.: HY-116069</p>	<p><b>Propargyl-PEG1-SS-PEG1-C2-Boc</b></p> <p>Cat. No.: HY-130690</p>
<p>Propargyl-PEG1-NH2 is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG1-SS-PEG1-C2-Boc is a Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG1-SS-PEG1-C2-Boc is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG1-THP</b></p> <p>Cat. No.: HY-138334</p>	<p><b>Propargyl-PEG10-acid</b></p> <p>Cat. No.: HY-140020</p>
<p>Propargyl-PEG1-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG10-alcohol</b></p> <p>Cat. No.: HY-138460</p>	<p><b>Propargyl-PEG10-amine</b></p> <p>Cat. No.: HY-133230</p>
<p>Propargyl-PEG10-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG10-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>





<p><b>Propargyl-PEG10-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140029</p> <p>Propargyl-PEG10-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG11-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138763</p> <p>Propargyl-PEG11-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG11-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-138366</p> <p>Propargyl-PEG11-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG11-amine</b></p> <p style="text-align: right;">Cat. No.: HY-138766</p> <p>Propargyl-PEG11-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG11-methane</b></p> <p style="text-align: right;">Cat. No.: HY-140061</p> <p>Propargyl-PEG11-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138764</p> <p>Propargyl-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG12-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140034</p> <p>Propargyl-PEG12-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG12-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-135823</p> <p>Propargyl-PEG12-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG12-methane</b></p> <p style="text-align: right;">Cat. No.: HY-130894</p> <p>Propargyl-PEG12-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG12-OH</b></p> <p style="text-align: right;">Cat. No.: HY-117045</p> <p>Propargyl-PEG12-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>










<p><b>Propargyl-PEG12-SH</b></p> <p style="text-align: right;">Cat. No.: HY-130911</p> <p>Propargyl-PEG12-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG13-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140021</p> <p>Propargyl-PEG13-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG13-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138733</p> <p>Propargyl-PEG13-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG13-OH</b></p> <p style="text-align: right;">Cat. No.: HY-138735</p> <p>Propargyl-PEG13-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG14-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140022</p> <p>Propargyl-PEG14-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG14-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140030</p> <p>Propargyl-PEG14-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG17-methane</b></p> <p style="text-align: right;">Cat. No.: HY-140062</p> <p>Propargyl-PEG17-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-118764</p> <p>Propargyl-PEG2-acid is a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG2-acid is also a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>
<p><b>Propargyl-PEG2-amine</b></p> <p style="text-align: right;">Cat. No.: HY-W051634</p> <p>Propargyl-PEG2-amine is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG2-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>	<p><b>Propargyl-PEG2-beta-D-glucose</b></p> <p style="text-align: right;">Cat. No.: HY-141131</p> <p>Propargyl-PEG2-beta-D-glucose is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>


<p><b>Propargyl-PEG2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130597</p> <p>Propargyl-PEG2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG2-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-W096164</p> <p>Propargyl-PEG2-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-41922</p> <p>Propargyl-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>	<p><b>Propargyl-PEG2-Ms</b></p> <p style="text-align: right;">Cat. No.: HY-130584</p> <p>Propargyl-PEG2-Ms is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>
<p><b>Propargyl-PEG2-N-bis(PEG2)</b></p> <p style="text-align: right;">Cat. No.: HY-140083</p> <p>Propargyl-PEG2-N-bis(PEG2) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG2-NHBoc</b></p> <p style="text-align: right;">Cat. No.: HY-118808</p> <p>Propargyl-PEG2-NHBoc is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG2-NHBoc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG2-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-138734</p> <p>Propargyl-PEG2-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG2-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130541</p> <p>Propargyl-PEG2-OH is a PEG-based PROTAC linker can be used in the synthesis of Thalidomide-O-PEG2-propargyl (HY-126458).</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Propargyl-PEG2-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140374</p> <p>Propargyl-PEG2-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG2-urea-C3-triethoxysilane</b></p> <p style="text-align: right;">Cat. No.: HY-134749</p> <p>Propargyl-PEG2-urea-C3-triethoxysilane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Propargyl-PEG24-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140035</p>	<p><b>Propargyl-PEG25-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130909</p>
<p>Propargyl-PEG24-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b>  <b>Size:</b> 50 mg, 100 mg</p>	<p>Propargyl-PEG25-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG3-1-o-(b-cyanoethyl-N,N-diisopropyl)phosphoramidite</b></p> <p style="text-align: right;">Cat. No.: HY-W096141</p>	<p><b>Propargyl-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-126975</p>
<p>Propargyl-PEG3-1-o-b-cyanoethyl-N,N-diisopropylphosphoramidite is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG3-acid is a non-cleavable (3 unit PEG) ADC linker and also a PEG-based PROTAC linker that can be used to synthesis 6-OHDA-PEG3-yne. 6-OHDA-PEG3-yne contains 6-OHDA (HY-B1081, HY-B1081A) and Propargyl-PEG3-acid.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Propargyl-PEG3-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-41921</p>	<p><b>Propargyl-PEG3-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140033</p>
<p>Propargyl-PEG3-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 96.83%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>	<p>Propargyl-PEG3-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-118781</p>	<p><b>Propargyl-PEG3-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140027</p>
<p>Propargyl-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG3-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG3-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-140036</p>	<p><b>Propargyl-PEG3-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130563</p>
<p>Propargyl-PEG3-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG3-CH2COOH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg</p>




<p><b>Propargyl-PEG3-methane</b></p> <p style="text-align: right;">Cat. No.: HY-140059</p> <p>Propargyl-PEG3-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG3-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140876</p> <p>Propargyl-PEG3-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG3-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-126974</p> <p>Propargyl-PEG3-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG3-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG3-OCH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140031</p> <p>Propargyl-PEG3-OCH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG3-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-132007</p> <p>Propargyl-PEG3-PFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG3-phosphonic acid</b></p> <p style="text-align: right;">Cat. No.: HY-133049</p> <p>Propargyl-PEG3-phosphonic acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG3-phosphonic acid diethyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-130146</p> <p>Propargyl-PEG3-phosphonic acid diethyl ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG3-SH</b></p> <p style="text-align: right;">Cat. No.: HY-130900</p> <p>Propargyl-PEG3-SH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-peg3-sulfone-peg3-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-140611</p> <p>Propargyl-peg3-sulfone-peg3-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG4-5-nitrophenyl carbonate</b></p> <p style="text-align: right;">Cat. No.: HY-140048</p> <p>Propargyl-PEG4-5-nitrophenyl carbonate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Propargyl-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130481</p> <p>Propargyl-PEG4-acid is a PEG-based PROTAC linker can be used in the synthesis of BTK-IAP PROTACs Ibrutinib (HY-10997)-based PROTAC 2 and an analogue PROTAC 3. PROTAC 3 causes BTK degradation with a DC<sub>50</sub> of 200 nM in THP-1 cells.</p>  <p><b>Purity:</b> ≥97.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>	<p><b>Propargyl-PEG4-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-133229</p> <p>Propargyl-PEG4-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 500 mg</p>
<p><b>Propargyl-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-114670</p> <p>Propargyl-PEG4-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p><b>Propargyl-PEG4-beta-D-glucose</b></p> <p style="text-align: right;">Cat. No.: HY-141132</p> <p>Propargyl-PEG4-beta-D-glucose is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140028</p> <p>Propargyl-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG4-Br</b></p> <p style="text-align: right;">Cat. No.: HY-130591</p> <p>Propargyl-PEG4-Br is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG4-Br is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>
<p><b>Propargyl-PEG4-CH2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140024</p> <p>Propargyl-PEG4-CH2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG4-CH2-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140063</p> <p>Propargyl-PEG4-CH2-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG4-CH2CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130293</p> <p>Propargyl-PEG4-CH2CH2-Boc is a non-cleavable ADC linker that can be used to synthesize ADC inhibitors of Galectin-3. Propargyl-PEG4-CH2CH2-Boc is a PEG- and Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG4-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-140023</p> <p>Propargyl-PEG4-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>


<p><b>Propargyl-PEG4-methylamine</b></p> <p>Cat. No.: HY-140045</p>	<p><b>Propargyl-PEG4-O-C1-Boc</b></p> <p>Cat. No.: HY-130389</p>
<p>Propargyl-PEG4-methylamine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG4-O-C1-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG4-O-C1-NHS ester</b></p> <p>Cat. No.: HY-130390</p>	<p><b>Propargyl-PEG4-S-PEG4-acid</b></p> <p>Cat. No.: HY-140597</p>
<p>Propargyl-PEG4-O-C1-NHS ester (compound 8) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG4-S-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG4-S-PEG4-Boc</b></p> <p>Cat. No.: HY-140599</p>	<p><b>Propargyl-PEG4-S-PEG4-propargyl</b></p> <p>Cat. No.: HY-140598</p>
<p>Propargyl-PEG4-S-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG4-S-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG4-Sulfone-PEG4-acid</b></p> <p>Cat. No.: HY-140610</p>	<p><b>Propargyl-PEG4-Sulfone-PEG4-Boc</b></p> <p>Cat. No.: HY-140612</p>
<p>Propargyl-PEG4-Sulfone-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG4-Sulfone-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG4-sulfonic acid</b></p> <p>Cat. No.: HY-140047</p>	<p><b>Propargyl-PEG4-tetra-Ac-beta-D-galactose</b></p> <p>Cat. No.: HY-141135</p>
<p>Propargyl-PEG4-sulfonic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG4-tetra-Ac-beta-D-galactose is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Propargyl-PEG4-tetra-Ac-beta-D-glucose</b></p> <p>Cat. No.: HY-141133</p> <p>Propargyl-PEG4-tetra-Ac-beta-D-glucose is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG4-thiol</b></p> <p>Cat. No.: HY-116427</p> <p>Propargyl-PEG4-thiol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG4-thiol is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG4-Tos</b></p> <p>Cat. No.: HY-130387</p> <p>Propargyl-PEG4-Tos is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG4-Tos is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG5-1-o-(b-cyanoethyl-n,n-diisopropyl)phosphoramidite</b></p> <p>Cat. No.: HY-W096140</p> <p>Propargyl-PEG5-1-o-b-cyanoethyl-nn-diisopropylphosphoramidite is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG5-acid</b></p> <p>Cat. No.: HY-101157</p> <p>Propargyl-PEG5-acid is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG5-acid can be used to synthesize ADC inhibitors of Galectin-3. Propargyl-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mg, 25 mg, 50 mg, 100 mg</p>	<p><b>Propargyl-PEG5-amine</b></p> <p>Cat. No.: HY-126976</p> <p>Propargyl-PEG5-amine is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Propargyl-PEG5-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 25 mg, 50 mg, 100 mg</p>
<p><b>Propargyl-PEG5-azide</b></p> <p>Cat. No.: HY-138738</p> <p>Propargyl-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG5-Br</b></p> <p>Cat. No.: HY-138736</p> <p>Propargyl-PEG5-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG5-Ms</b></p> <p>Cat. No.: HY-140046</p> <p>Propargyl-PEG5-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Propargyl-PEG5-NHS ester</b></p> <p>Cat. No.: HY-130388</p> <p>Propargyl-PEG5-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG5-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 25 mg, 50 mg, 100 mg</p>











<p><b>Propargyl-PEG5-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130147</p>	<p><b>Propargyl-PEG5-PFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-132043</p>
<p>Propargyl-PEG5-OH is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG5-PFP ester is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130386</p>	<p><b>Propargyl-PEG6-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-130382</p>
<p>Propargyl-PEG6-acid is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs. Propargyl-PEG6-acid is a cleavable <b>ADC linker</b> used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG6-alcohol is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG6-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130384</p>	<p><b>Propargyl-PEG6-Br</b></p> <p style="text-align: right;">Cat. No.: HY-138321</p>
<p>Propargyl-PEG6-Boc is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG6-Br is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG6-N3</b></p> <p style="text-align: right;">Cat. No.: HY-130183</p>	<p><b>Propargyl-PEG6-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-130180</p>
<p>Propargyl-PEG6-N3 is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG6-NH2 is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 250 mg</p>
<p><b>Propargyl-PEG6-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130385</p>	<p><b>Propargyl-PEG6-SH</b></p> <p style="text-align: right;">Cat. No.: HY-135917</p>
<p>Propargyl-PEG6-NHS ester is a PEG/Alkyl/ether-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs. Propargyl-PEG6-NHS ester is a cleavable <b>ADC linker</b> used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG6-SH is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>











<p><b>Propargyl-PEG7-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130383</p>	<p><b>Propargyl-PEG7-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-130378</p>
<p>Propargyl-PEG7-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG7-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG7-alcohol is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG7-amine</b></p> <p style="text-align: right;">Cat. No.: HY-138765</p>	<p><b>Propargyl-PEG7-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130380</p>
<p>Propargyl-PEG7-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG7-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG7-Br</b></p> <p style="text-align: right;">Cat. No.: HY-138737</p>	<p><b>Propargyl-PEG7-methane</b></p> <p style="text-align: right;">Cat. No.: HY-140060</p>
<p>Propargyl-PEG7-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG7-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG7-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130381</p>	<p><b>Propargyl-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130379</p>
<p>Propargyl-PEG7-NHS ester is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG7-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG8-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG8-acid is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). The ADCs can be used in bacterial infections caused by Gram-negative bacteria.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG8-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130375</p>	<p><b>Propargyl-PEG8-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-130377</p>
<p>Propargyl-PEG7-Boc is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG8-bromide is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Propargyl-PEG8-bromide is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>








<p><b>Propargyl-PEG8-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-130182</p>	<p><b>Propargyl-PEG8-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130376</p>
<p>Propargyl-PEG8-NH2 (compound 3b) is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs. Propargyl-PEG8-NH2 is a non-cleavable <b>ADC linker</b> used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG8-NHS ester is a PEG/Alkyl/ether-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs. Propargyl-PEG8-NHS ester is a cleavable <b>ADC linker</b> used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG8-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130374</p>	<p><b>Propargyl-PEG8-SH</b></p> <p style="text-align: right;">Cat. No.: HY-130910</p>
<p>Propargyl-PEG8-OH is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG8-SH is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG9-acid</b></p> <p style="text-align: right;">Cat. No.: HY-132091</p>	<p><b>Propargyl-PEG9-amine</b></p> <p style="text-align: right;">Cat. No.: HY-130373</p>
<p>Propargyl-PEG9-acid is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG9-amine is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG9-bromide</b></p> <p style="text-align: right;">Cat. No.: HY-130372</p>	<p><b>Propargyl-PEG9-OH</b></p> <p style="text-align: right;">Cat. No.: HY-126977</p>
<p>Propargyl-PEG9-bromide is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs. Propargyl-PEG9-bromide is a non-cleavable <b>ADC linker</b> used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propargyl-PEG9-OH is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Propargyl-PEG9-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138459</p>	<p><b>Propenyl-PEG3-Propenyl</b> (Triethylene glycol diallyl ether)</p> <p style="text-align: right;">Cat. No.: HY-134676</p>
<p>Propargyl-PEG9-THP is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Propenyl-PEG3-Propenyl is a PEG-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Propynol Ethoxylate</b></p> <p>Cat. No.: HY-43614</p>	<p><b>Propynyl-PEG1-Ac</b></p> <p>Cat. No.: HY-W096146</p>
<p>Propynol Ethoxylate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg, 1 g</p>	<p>Propynyl-PEG1-Ac is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Pyrene azide 3</b></p> <p>Cat. No.: HY-D1295</p>	<p><b>Pyrene-amido-PEG4-azide</b></p> <p>Cat. No.: HY-123609</p>
<p>Pyrene azide 3 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Pyrene-amido-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Pyrene-amido-PEG4-CH2CH2COOH</b></p> <p>Cat. No.: HY-130397</p>	<p><b>Pyrene-PEG2-azide</b></p> <p>Cat. No.: HY-141092</p>
<p>Pyrene-amido-PEG4-CH2CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Pyrene-PEG2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Pyrene-PEG5-alcohol</b></p> <p>Cat. No.: HY-141093</p>	<p><b>Pyrene-PEG5-biotin</b></p> <p>Cat. No.: HY-141094</p>
<p>Pyrene-PEG5-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Pyrene-PEG5-biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Pyrene-PEG5-propargyl</b></p> <p>Cat. No.: HY-141095</p>	<p><b>Pyrroline-5-carboxylate</b></p> <p>Cat. No.: HY-138354</p>
<p>Pyrene-PEG5-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Pyrroline-5-carboxylate is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>rel-Biotin-PEG3-C3-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-140902</p>	<p><b>Rhodamine B PEG2-NH2</b></p> <p style="text-align: right;">Cat. No.: HY-138396</p>
<p>rel-Biotin-PEG3-C3-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Rhodamine B PEG2-NH2 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 25 mg, 50 mg</p>
<p><b>S-acetyl-PEG12-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141342</p>	<p><b>S-acetyl-PEG16-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141343</p>
<p>S-acetyl-PEG12-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>S-acetyl-PEG16-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>S-acetyl-PEG2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141345</p>	<p><b>S-acetyl-PEG20-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141344</p>
<p>S-acetyl-PEG2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>S-acetyl-PEG20-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>S-acetyl-PEG3-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-141340</p>	<p><b>S-Acetyl-PEG3-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140859</p>
<p>S-acetyl-PEG3-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>S-Acetyl-PEG3-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>S-acetyl-PEG3-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-141346</p>	<p><b>S-Acetyl-PEG3-C2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141349</p>
<p>S-acetyl-PEG3-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>S-Acetyl-PEG3-C2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>


<p><b>S-acetyl-PEG3-phosphonic acid ethyl ester</b></p> <p>Cat. No.: HY-141317</p>	<p><b>S-acetyl-PEG4-alcohol</b></p> <p>Cat. No.: HY-141341</p>
<p>S-acetyl-PEG3-phosphonic acid ethyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>S-acetyl-PEG4-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>S-acetyl-PEG4-amine</b></p> <p>Cat. No.: HY-138752</p>	<p><b>S-acetyl-PEG4-Boc</b></p> <p>Cat. No.: HY-141347</p>
<p>S-acetyl-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>S-acetyl-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>S-acetyl-PEG4-NHBoc</b></p> <p>Cat. No.: HY-138755</p>	<p><b>S-acetyl-PEG4-propargyl</b></p> <p>Cat. No.: HY-141350</p>
<p>S-acetyl-PEG4-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>S-acetyl-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>S-acetyl-PEG4-Thiol</b></p> <p>Cat. No.: HY-138758</p>	<p><b>S-acetyl-PEG5-alcohol</b></p> <p>Cat. No.: HY-138748</p>
<p>S-acetyl-PEG4-Thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>S-acetyl-PEG5-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>S-acetyl-PEG6</b></p> <p>Cat. No.: HY-W096147</p>	<p><b>S-acetyl-PEG6-Boc</b></p> <p>Cat. No.: HY-141348</p>
<p>S-acetyl-PEG6 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>S-acetyl-PEG6-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>

<p><b>S-acetyl-PEG6-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-133473</p>	<p><b>S-Acetyl-PEG8-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130209</p>
<p>S-acetyl-PEG6-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>S-Acetyl-PEG8-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>S-Bis-(PEG4-Boc)</b></p> <p style="text-align: right;">Cat. No.: HY-140600</p>	<p><b>SMPH Crosslinker</b></p> <p style="text-align: right;">Cat. No.: HY-124318</p>
<p>S-Bis-(PEG4-Boc) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>SMPH Crosslinker is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>SPDP-C6-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-124377</p>	<p><b>SPDP-PEG12-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141356</p>
<p>SPDP-C6-NHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.09%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg</p>	<p>SPDP-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>SPDP-PEG24-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141354</p>	<p><b>SPDP-PEG24-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141357</p>
<p>SPDP-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>SPDP-PEG24-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>SPDP-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141352</p>	<p><b>SPDP-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130195</p>
<p>SPDP-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>SPDP-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>SPDP-PEG5-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133384</p> <p>SPDP-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>SPDP-PEG6-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-126717</p> <p>SPDP-PEG6-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>SPDP-PEG7-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133385</p> <p>SPDP-PEG7-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>SPDP-PEG9-acid</b></p> <p style="text-align: right;">Cat. No.: HY-133386</p> <p>SPDP-PEG9-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Sulfo DBCO-amine</b></p> <p style="text-align: right;">Cat. No.: HY-127056</p> <p>Sulfo DBCO-amine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Sulfo DBCO-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140286</p> <p>Sulfo DBCO-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 50 mg, 100 mg</p>
<p><b>Sulfo DBCO-PEG4-Maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-140307</p> <p>Sulfo DBCO-PEG4-Maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Sulfo-NHS-Acetate</b></p> <p style="text-align: right;">Cat. No.: HY-140341</p> <p>Sulfo-NHS-Acetate is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Sulfone-Bis-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140601</p> <p>Sulfone-Bis-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>t-Boc-amido-PEG10-acid</b></p> <p style="text-align: right;">Cat. No.: HY-144076</p> <p>t-Boc-amido-PEG10-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>



<p><b>t-Boc-Aminoxy-PEG11-amine</b></p> <p style="text-align: right;">Cat. No.: HY-143825</p>	<p><b>t-Boc-Aminoxy-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140414</p>
<p>t-Boc-Aminoxy-PEG11-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>t-Boc-Aminoxy-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>t-Boc-Aminoxy-PEG12-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-140419</p>	<p><b>t-Boc-Aminoxy-PEG12-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140418</p>
<p>t-Boc-Aminoxy-PEG12-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>t-Boc-Aminoxy-PEG12-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>t-Boc-Aminoxy-PEG2-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140431</p>	<p><b>t-Boc-Aminoxy-PEG3-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-140422</p>
<p>t-Boc-Aminoxy-PEG2-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>t-Boc-Aminoxy-PEG3-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>t-Boc-Aminoxy-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-143821</p>	<p><b>t-Boc-Aminoxy-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-140417</p>
<p>t-Boc-Aminoxy-PEG4-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>t-Boc-Aminoxy-PEG4-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>t-Boc-Aminoxy-PEG4-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-W190966</p>	<p><b>t-Boc-Aminoxy-PEG5-azide</b></p> <p style="text-align: right;">Cat. No.: HY-140434</p>
<p>t-Boc-Aminoxy-PEG4-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>t-Boc-Aminoxy-PEG5-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>


<p><b>t-Boc-aminoxy-PEG6-propargyl</b></p> <p>Cat. No.: HY-140054</p>	<p><b>t-Boc-Aminoxy-PEG7-amine</b></p> <p>Cat. No.: HY-140429</p>
<p>t-Boc-aminoxy-PEG6-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>t-Boc-Aminoxy-PEG7-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>t-Boc-Aminoxy-PEG7-bromide</b></p> <p>Cat. No.: HY-140436</p>	<p><b>t-Boc-Aminoxy-PEG7-methane</b></p> <p>Cat. No.: HY-140437</p>
<p>t-Boc-Aminoxy-PEG7-bromide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>t-Boc-Aminoxy-PEG7-methane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>t-Boc-Aminoxy-PEG8-alcohol</b></p> <p>Cat. No.: HY-140424</p>	<p><b>t-Boc-Aminoxy-PEG8-Ms</b></p> <p>Cat. No.: HY-140378</p>
<p>t-Boc-Aminoxy-PEG8-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>t-Boc-Aminoxy-PEG8-Ms is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>t-Boc-N-amido-PEG10-Br</b></p> <p>Cat. No.: HY-143828</p>	<p><b>t-Boc-N-amido-PEG15-Br</b></p> <p>Cat. No.: HY-143829</p>
<p>t-Boc-N-amido-PEG10-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>t-Boc-N-amido-PEG15-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>t-Boc-N-amido-PEG2-C6-Cl</b></p> <p>Cat. No.: HY-143830</p>	<p><b>t-Boc-N-amido-PEG5-acetic acid</b></p> <p>Cat. No.: HY-W190962</p>
<p>t-Boc-N-amido-PEG2-C6-Cl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 5 mg, 10 mg, 50 mg, 100 mg</p>	<p>BocNH-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>

<p><b>t-Boc-N-amido-PEG6-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-W096165</p>	<p><b>t-Butyl acetate-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W096080</p>
<p>t-Boc-N-amido-PEG6-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>t-Butyl acetate-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>t-Butyl acetate-PEG3-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-W096082</p>	<p><b>TAMRA-Azide-PEG-biotin</b></p> <p style="text-align: right;">Cat. No.: HY-140947</p>
<p>t-Butyl acetate-PEG3-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TAMRA-Azide-PEG-biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TAMRA-PEG3-Azide</b></p> <p style="text-align: right;">Cat. No.: HY-123629</p>	<p><b>TAMRA-PEG3-biotin</b></p> <p style="text-align: right;">Cat. No.: HY-140946</p>
<p>TAMRA-PEG3-Azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TAMRA-PEG3-biotin is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TAMRA-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140509</p>	<p><b>TAMRA-PEG4-Alkyne</b></p> <p style="text-align: right;">Cat. No.: HY-120666</p>
<p>TAMRA-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TAMRA-PEG4-Alkyne is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TAMRA-PEG4-methyltetrazine</b></p> <p style="text-align: right;">Cat. No.: HY-141286</p>	<p><b>TAMRA-PEG4-tetrazine</b></p> <p style="text-align: right;">Cat. No.: HY-141285</p>
<p>TAMRA-PEG4-methyltetrazine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TAMRA-PEG4-tetrazine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

### tans-4-Hydroxy-D-proline hydrochloride

Cat. No.: HY-W003511

tans-4-Hydroxy-D-proline hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs). tans-4-Hydroxy-D-proline hydrochloride is also a alkyl chain-based PROTAC linker that can be used in the synthesis of PR.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 100 mg, 500 mg

### tans-4-Hydroxy-D-proline methyl ester hydrochloride

Cat. No.: HY-W006629

tans-4-Hydroxy-D-proline methyl ester hydrochloride is a non-cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 250 mg, 500 mg

### TBDMS-PEG4-OH

Cat. No.: HY-W042579

TBDMS-PEG4-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### TCO-amine

Cat. No.: HY-141176

TCO-amine is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### TCO-C3-PEG3-C3-amine

Cat. No.: HY-141182

TCO-C3-PEG3-C3-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### TCO-NHS ester

Cat. No.: HY-141165

TCO-NHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 50 mg, 100 mg

### TCO-OH

Cat. No.: HY-141186

TCO-OH is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 50 mg, 100 mg

### TCO-PEG-TCO (MW 5000)

Cat. No.: HY-140731

TCO-PEG-TCO (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### TCO-PEG11-TCO

Cat. No.: HY-133478

TCO-PEG11-TCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

### TCO-PEG12-acid

Cat. No.: HY-141162










TCO-PEG12-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.













**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

<p><b>TCO-PEG12-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141170</p>	<p><b>TCO-PEG12-TFP ester</b></p> <p style="text-align: right;">Cat. No.: HY-141174</p>
<p>TCO-PEG12-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. TCO-PEG12-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG12-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG2-amine</b></p> <p style="text-align: right;">Cat. No.: HY-141177</p>	<p><b>TCO-PEG2-Sulfo-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141172</p>
<p>TCO-PEG2-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG2-Sulfo-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg</p>
<p><b>TCO-PEG2-Sulfo-NHS ester sodium</b></p> <p style="text-align: right;">Cat. No.: HY-141172A</p>	<p><b>TCO-PEG24-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141163</p>
<p>TCO-PEG2-Sulfo-NHS ester sodium is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG24-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG24-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141171</p>	<p><b>TCO-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140015</p>
<p>TCO-PEG24-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG3-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg</p>
<p><b>TCO-PEG3-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-133476</p>	<p><b>TCO-PEG3-amide-C3-triethoxysilane</b></p> <p style="text-align: right;">Cat. No.: HY-141185</p>
<p>TCO-PEG3-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG3-amide-C3-triethoxysilane is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>








<p><b>TCO-PEG3-amine</b></p> <p style="text-align: right;">Cat. No.: HY-141178</p>	<p><b>TCO-PEG3-maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-133477</p>
<p>TCO-PEG3-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg, 25 mg, 50 mg, 100 mg</p>	<p>TCO-PEG3-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG3-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141166</p>	<p><b>TCO-PEG3-oxyamine</b></p> <p style="text-align: right;">Cat. No.: HY-133474</p>
<p>TCO-PEG3-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg</p>	<p>TCO-PEG3-oxyamine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG3-TCO</b></p> <p style="text-align: right;">Cat. No.: HY-141187</p>	<p><b>TCO-PEG36-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141164</p>
<p>TCO-PEG3-TCO is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG36-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141159</p>	<p><b>TCO-PEG4-amine</b></p> <p style="text-align: right;">Cat. No.: HY-141179</p>
<p>TCO-PEG4-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG4-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG4-biotin</b></p> <p style="text-align: right;">Cat. No.: HY-141183</p>	<p><b>TCO-PEG4-DBCO</b></p> <p style="text-align: right;">Cat. No.: HY-140310</p>
<p>TCO-PEG4-biotin is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg</p>	<p>TCO-PEG4-DBCO is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. TCO-PEG4-DBCO is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>










<p><b>TCO-PEG4-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141167</p>	<p><b>TCO-PEG4-TCO</b></p> <p style="text-align: right;">Cat. No.: HY-141188</p>
<p>TCO-PEG4-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. TCO-PEG4-NHS ester is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p>  <p><b>Purity:</b> 99.58%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG4-TCO is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG5-maleimide</b></p> <p style="text-align: right;">Cat. No.: HY-133475</p>	<p><b>TCO-PEG6-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141160</p>
<p>TCO-PEG5-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG6-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG6-amine</b></p> <p style="text-align: right;">Cat. No.: HY-141180</p>	<p><b>TCO-PEG6-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141168</p>
<p>TCO-PEG6-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG6-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141161</p>	<p><b>TCO-PEG8-amine</b></p> <p style="text-align: right;">Cat. No.: HY-141181</p>
<p>TCO-PEG8-acid is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG8-amine is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PEG8-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-141169</p>	<p><b>TCO-PEG8-TCO</b></p> <p style="text-align: right;">Cat. No.: HY-141189</p>
<p>TCO-PEG8-NHS ester is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 25 mg, 50 mg, 100 mg</p>	<p>TCO-PEG8-TCO is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>TCO-PEG8-TFP ester</b></p> <p>Cat. No.: HY-141173</p>	<p><b>TCO-PEG9-maleimide</b></p> <p>Cat. No.: HY-141184</p>
<p>TCO-PEG8-TFP ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>TCO-PEG9-maleimide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>TCO-PNB ester</b></p> <p>Cat. No.: HY-141175</p>	<p><b>tert-Butyl 11-aminoundecanoate</b></p> <p>Cat. No.: HY-130715</p>
<p>TCO-PNB ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>tert-Butyl 11-aminoundecanoate (compound 6b) is a PROTAC linker, which refers to the PEG composition. tert-Butyl 11-aminoundecanoate can be used in the synthesis of a series of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 100 mg</p>
<p><b>Tetra(cyanoethoxymethyl) methane</b></p> <p>Cat. No.: HY-124531</p>	<p><b>Tetraethyl butane-1,4-diylbis(phosphonate)</b></p> <p>Cat. No.: HY-140003</p>
<p>Tetra(cyanoethoxymethyl) methane is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Tetraethyl butane-1,4-diylbis(phosphonate) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tetraethyl butane-1,4-diylbis(phosphoramidate)</b></p> <p>Cat. No.: HY-141324</p>	<p><b>Tetraethyl decane-1,10-diylbis(phosphonate)</b></p> <p>Cat. No.: HY-140325</p>
<p>Tetraethyl butane-1,4-diylbis(phosphoramidate) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b> No Development Reported</p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Tetraethyl decane-1,10-diylbis(phosphonate) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tetraethyl heptane-1,7-diylbis(phosphonate)</b></p> <p>Cat. No.: HY-140323</p>	<p><b>Tetraethyl octane-1,8-diylbis(phosphonate)</b></p> <p>Cat. No.: HY-140324</p>
<p>Tetraethyl heptane-1,7-diylbis(phosphonate) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>	<p>Tetraethyl octane-1,8-diylbis(phosphonate) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%</p> <p><b>Clinical Data:</b></p> <p><b>Size:</b> 1 mg, 5 mg</p>
























<p><b>Tetrazine-Ph-NHCO-PEG4-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133462</p> <p>Tetrazine-Ph-NHCO-PEG4-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Tetrazine-Ph-NHCO-PEG6-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133461</p> <p>Tetrazine-Ph-NHCO-PEG6-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tetrazine-Ph-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-126908</p> <p>Tetrazine-Ph-NHS ester is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg</p>	<p><b>Tetrazine-Ph-PEG4-Ph-aldehyde</b></p> <p style="text-align: right;">Cat. No.: HY-133465</p> <p>Tetrazine-Ph-PEG4-Ph-aldehyde is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tetrazine-Ph-PEG5-NHS ester</b></p> <p style="text-align: right;">Cat. No.: HY-130551</p> <p>Tetrazine-Ph-PEG5-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Tetrazine-Ph-PEG5-Ph-tetrazine</b></p> <p style="text-align: right;">Cat. No.: HY-133463</p> <p>Tetrazine-Ph-PEG5-Ph-tetrazine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>TFP-PEG3-TFP</b></p> <p style="text-align: right;">Cat. No.: HY-W096112</p> <p>TFP-PEG3-TFP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>TGN-020</b></p> <p style="text-align: right;">Cat. No.: HY-W008574</p> <p>TGN-020 is a selective <b>Aquaporin 4 (AQP4)</b> inhibitor with an <math>IC_{50}</math> of 3.1 <math>\mu</math>M. TGN-020 is an alkyl chain-based <b>PROTAC linker</b> that can be used in the synthesis of PROTACs. TGN-020 alleviates edema and inhibits glial scar formation after spinal cord compression injury in rats.</p>  <p><b>Purity:</b> 98.03%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM <math>\times</math> 1 mL, 10 mg, 50 mg, 100 mg</p>
<p><b>Thalidomide-O-amido-PEG4-azide</b></p> <p style="text-align: right;">Cat. No.: HY-141011</p> <p>Thalidomide-O-amido-PEG4-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Thalidomide-O-amido-PEG4-propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-141013</p> <p>Thalidomide-O-amido-PEG4-propargyl is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Thiol-C10-amide-PEG8</b></p> <p style="text-align: right;">Cat. No.: HY-138533</p> <p>Thiol-C10-amide-PEG8 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Thiol-C2-PEG2-OH</b></p> <p style="text-align: right;">Cat. No.: HY-135085</p> <p>Thiol-C2-PEG2-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Thiol-C9-PEG4</b></p> <p style="text-align: right;">Cat. No.: HY-138527</p> <p>Thiol-C9-PEG4 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Thiol-C9-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138526</p> <p>Thiol-C9-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Thiol-C9-PEG5</b></p> <p style="text-align: right;">Cat. No.: HY-138529</p> <p>Thiol-C9-PEG5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Thiol-C9-PEG5-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138528</p> <p>Thiol-C9-PEG5-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Thiol-C9-PEG7</b></p> <p style="text-align: right;">Cat. No.: HY-138517</p> <p>Thiol-C9-PEG7 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Thiol-PEG-CH2COOH (MW 2000)</b></p> <p style="text-align: right;">Cat. No.: HY-140728</p> <p>Thiol-PEG-CH2COOH (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 2000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Thiol-PEG-CH2COOH (MW 3400)</b></p> <p style="text-align: right;">Cat. No.: HY-140729</p> <p>Thiol-PEG-CH2COOH (MW 3400) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 3400</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Thiol-PEG-CH2COOH (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-140730</p> <p>Thiol-PEG-CH2COOH (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p style="text-align: center;">MW 5000</p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg, 250 mg</p>

<p><b>Thiol-PEG-COOH (MW 5000)</b></p> <p style="text-align: right;">Cat. No.: HY-138310</p>	<p><b>Thiol-PEG10-alcohol</b> (HS-PEG10-OH)</p> <p style="text-align: right;">Cat. No.: HY-133295</p>
<p>Thiol-PEG-COOH (MW 5000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  <p>MW 5000</p> </div> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Thiol-PEG10-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Thiol-PEG12-acid</b></p> <p style="text-align: right;">Cat. No.: HY-141326</p>	<p><b>Thiol-PEG12-alcohol</b> (HS-PEG12-OH)</p> <p style="text-align: right;">Cat. No.: HY-133297</p>
<p>Thiol-PEG12-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> <b>Size:</b> 1 mg, 5 mg</p>	<p>Thiol-PEG12-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Thiol-PEG2-acid</b></p> <p style="text-align: right;">Cat. No.: HY-138524</p>	<p><b>Thiol-PEG2-t-butyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-132100</p>
<p>Thiol-PEG2-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Thiol-PEG2-t-butyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Thiol-PEG3-acetic acid</b></p> <p style="text-align: right;">Cat. No.: HY-W190729</p>	<p><b>Thiol-PEG3-acid</b></p> <p style="text-align: right;">Cat. No.: HY-140018</p>
<p>Thiol-PEG3-acetic acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Thiol-PEG3-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> ≥95.0% <b>Clinical Data:</b> <b>Size:</b> 50 mg, 100 mg, 500 mg</p>
<p><b>Thiol-PEG3-Boc</b> (Thiol-PEG3-t-butyl ester)</p> <p style="text-align: right;">Cat. No.: HY-141327</p>	<p><b>Thiol-PEG3-NHBoc</b></p> <p style="text-align: right;">Cat. No.: HY-138759</p>
<p>Thiol-PEG3-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Thiol-PEG3-NHBoc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <div style="text-align: center;">  </div> <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>













<p><b>Thiol-PEG8-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130542</p> <p>Thiol-PEG8-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>Thiol-PEG8-alcohol</b> (HS-PEG8-OH)</p> <p style="text-align: right;">Cat. No.: HY-133294</p> <p>Thiol-PEG8-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> ≥98.0%  <b>Clinical Data:</b>  <b>Size:</b> 100 mg</p>
<p><b>Thiol-PEG9-alcohol</b> (HS-PEG9-OH)</p> <p style="text-align: right;">Cat. No.: HY-133296</p> <p>Thiol-PEG9-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>THP-C1-PEG5</b></p> <p style="text-align: right;">Cat. No.: HY-138403</p> <p>THP-C1-PEG5 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-C4-PEG4</b></p> <p style="text-align: right;">Cat. No.: HY-138465</p> <p>THP-C4-PEG4 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>THP-CH3-ethyl propionate</b></p> <p style="text-align: right;">Cat. No.: HY-138534</p> <p>THP-CH3-ethyl propionate is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG1-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-W038771</p> <p>THP-PEG1-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>THP-PEG1-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138368</p> <p>THP-PEG1-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG1-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138531</p> <p>THP-PEG1-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p><b>THP-PEG10-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138347</p> <p>THP-PEG10-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>THP-PEG10-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138316</p>	<p><b>THP-PEG10-OH</b></p> <p style="text-align: right;">Cat. No.: HY-138413</p>
<p>THP-PEG10-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG10-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG10-THP</b></p> <p style="text-align: right;">Cat. No.: HY-134747</p>	<p><b>THP-PEG11-OH</b></p> <p style="text-align: right;">Cat. No.: HY-141227</p>
<p>THP-PEG10-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG11-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG11-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132014</p>	<p><b>THP-PEG12-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-132104</p>
<p>THP-PEG11-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG12-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG12-THP</b></p> <p style="text-align: right;">Cat. No.: HY-134699</p>	<p><b>THP-PEG13-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138478</p>
<p>THP-PEG12-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG13-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG13-OH</b></p> <p style="text-align: right;">Cat. No.: HY-138389</p>	<p><b>THP-PEG16-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-132058</p>
<p>THP-PEG13-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG16-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>THP-PEG16-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132077</p>	<p><b>THP-PEG2-Mal</b></p> <p style="text-align: right;">Cat. No.: HY-138404</p>
<p>THP-PEG16-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG2-Mal is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG2-methyl propionate</b></p> <p style="text-align: right;">Cat. No.: HY-138348</p>	<p><b>THP-PEG24-THP</b></p> <p style="text-align: right;">Cat. No.: HY-138329</p>
<p>THP-PEG2-methyl propionate is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG24-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG3-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130216</p>	<p><b>THP-PEG4-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138352</p>
<p>THP-PEG3-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG4-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG4-C1-OH</b></p> <p style="text-align: right;">Cat. No.: HY-134675</p>	<p><b>THP-PEG4-OH</b></p> <p style="text-align: right;">Cat. No.: HY-130230</p>
<p>THP-PEG4-C1-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG4-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG4-Pyrrolidine(N-Boc)-CH2OH</b></p> <p style="text-align: right;">Cat. No.: HY-130820</p>	<p><b>THP-PEG4-Pyrrolidine(N-Me)-CH2OH</b></p> <p style="text-align: right;">Cat. No.: HY-130821</p>
<p>THP-PEG4-Pyrrolidine(N-Boc)-CH2OH is a PEG-based PROTAC linker can be used in the synthesis of PROTAC K-Ras Degradier-1 (HY-129523).</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p>THP-PEG4-Pyrrolidine(N-Me)-CH2OH is a PEG-based PROTAC linker can be used in the synthesis of PROTAC K-Ras Degradier-1 (HY-129523).</p>  <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>



<p><b>THP-PEG5-OH</b></p> <p style="text-align: right;">Cat. No.: HY-126917</p>	<p><b>THP-PEG6-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138349</p>
<p>THP-PEG5-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG6-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG6-OH</b></p> <p style="text-align: right;">Cat. No.: HY-126918</p>	<p><b>THP-PEG7-alcohol</b></p> <p style="text-align: right;">Cat. No.: HY-132068</p>
<p>THP-PEG6-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. THP-PEG6-OH is also a non-cleavable 3 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG7-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG8-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-138340</p>	<p><b>THP-PEG8-OH</b></p> <p style="text-align: right;">Cat. No.: HY-126919</p>
<p>THP-PEG8-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG8-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG8-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132072</p>	<p><b>THP-PEG8-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-141228</p>
<p>THP-PEG8-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG8-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>THP-PEG9-OH</b></p> <p style="text-align: right;">Cat. No.: HY-138483</p>	<p><b>THP-PEG9-THP</b></p> <p style="text-align: right;">Cat. No.: HY-134698</p>
<p>THP-PEG9-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>THP-PEG9-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: center;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

**TLR4-IN-C34-C2-amide-C6-OH**


Cat. No.: HY-145245

TLR4-IN-C34-C2-amide-C6-OH is a linker that incorporates TLR4 inhibitor TLR4-IN-C34. TLR4-IN-C34 inhibits TLR4 in enterocytes and macrophages, and reduces systemic inflammation in mouse models of endotoxemia and necrotizing enterocolitis.

**Purity:** >98%

**Clinical Data:** No Development Reported

**Size:** 25 mg, 50 mg, 100 mg, 500 mg



**TLR4-IN-C34-C2-COOH**


Cat. No.: HY-W092043

TLR4-IN-C34-C2-COOH is a linker that incorporates TLR4 inhibitor TLR4-IN-C34. TLR4-IN-C34 inhibits TLR4 in enterocytes and macrophages, and reduces systemic inflammation in mouse models of endotoxemia and necrotizing enterocolitis.

**Purity:** >98%

**Clinical Data:** No Development Reported

**Size:** 25 mg, 50 mg, 100 mg, 500 mg, 1 g



**Tos-aminoxy-Boc-PEG4-Tos**


Cat. No.: HY-140380

Tos-aminoxy-Boc-PEG4-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.

**Purity:** >98%

**Clinical Data:** No Development Reported

**Size:** 1 mg, 5 mg



**Tos-O-C4-NH-Boc**


Cat. No.: HY-132004

Tos-O-C4-NH-Boc is an alkyl ether-based PROTAC linker that can be used in the synthesis of PROTACs, such as BSJ-03-204 (HY-136250).

**Purity:** 96.31%

**Clinical Data:** No Development Reported

**Size:** 500 mg



**Tos-PEG1-CH2-Boc**


Cat. No.: HY-130158

Tos-PEG1-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.

**Purity:** >98%

**Clinical Data:** No Development Reported

**Size:** 1 mg, 5 mg



**Tos-PEG1-O-CH2COOH**  
(PROTAC Linker 28)


Cat. No.: HY-125844

Tos-PEG1-O-CH2COOH (PROTAC Linker 28) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.

**Purity:** >98%

**Clinical Data:** No Development Reported

**Size:** 1 mg, 5 mg



**Tos-PEG12-Tos**


Cat. No.: HY-140371

Tos-PEG12-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.

**Purity:** >98%

**Clinical Data:** No Development Reported

**Size:** 1 mg, 5 mg



**Tos-PEG13-Boc**


Cat. No.: HY-140365

Tos-PEG13-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.

**Purity:** >98%

**Clinical Data:** No Development Reported

**Size:** 1 mg, 5 mg



**Tos-PEG14-OH**


Cat. No.: HY-134713

Tos-PEG14-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.

**Purity:** >98%

**Clinical Data:** No Development Reported

**Size:** 1 mg, 5 mg



**Tos-PEG2-Boc**


Cat. No.: HY-130534

Tos-PEG2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.

**Purity:** >98%











**Clinical Data:** No Development Reported

**Size:** 1 mg, 5 mg




<p><b>Tos-PEG2-C2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-133069</p>	<p><b>Tos-PEG2-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-130532</p>
<p>Tos-PEG2-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG2-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tos-PEG2-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130415</p>	<p><b>Tos-PEG2-NH-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-135798</p>
<p>Tos-PEG2-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG2-NH-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 98.23%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 50 mg, 100 mg</p>
<p><b>Tos-PEG2-NH2</b> (PROTAC Linker 27)</p> <p style="text-align: right;">Cat. No.: HY-125842</p>	<p><b>Tos-PEG2-O-Propargyl</b></p> <p style="text-align: right;">Cat. No.: HY-130162</p>
<p>Tos-PEG2-NH2 (PROTAC Linker 27) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 250 mg</p>	<p>Tos-PEG2-O-Propargyl is a PEG-based PROTAC linker that can be used in the synthesis of Thalidomide-O-PEG2-propargyl (HY-126458).</p>  <p><b>Purity:</b> 98.47%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 5 mg, 10 mg, 50 mg, 100 mg</p>
<p><b>Tos-PEG2-OH</b></p> <p style="text-align: right;">Cat. No.: HY-W090623</p>	<p><b>Tos-PEG2-THP</b></p> <p style="text-align: right;">Cat. No.: HY-132020</p>
<p>Tos-PEG2-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG2-THP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tos-PEG20-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140372</p>	<p><b>Tos-PEG21-Tos</b></p> <p style="text-align: right;">Cat. No.: HY-140373</p>
<p>Tos-PEG20-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG21-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Tos-PEG3</b></p> <p style="text-align: right;">Cat. No.: HY-23408</p>	<p><b>Tos-PEG3-C2-methyl ester</b></p> <p style="text-align: right;">Cat. No.: HY-140375</p>
<p>Tos-PEG3 is a PEG-based PROTAC linker can be used in the synthesis of PROTACs. Tos-PEG3 (structure 1) can be used for the synthesis of 3'-aminoxy oligonucleotides solid supports.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> 97.39%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>	<p>Tos-PEG3-C2-methyl ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tos-PEG3-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130674</p>	<p><b>Tos-PEG3-CH2COOtBu</b></p> <p style="text-align: right;">Cat. No.: HY-45018</p>
<p>Tos-PEG3-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG3-CH2COOtBu is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg</p>
<p><b>Tos-PEG3-NH-Boc</b> (PROTAC Linker 9)</p> <p style="text-align: right;">Cat. No.: HY-128805</p>	<p><b>Tos-PEG3-O-C1-CH3COO</b> (PROTAC Linker 6)</p> <p style="text-align: right;">Cat. No.: HY-128802</p>
<p>Tos-PEG3-NH-Boc (PROTAC Linker 9) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> 99.66%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 50 mg, 100 mg, 250 mg, 500 mg</p>	<p>Tos-PEG3-O-C1-CH3COO (PROTAC Linker 6) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tos-PEG4-acid</b></p> <p style="text-align: right;">Cat. No.: HY-130417</p>	<p><b>Tos-PEG4-CH2-Boc</b></p> <p style="text-align: right;">Cat. No.: HY-42620</p>
<p>Tos-PEG4-acid is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG4-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> ≥95.0%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 250 mg, 1 g</p>
<p><b>Tos-PEG4-CH2COOH</b></p> <p style="text-align: right;">Cat. No.: HY-130473</p>	<p><b>Tos-PEG4-NH-Boc</b> (PROTAC Linker 7)</p> <p style="text-align: right;">Cat. No.: HY-128803</p>
<p>Tos-PEG4-CH2COOH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG4-NH-Boc (PROTAC Linker 7) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p> <p style="text-align: right;"></p> <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>

<p><b>Tos-PEG4-t-butyl ester</b> (Tos-PEG4-Boc) <span style="float: right;">Cat. No.: HY-130422</span></p>	<p><b>Tos-PEG4-THP</b> <span style="float: right;">Cat. No.: HY-130819</span></p>
<p>Tos-PEG4-t-butyl ester (Tos-PEG4-Boc) is a <b>PROTAC linker</b>, which refers to the PEG composition. Tos-PEG4-t-butyl ester (Tos-PEG4-Boc) can be used in the synthesis of a series of PROTACs, such as BI-3663 (HY-111546).</p>  <p><b>Purity:</b> 97.44% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 10 mg, 25 mg, 50 mg, 100 mg</p>	<p>Tos-PEG4-THP is a PEG-based <b>PROTAC linker</b> can be used in the synthesis of PROTAC K-Ras Degradar-1 (HY-129523).</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 10 mM × 1 mL, 500 mg</p>
<p><b>Tos-PEG5-Boc</b> <span style="float: right;">Cat. No.: HY-130523</span></p>	<p><b>Tos-PEG5-C2-Boc</b> <span style="float: right;">Cat. No.: HY-133070</span></p>
<p>Tos-PEG5-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG5-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tos-PEG5-CH2COOtBu</b> <span style="float: right;">Cat. No.: HY-140366</span></p>	<p><b>Tos-PEG6-C2-Boc</b> <span style="float: right;">Cat. No.: HY-133071</span></p>
<p>Tos-PEG5-CH2COOtBu is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG6-C2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tos-PEG6-CH2-Boc</b> <span style="float: right;">Cat. No.: HY-130526</span></p>	<p><b>Tos-PEG6-OH</b> <span style="float: right;">Cat. No.: HY-140356</span></p>
<p>Tos-PEG6-CH2-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG6-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tos-PEG7-OH</b> <span style="float: right;">Cat. No.: HY-117028</span></p>	<p><b>Tos-PEG8-Tos</b> <span style="float: right;">Cat. No.: HY-140369</span></p>
<p>Tos-PEG7-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>	<p>Tos-PEG8-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98% <b>Clinical Data:</b> No Development Reported <b>Size:</b> 1 mg, 5 mg</p>

**Tos-PEG9** Cat. No.: HY-140357


Tos-PEG9 is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:**  
**Size:** 1 mg, 5 mg

**Tos-PEG9-Boc** Cat. No.: HY-125534


Tos-PEG9-Boc is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Tos-PEG9-Tos** Cat. No.: HY-140370


Tos-PEG9-Tos is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Tr-PEG2-OH** Cat. No.: HY-114995


Tr-PEG2-OH is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs. Tr-PEG2-OH is also a non-cleavable 2 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Tr-PEG4-OH** Cat. No.: HY-126883


Tr-PEG4-OH is a non-cleavable 4 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs).



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Tr-PEG5-OH** Cat. No.: HY-120845


Tr-PEG5-OH is a non-cleavable 5 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Tr-PEG5-OH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Tr-PEG8-OH** Cat. No.: HY-130165


Tr-PEG8-OH is a non-cleavable 8 unit PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Tr-PEG8-OH is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Tri(Amino-PEG3-amide)-amine** Cat. No.: HY-140249


Tri(Amino-PEG3-amide)-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** ≥98.0%  
**Clinical Data:** No Development Reported  
**Size:** 100 mg, 250 mg

**Tri(Amino-PEG3-amide)-amine TFA** Cat. No.: HY-140249A


Tri(Amino-PEG3-amide)-amine TFA is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Tri(Amino-PEG4-amide)-amine** Cat. No.: HY-140250

Tri(Amino-PEG4-amide)-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** ≥95.0%  
**Clinical Data:**  
**Size:** 100 mg, 250 mg

<p><b>Tri(Amino-PEG4-amide)-amine TFA</b></p> <p style="text-align: right;">Cat. No.: HY-140250A</p>	<p><b>Tri(Amino-PEG5-amide)-amine</b></p> <p style="text-align: right;">Cat. No.: HY-140251</p>
<p>Tri(Amino-PEG4-amide)-amine TFA is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>	<p>Tri(Amino-PEG5-amide)-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tri(Mal-PEG2-amide)-amine</b></p> <p style="text-align: right;">Cat. No.: HY-141008</p>	<p><b>Tri(t-butoxycarbonylethoxymethyl) ethanol</b></p> <p style="text-align: right;">Cat. No.: HY-113924</p>
<p>Tri(Mal-PEG2-amide)-amine is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Tri(t-butoxycarbonylethoxymethyl) ethanol is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 1 mg, 5 mg</p>
<p><b>Tri(TLR4-IN-C34-C2-amide-C3-amide-PEG1)-amide-C3-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-145255</p>	<p><b>Tri(TLR4-IN-C34-C2-amide-PEG1)-amide-C3-COOH</b></p> <p style="text-align: right;">Cat. No.: HY-145253</p>
<p>Tri(TLR4-IN-C34-C2-amide-C3-amide-PEG1)-amide-C3-COOH is a linker that incorporates TLR4 inhibitor TLR4-IN-C34. TLR4-IN-C34 inhibits TLR4 in enterocytes and macrophages, and reduces systemic inflammation in mouse models of endotoxemia and necrotizing enterocolitis.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg, 50 mg, 100 mg</p>	<p>Tri(TLR4-IN-C34-C2-amide-PEG1)-amide-C3-COOH is a linker that incorporates TLR4 inhibitor TLR4-IN-C34. TLR4-IN-C34 inhibits TLR4 in enterocytes and macrophages, and reduces systemic inflammation in mouse models of endotoxemia and necrotizing enterocolitis.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 5 mg, 10 mg, 50 mg, 100 mg</p>
<p><b>Tri-(PEG1-C2-acid)</b></p> <p style="text-align: right;">Cat. No.: HY-140541</p>	<p><b>Triethylene glycol (PROTAC Linker 25)</b></p> <p style="text-align: right;">Cat. No.: HY-W017440</p>
<p>Tri-(PEG1-C2-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> &gt;98%  <b>Clinical Data:</b>  <b>Size:</b> 1 mg, 5 mg</p>	<p>Triethylene glycol (PROTAC Linker 25) is a PEG-based PROTAC linker can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.88%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 10 mM × 1 mL, 100 mg</p>
<p><b>Triethylene glycol bis(p-toluenesulfonate)</b> (Bis-Tos-PEG3)</p> <p style="text-align: right;">Cat. No.: HY-W013731</p>	<p><b>Triethylene glycol monobenzyl ether</b></p> <p style="text-align: right;">Cat. No.: HY-W044459</p>
<p>Triethylene glycol bis(p-toluenesulfonate) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.42%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 500 mg</p>	<p>Triethylene glycol monobenzyl ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.</p>  <p><b>Purity:</b> 99.25%  <b>Clinical Data:</b> No Development Reported  <b>Size:</b> 100 mg</p>

**Triethylene glycol monodecyl ether**

Cat. No.: HY-W190971

Triethylene glycol monodecyl ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Triethylene glycol monododecyl ether**

Cat. No.: HY-W142506

Triethylene glycol monododecyl ether is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Tris[[2-(tert-butoxycarbonyl)ethoxy]methyl]methylamine**

Cat. No.: HY-21577

Tris[[2-(tert-butoxycarbonyl)ethoxy]methyl]methylamine is a cleavable PEG ADC linker used in the synthesis of antibody-drug conjugates (ADCs). Amino-Tri-(t-butoxycarbonylethoxymethyl)-methane is also a PEG/Alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** ≥97.0%  
**Clinical Data:** No Development Reported  
**Size:** 50 mg

**Trityl-PEG10-azide**

Cat. No.: HY-140861

Trityl-PEG10-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Trityl-PEG8-azide**

Cat. No.: HY-140860

Trityl-PEG8-azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**Trt-PEG4-C2-acid hydrate**

Cat. No.: HY-W096111A

Trt-PEG4-C2-acid (hydrate) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**UV Cleavable Biotin-PEG2-Azide**

Cat. No.: HY-140920

UV Cleavable Biotin-PEG2-Azide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** ≥95.0%  
**Clinical Data:** No Development Reported  
**Size:** 25 mg, 100 mg

**VES-POFP**

Cat. No.: HY-134751

VES-POFP is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**WSPC Biotin-PEG3-DBCO**

Cat. No.: HY-140137

WSPC Biotin-PEG3-DBCO is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.




**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg

**[10-(Diethoxy-phosphoryl)-decyl]-phosphonic acid**

Cat. No.: HY-140327

[10-(Diethoxy-phosphoryl)-decyl]-phosphonic acid is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%  
**Clinical Data:** No Development Reported  
**Size:** 1 mg, 5 mg




### $\alpha$ -Lipoic acid-NHS

(DL- $\alpha$ -Lipoic acid-NHS)

Cat. No.: HY-141336

$\alpha$ -Lipoic acid-NHS is an alkyl/ether-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%


**Clinical Data:**

**Size:** 1 mg, 5 mg

### $\beta$ -Estradiol-6-one 6-(O-carboxymethyloxime)

Cat. No.: HY-133407

$\beta$ -Estradiol-6-one 6-(O-carboxymethyloxime) is an alkyl chain-based PROTAC linker that can be used in the synthesis of PROTACs.



**Purity:** >98%

**Clinical Data:** No Development Reported

**Size:** 1 mg, 5 mg